



GOVERNMENT COLLEGE FOR WOMEN (AUTONOMOUS)

KUMBAKONAM – 612 001

Affiliated to Bharathidasan University

DST - CURIE Sponsored Institution

IV Cycle of Accreditation

 **0435 – 2401391**

 **principal@gwk.ac.in**



CRITERION III – RESEARCH, INNOVATIONS AND EXTENSION

FRONT PAGE OF PUBLISHED PAPERS

2022-2023

Article

Effects of Dietary Blend of Algae Extract Supplementation on Growth, Biochemical, Haemato-Immunological Response, and Immune Gene Expression in *Labeo rohita* with *Aeromonas hydrophila* Post-Challenges

Govindharajan Sattanathan ^{1,†}, Wen-Chao Liu ^{2,†}, Swaminathan Padmapriya ³, Karthika Pushparaj ⁴, Shanmugam Sureshkumar ⁵, Jang-Won Lee ⁶, Balamuralikrishnan Balasubramanian ^{7,*} and In Ho Kim ^{5,*}

¹ Department of Zoology, St. Joseph University, Chumukedima 797115, Nagaland, India

² Department of Animal Science, College of Coastal Agricultural Sciences, Guangdong Ocean University, Zhanjiang 524088, China

³ Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

⁴ Department of Zoology, School of Biosciences, Avinashilingam Institute for Home Science and Higher Education for Women, Coimbatore 641043, Tamil Nadu, India

⁵ Department of Animal Resource and Science, Dankook University, Cheonan-si, Chungnam 31116, Republic of Korea

⁶ Department of Integrative Biological Sciences and Industry, Sejong University, Seoul 05006, Republic of Korea

⁷ Department of Food Science and Biotechnology, College of Life Science, Sejong University, Seoul 05006, Republic of Korea

* Correspondence: bala.m.k@sejong.ac.kr or geneticsmurali@gmail.com (B.B.); inhokim@dankook.ac.kr (I.H.K)

† Equally contributed as first author.



Citation: Sattanathan, G.; Liu, W.-C.; Padmapriya, S.; Pushparaj, K.; Sureshkumar, S.; Lee, J.-W.; Balasubramanian, B.; Kim, I.H. Effects of Dietary Blend of Algae Extract Supplementation on Growth, Biochemical, Haemato-Immunological Response, and Immune Gene Expression in *Labeo rohita* with *Aeromonas hydrophila* Post-Challenges. *Fishes* 2023, 8, 7. <https://doi.org/10.3390/fishes8010007>

Academic Editor: Jong Chen

Received: 28 October 2022

Revised: 16 December 2022

Accepted: 20 December 2022

Published: 22 December 2022



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: In this study, the effects of a mixed algal blend (*Chlorella vulgaris*, *Euglena viridis*, and *Spirulina platensis*) at different levels were evaluated on growth, hematological immune responses, and expression of immune genes in *Labeo rohita* against post-challenges of *Aeromonas hydrophila*. Fish samples were fed a diet containing different levels of mixed blend algal (0, 0.01, 0.02, 0.04 and 0.08% of basal diet). At the end of the feeding period, the fish were challenged with *A. hydrophila* and fish mortality was recorded over a 14-days period. To evaluate the serum biochemical (albumin, globulin), hematological parameters (Hb, RBC and WBC) and immune parameters (neutrophil activity, lysozyme activity, myeloperoxidase activity, antiprotease activity, ceruloplasmin activity, and bactericidal activity), as well as the expression of immune genes (NKEF-B, Lysozyme C and G, TNF α , TLR22, β 2M, and β -actin), fish were sampled on Day 7, 14, 21 and 28. Fish were challenged with virulent *A. hydrophila* 30 days post-feeding and mortalities were recorded over 30 days post-infection. Results demonstrate that fish fed with a mixed algal blend showed that total body weight gain, specific growth rate, total serum protein, globulin, total hemoglobin content, white blood cells, neutrophil, lysozyme, bactericidal, myeloperoxidase, and antiprotease activity in dietary algae blended application was higher than in the control ($p < 0.05$). According to the results, relative expression of target genes showed significant increases of 0.02 to 0.04% in the treatment group compared to the control group ($p < 0.05$). At the end of the 30-day exposure to *A. hydrophila*, the fish that received the mixed algal blend had a significantly higher rate of survival than the control group, with the highest survival rate recorded in the 0.02% mixed algal blend ($p < 0.05$). According to the effective results of the mixed algal blend on stimulating the immune system and increasing fish resistance to *A. hydrophila*, it is recommended to use 0.02 to 0.04% of this mixed algal blend in rohu, *L. rohita* diets.

Keywords: mixed algae; rohu; *Aeromonas hydrophila*; immune gene expression



TOXIC EFFECT OF LAMBDA CYHALOTHRIN AND PRETILACHLOR ON BIOMASS AND HISTOLOGY OF TESTIS OF *Lampito mauritii* (KINBERG)

G. INDHUMATHI ¹*, V. KAVITHA ¹** AND R. ANANDHAN ¹#

¹ Department of Zoology, Annamalai University, Annamalai Nagar - 608 002, Tamil Nadu, India.

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.56557/U.PJZ/2022/v43i173157

Editor(s):

(1) Dr. Juan Carlos Trilano, University of Buenos Aires, Argentina.

Reviewer(s):

(1) Sarangdha Navren, University of Malaya, Malaysia.

(2) Cenk Kilig, Türkiye.

Received: 05 July 2022

Accepted: 09 September 2022

Published: 17 September 2022

Original Research Article

ABSTRACT

Lambda cyhalothrin and pretilachlor is widely used to control agricultural pest but also affect non target organisms like earthworms. The present laboratory study was conducted to determine the effect of Lambda cyhalothrin and Pretilachlor on biomass and histology of testis of *Lampito mauritii*. Lower and higher sub-lethal concentrations of Lambda cyhalothrin (T1 and T2) and Pretilachlor (T3 and T4) was mixed with soil substrate. Biomass was observed once in 10 days up to 90 days. 10 non clitellate *L.mauritii* was introduced into each treatment. Soil substrate without lambda cyhalothrin and pretilachlor served as control. The present study was revealed that lambda cyhalothrin and pretilachlor was highly affected the growth of *L. mauritii*. For histopathology study, four adult *L. mauritii* were introduced in to C, T1, T2, T3 and T4. After 5th, 15th and 30th day, testis was dissected out. The results reported that histology of testis was highly damaged on 5th and 15th day of experiment than 30th day.

Keywords: Earthworm; *Lampito mauritii*; lambda cyhalothrin; pretilachlor; biomass; testis.

1. INTRODUCTION

Our nation father GandhiG told that agriculture is a backbone of India. The growing human population is expected to be 9.1 billion by 2050 (UN [1]). It needs more food production. So, chemical fertilizers and

pesticides are used in the agricultural field to avoid increasing demand for food which leads to soil contamination. India is the largest manufacturer of pesticides. More than 128 pesticides are registered in India Laxmi [2]. They are not only destroy pest and boostup plant growth also affect farmer's friend as

*Research Scholar;

#Assistant Professor;

**Corresponding author: Email: kavitha@kslso@gmail.com;



Synthesis and characterization of *Vitis vinifera* exocarp-mediated ZnO nanoparticles: An evaluation of biological potential and ecotoxicity

Kalyanaraman Nivetha^a, Viswanathan Vinotha^a, Mohammed F. Albeshr^b, Shahid Mahboob^b, Irfan Manzoor^c, Marimuthu Govindarajan^{d,e}, Baskaralingam Vaseeharan^{a,*}

^a Nanobioscience and Nanopharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Department of Biology, Indiana University Bloomington, Indiana, USA

^d Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Green synthesis
Nanoparticles
Anticancer activity
Antibacterial activity
Acute toxicity
Bio-monitoring

ABSTRACT

The popularity of plant-based nanoparticle synthesis has increased over the years due to the quantity of bioactive compounds. It is economical and environmentally friendly, which is an added feature. In the present study, the polyphenol-rich exocarp extract of *Vitis vinifera* was used to synthesize zinc oxide nanoparticles (Vv-ZnO NPs) and evaluate their potential usage in various biological applications. UV-Vis spectrophotometer observations of Vv-ZnO NPs revealed an SPR peak at 355 nm. FTIR analysis confirmed the presence of *V. vinifera* phytochemicals in the produced ZnO NPs. The electron microscope (SEM, TEM) revealed that the Vv-ZnO NPs had a conical shape and varied in size between 30 and 73.5 nm. The current investigation indicated that Vv-ZnO NPs are a more effective antioxidant than grape peel extract when exposed to DPPH solution, with an 87% inhibition rate at 100 µg/ml. At 200 µg/ml concentration, Vv-ZnO NPs demonstrated good antibacterial activity against *Staphylococcus aureus* (6 mm) and *Pseudomonas aeruginosa* (4 mm). The total destruction of biofilm architecture by Vv-ZnO NPs at high concentrations (100 µg/ml) was clearly seen in light microscopic images. The IC₅₀ value for its anticancer activity against the MDA-MB-231 breast cancer cell line was 12.90 µg/ml. The nanoparticles demonstrated acceptable larvicidal effectiveness against the Zika virus vector *Aedes aegypti* (LC₅₀-80.27-LC₉₀-136.39 µg/ml). Vv-ZnO NPs exhibited decreased ecotoxicity against non-target bioindicators such as *Daphnia similis*, *Artemia salina*, and *Perionyx excavatus*. Overall our studies revealed the eco-friendly synthesis of ecologically sound Vv-ZnO nanoparticles for diverse biological applications.

1. Introduction

The increasing urgency of developing eco-friendly technologies has helped the emerging science of nanotechnology gain ground in recent years. Nanomaterials are often manufactured via physical, chemical, and mechanical processes that need significant amounts of energy. These techniques are often risky since they call for high pressure, varying temperatures, and toxic mediators [1–4]. The detrimental impacts of such procedures on the environment and living organisms have prompted researchers to develop more cost-effective, biocompatible, and eco-conscious strategies [5]. Applications like semiconductors, catalysts, surfactants, and so on have been documented using individual

nanoparticles, bimetallic nanoparticles, and polymer-based nanocomposites [6–11].

Biopolymers produced from different types of life are used as capping agents in the environmentally friendly production of nanoparticles. Researchers have successfully created metal and metal oxide nanoparticles using plant-based biopolymers [12–14]. The plants are readily available and pose no health risks to those working with them. Zinc oxide nanoparticles (ZnO NPs) are a kind of metal oxide that has been shown to be both eco-friendly and biocompatible [15,16]. It has been found that biosynthesized ZnO NPs treat cancer and wounds and have antibacterial, antifungal, anti-inflammatory, and antidiabetic properties [17–20]. ZnO NPs are also employed as a packaging material for the

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jddst.2022.103846>

Received 17 August 2022; Received in revised form 21 September 2022; Accepted 27 September 2022


Available online 30 September 2022

1773-2247/© 2022 Elsevier B.V. All rights reserved.



Contents lists available at ScienceDirect

Journal of Trace Elements in Medicine and Biology

journal homepage: www.elsevier.com/locate/jtemb

The dietary supplementation of zinc oxide and selenium nanoparticles enhance the immune response in freshwater fish *Oreochromis mossambicus* against aquatic pathogen *Aeromonas hydrophila*

Mariappan Yazhini Prabha^a, Narayanan Gopi^a, Shahid Mahboob^b, Khalid A. Al-Ghanim^b, Fahad Al-Misned^b, Zubair Ahmed^b, Mian Nadeem Riaz^c, Jeyachandran Sivakamavalli^d, Marimuthu Govindarajan^{e,f}, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Texas A&M University, College Station, TX, United States

^d Department of Fisheries, Chonnam National University, Yeosu, South Korea

^e Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002 Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Immunostimulant
M. koenigii berry
 Nanoparticles
 Supplemented diet
A. hydrophila
O. mossambicus

ABSTRACT

Background: Green nanoparticles are subjected as an immunostimulant against bacterial pathogens.

Methods: *Murraya koenigii* berry extract-based synthesized zinc oxide nanoparticles (Mb-ZnO NPs) and selenium nanoparticles (Mb-Se NPs) were relatively analyzed for immunostimulation in serum and mucus fish *Oreochromis mossambicus* against *Aeromonas hydrophila* infections. Initial minimum inhibitory concentration (MIC) was determined for both Mb-ZnO NPs and Mb-Se NPs followed by specific growth rate (SGR), antioxidant level (Superoxide dismutase activity (SOD), Catalase activity (CA), and Glutathione peroxidase activity (GPx)), and immune parameters Myeloperoxidase activity (MPO), Respiratory burst activity (RBA), Lysozyme activity (LYZ), Alkaline phosphatase activity (ALP), Serum antiprotease activity and Natural complement activity (NAC).

Results: The potential bacterial inhibition property of Mb-ZnO NPs and Mb-Se NPs exhibited the most negligible concentration of 25 and 15 $\mu\text{g mL}^{-1}$, respectively, against *A. hydrophila*. In addition, Mb-ZnO NPs and Mb-Se NPs exhibited 70–80 % and 90–95 % diminished biofilm activity at 50 $\mu\text{g mL}^{-1}$ that was viewed under an inverted research microscope and confocal laser scanning microscopy (CLSM). Protein leakage and nucleic acid leakage assay quantified oozed out protein and nucleic acid from *A. hydrophila* that confirms Mb-Se NPs exhibited vigorous antibacterial activity than Mb-ZnO NPs at tested concentrations. *Oreochromis mossambicus* fed with Mb-ZnO NPs and Mb-Se NPs supplemented diet at different concentrations (0.5 mg/kg, 1 mg/kg and 2 mg/kg) improved SGR along with a rise in the immune response of those fishes against *A. hydrophila* infection. Serum and mucus of fish fed with Mb-Se NPs supplemented diet exhibited a significant rise in antioxidant level SOD, CA and GPx at a dosage of 2 mg/kg. Likewise, lipid peroxidation assay detected significantly diminished oxidative stress in the serum and mucus of fish fed with Mb-Se NPs supplemented diet (2 mg/kg). Enhanced immune parameters in serum and mucus of fish fed with Mb-Se NPs supplemented diet determined by MPO, RBA, LYZ, ALP, Serum antiprotease activity and NAC.

Conclusion: Thus *O. mossambicus* fed with Mb-Se NPs supplemented diet was less prone to become infected by aquatic pathogen *A. hydrophila* established by challenge study. On the whole, Mb-Se NPs supplemented diet ensured the rise in antioxidant response that boosts the immune responses and reduces the chance of getting infected against *A. hydrophila* infections.

* Corresponding author.


E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jtemb.2021.126878>

Received 9 July 2021; Received in revised form 31 August 2021; Accepted 12 October 2021

Available online 14 October 2021

0946-672X/© 2021 Elsevier GmbH. All rights reserved.



Characterization of secondary metabolites from Lamiaceae plant leaf essential oil: A novel perspective to combat medical and agricultural pests

Kuppusamy Elumalai^a, Kaliyamoorthy Krishnappa^b, Jeganathan Pandiyan^b, Naiyf S. Alharbi^c, Shine Kadaikunnan^c, Jamal M. Khaled^c, Donald R. Barnard^d, Natesan Vijayakumar^e, Marimuthu Govindarajan^{e,g,*}

^a Department of Advanced Zoology & Biotechnology, Government Arts College (Autonomous), Chennai, 600035, Tamil Nadu, India

^b Department of Zoology and Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai, 609305, India

^c Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^d United States Department of Agriculture, Agricultural Research Service, P. O Box 141498, Gainesville, FL, 32614-1498, USA

^e Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalainagar, 608002, Tamil Nadu, India

^f Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^g Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Aedes vittatus

Anopheles maculatus

Spodoptera litura

Helicoverpa armigera

Insecticidal activity

ABSTRACT

The insecticidal bio-efficacy of Lamiaceae essential oils (LEOs) and its major compounds (MCs) were tested against mosquitoes (*Aedes vittatus* and *Anopheles maculatus*) and agricultural pests (*Spodoptera litura* and *Helicoverpa armigera*). The naturally gained LEOs were confirmed through GC-MS analysis. The larval and pupal toxicity of LEOs and their major compounds LC₅₀/LC₉₀ value was calculated by standard protocol. A total of 54, 47 and 31 compounds were identified from *M. officinalis*, *L. latifolia* and *O. majorana*, respectively. The MCs of LEOs: (*M. officinalis*) 2,6-Octadienal, 3,7-dimethyl-, (Z)- (29.4354%) and Caryophyllene (20.2749%); (*L. latifolia*) 1,6-Octadien-3-ol, 3,7-dimethyl- (34.4992%) and 1-Cyclohexyl-2-buten-1-ol (c,t) (38.0339%); (*O. majorana*) Eucalyptol (57.1278%) and p-menth-1-en-8-ol (17.9683%). The LEOs and their MCs were tested individually against larvae and pupae of *Ae. vittatus*, *An. maculatus*, *S. litura* and *H. armigera*, which produced maximum toxic effects. Predominantly, the MCs were safer to the non-target organisms and target-specific insecticidal properties against selected insect pests. On the whole, our findings highlight that LEOs and their MCs represent promising eco-friendly and target specific insecticidal activities.

1. Introduction

The nuisance of growing agricultural and medical insect pest populations is exceedingly threatening to the human community in different ways. Globally, above 30% of economic losses in many cultivated crops are severely damaged by agricultural insect pests [1]. Human vector mosquitoes are well-known for transmitting plenty of frightful human diseases and creating negative impacts on human health, therefore which have been declared as “Public Enemy” [2]. Mosquito biting not only transmits the disease, consequently causing itching, irritation, and allergic skin response [3]. Mainly, tropical and subtropical zones are preferable hotspots for the occurrence of high levels of illness [4]. In addition, globally *Aedes* species are the predominant vector of many diseases as a shred of single evidence. Around 390 million people are

highly ripped by dengue infection in all continents [5]. In order to control the mosquitoes, people frequently use plenty of unadvisable synthetic insecticides, which are non-selective and cause drastic negative impacts on the environment/diverse eco-system [6]. According to the WHO [7], worldwide, nearly 3 million deaths and around 500 million infections are mainly raised by the only key vector of *Anopheles* species that remains endemic in above 100 developing tropical zones.

Spodoptera litura is a leading polyphagous Lepidopteron pest that drastically consumes all kinds of green flora and causes active infestation, consequently declining yield and significant profitable losses [8]. It exploits above 200 host flora and belongs to nearly 30 economically substantial vegetations families [9]. It has a wide range of hosts and almost 40 species are known from India only. Recently in southern India, it was noticed on brinjal, which is not an actual host plant of this pest

* Corresponding author. Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India.



E-mail address: drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.pmpp.2021.101752>

Received 2 October 2021; Received in revised form 24 October 2021; Accepted 9 November 2021

Available online 15 November 2021

0885-5765/© 2021 Elsevier Ltd. All rights reserved.



β -1,3-Glucan binding protein-based silver nanoparticles enhance the wound healing potential and disease resistance in *Oreochromis mossambicus* against *Aeromonas hydrophilla*

Arokiadhas Iswarya^a, Mahalingam Anjugam^a, Narayanan Gopi^a, Sathappan Shanthi^b, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Mohammed S. Alharbi^f, Jeyachandran Sivakamavalli^g, Baskaralingam Vaseeharan^{a,*}

^a Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th Floor, Burma Colony, Karaikudi – 630004, Tamil Nadu, India

^b Department of Zoology, University of Madras, Guindy Campus, Chennai-600 025, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar-608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^f Microbiology Department, King Khaled General Hospital, Hafer Albatin, Saudi Arabia

^g Department of Fisheries, Chonnam National University, Yeosu, South Korea

ARTICLE INFO

Keywords:
Skin lesion
Nanotechnology
Wound healing
Immunostimulant
Toxicity

ABSTRACT

Here we attempted to synthesize β -1,3-glucan binding protein-based silver nanoparticles ($Ph\beta$ -GBP-AgNPs) and evaluate its wound healing and disease resistance prompting ability in *Oreochromis mossambicus*. Using a column chromatography technique, an immune molecule, $Ph\beta$ -GBP was purified from the haemolymph of rice field crab, *Paratelphusa hydrodromus*. $Ph\beta$ -GBP-AgNPs were synthesized and described through SDS-PAGE, UV-vis spectroscopy, HR-TEM, XRD and FTIR analysis. HR-TEM revealed that the synthesized $Ph\beta$ -GBP-AgNPs were spherical with a 20–40 nm size range and the particles were not aggregated. Wound and infection were experimentally generated in *O. mossambicus* and treated with $Ph\beta$ -GBP, chem-AgNPs and $Ph\beta$ -GBP-AgNPs for 20 days. The immune parameters (peroxidase, lysozyme and protease) and antioxidant enzymes (SOD and catalase) were examined in the serum of experimental fish. $Ph\beta$ -GBP-AgNPs elevated the immune and antioxidant enzymes during the healing process and enhanced the wound healing percentage in fish than $Ph\beta$ -GBP and chem-AgNPs. The immune parameters and antioxidant enzymes were declined in the serum of fish (treated with $Ph\beta$ -GBP-AgNPs) after the mid-period of wound healing. Compared to others, relative percentage survival was increased in experimentally wounded and infected fish treated with $Ph\beta$ -GBP-AgNPs against *Aeromonas hydrophilla*. Moreover, $Ph\beta$ -GBP-AgNPs exhibited less toxicity towards *Artemia salina* than chem-AgNPs during 24 h exposure period. As a result, $Ph\beta$ -GBP-AgNPs may act as an alternative to commercial antibiotics and be considered an effective immunostimulant in treating skin lesions in intensive farming.

1. Introduction

Aquaculture is the fastest expanding food sector that supplies protein-rich food to meet the protein requirement for the growing population. Increased demand for protein leads to overfishing from wild sources results in unbalanced fish species [1]. Aquaculture introduces extensive and intensive fish farming to maintain the fish diversity in the wild from underbalanced conditions and meet the protein demand.

Intensive fish farming is highly practiced to increase fish production in a controlled area with regular aeration, water and food supply [2]. However, stress, disease, skin lesions are more prevalent in an intensive type of fish farming due to many fish populations [3]. Fish farmed in the intensive culture was more prone to skin damage, which may cause due to handling, transportation and infections [4,5]. Skin acts as a primary barrier against pathogens and its damage results in secondary infections caused by microorganisms. Infected or injured fish affect the wealth of

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.micpath.2021.105360>

Received 9 July 2021; Received in revised form 10 December 2021; Accepted 12 December 2021

Available online 15 December 2021

0882-4010/© 2021 Elsevier Ltd. All rights reserved.

Research Article

Balachandran Kartha, Kalaivani Thanikachalam, Natesan Vijayakumar*, Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, Kasi Gopinath, and **Marimuthu Govindarajan***

Synthesis and characterization of Ce-doped TiO₂ nanoparticles and their enhanced anticancer activity in Y79 retinoblastoma cancer cells

<https://doi.org/10.1515/gps-2022-0011>

received August 26, 2021; accepted December 16, 2021

Abstract: Rare earth metal cerium-doped titania nanoparticles (titanium dioxide [TiO₂]) were produced utilizing a low-cost and straightforward sol–gel technique, and its enhanced photodynamic anticancer activity was tested on Y79 retinoblastoma cancer cells. The structural, optical, morphological, anticancer activity, and cytotoxicity of pure and cerium-doped TiO₂ (Ce-doped TiO₂) were investigated. In X-ray diffraction (XRD) measurements, apparent doping of cerium in TiO₂ was detected, with reported anatase patterns shifting toward a lower angle in the anatase structure. Raman spectra verify the presence of cerium doping in TiO₂ by revealing greater wave number shifting. The scanning electron microscope (SEM) and transmission electron microscope (TEM) analysis showed that the synthesized TiO₂ and Ce-doped TiO₂ nearly spherical. TiO₂ and Ce-doped TiO₂ were studied for their photodynamic anticancer activities, and the results suggest that cerium doping in TiO₂ improves anticancer activity.

Keywords: cerium, nanocomposites, TiO₂, TEM, cytotoxicity

1 Introduction

Due to their innovative biological applications, semiconductor nanostructures with diameters smaller than 100 nm have emerged as nanobiomaterials [1,2]. In recent years, a wide range of nanocomposite semiconductor materials has been created to increase the photocatalytic activity efficiency [3,4]. By covering metal or semiconductors nanoclusters with some other layer of appropriate materials, the functional characteristics of such materials can be substantially enhanced [5,6].

Titanium dioxide (TiO₂) has been extensively used and demonstrated as a critical perspective photosensitizer [7–9], photostability, low cost, and nontoxicity. According to a recent study, the particle size of TiO₂ has a significant impact on its photocatalytic activity [10]. The reduction in particle size suggests an increase in surface area and a high redox potential, resulting in a strong photocatalytic activity.

Several methods, including doping, surface modification with metal particles, and particle size reduction to the nanoscale, have been proposed to increase the effectiveness of photocatalytic reactions utilizing TiO₂ when exposed to visible light [11]. Chemically modified TiO₂ nanoparticles are used in several environmental applications due to their self-cleaning properties [12,13]. Because of their nontoxicity, excellent optical absorption, cheap cost, and good chemical stability, metal and metal oxide nanoparticles have been widely investigated [14–19]. Among them, the medical applications of TiO₂ are undeniably promising, with the potential to significantly enhance health care, notably cancer therapy. Doping of Ce in TiO₂ enhances the photocatalysis and photodegradation of TiO₂ [20].

Sol–gel, direct aqueous solution depositions, ultrasonic spray pyrolysis, and sputtering are some of the

* **Corresponding author: Natesan Vijayakumar**, Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalainagar-608002, Tamil Nadu, India, e-mail: nvkbiochem@yahoo.co.in

* **Corresponding author: Marimuthu Govindarajan**, Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar-608002, Tamil Nadu, India; Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

Balachandran Kartha: Department of Chemistry, Vivekanandha College of Engineering for Women, Namakkal 637205, Tamil Nadu, India

Kalaivani Thanikachalam: Department of Chemistry, Jairams College of Arts and Science, Karur, Tamil Nadu, India

Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled: Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

Kasi Gopinath: Department of Packaging, Yonsei University, 1 Yonsei-dae-gil, Wonju-si, Gangwon-do 26493, South Korea

Research Article

Kuppusamy Elumalai, M. R. Kavipriya*, A. Lakshmi Prabha, Kaliyamoorthy Krishnappa, Jeganathan Pandiyan, Marcello Nicoletti, Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, and **Marimuthu Govindarajan***

Green synthesis of silver nanoparticles using *Atalantia monophylla*: A potential eco-friendly agent for controlling blood-sucking vectors

<https://doi.org/10.1515/gps-2022-0078>

received March 02, 2022; accepted August 22, 2022

Abstract: Developing floral-based replacement molecules might manage blood-sucking vectors in an eco-friendly way. *Atalantia monophylla* (*Am*) aqueous leaf extract (ALE) and silver nanoparticles (AgNPs) were evaluated against mosquitoes (*Aedes vittatus*, *Anopheles subpictus*, and *Culex vishnui*) and ticks (*Haemaphysalis bispinosa*, *Rhipicephalus microplus*, and *R. sanguineus*) at different concentrations. Phytochemical screening and AgNPs' synthesis were performed on ALE of *A. monophylla*. UV-visible spectroscopy, Fourier-transform infrared (FTIR) spectroscopy, scanning electron microscope, and transmission electron microscope were used to examine the synthesized *Am*-AgNPs. *A. monophylla*'s ALE included alkaloids, flavonoids, saponins, tannins, triterpenes,

coumarins, anthraquinones, and phenolics. *Am*-AgNPs had a higher LC_{50} (22.19, 23.92, 26.09, 40.25, 51.87, and 60.53 $\mu\text{g}\cdot\text{mL}^{-1}$, respectively) than leaf aqueous extract (LAE) against *Ae. vittatus*, *An. subpictus*, *Cx. vishnui*, *H. bispinosa*, *R. microplus*, and *R. sanguineus* larvae. *A. monophylla* ALE and *Am*-AgNPs' bio-toxicity was investigated against aquatic and terrestrial non-target species (*Acilius sulcatus*, *Anisops bouvieri*, *Araneus mitificus*, and *Cyrtophora moluccensis*) with LC_{50} values ranging from 2,094.5 to 10,532.8 $\mu\text{g}\cdot\text{mL}^{-1}$, respectively. *A. monophylla* ALE and *Am*-AgNPs had little negative impacts on the chosen non-target fauna. Environmental protection is important nowadays. Green AgNPs are low-cost, readily accessible, environmentally safe, and effective pesticides. *Am*-AgNPs are effective alternative insecticides, requiring a considerable study on this plant to control blood-sucking vectors for worldwide human/animal health importance.

Keywords: greener nanoparticles, blood-sucking vectors, larval toxicity, environmental safety, non-target fauna

* Corresponding author: M. R. Kavipriya, Department of Botany, School of Life Sciences, Bharathidasan University, Tiruchirappalli, 620024, Tamil Nadu, India, e-mail: kpresearch2126@gmail.com

* Corresponding author: Marimuthu Govindarajan, Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar-608 002, Tamil Nadu, India; Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

Kuppusamy Elumalai: Department of Advanced Zoology and Biotechnology, Government Arts College (Autonomous), Chennai, 600035, Tamil Nadu, India

A. Lakshmi Prabha: Department of Botany, School of Life Sciences, Bharathidasan University, Tiruchirappalli, 620024, Tamil Nadu, India

Kaliyamoorthy Krishnappa, Jeganathan Pandiyan: Department of Zoology and Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai, 609305, Tamil Nadu, India

Marcello Nicoletti: Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled: Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

1 Introduction

Blood-sucking vectors (BSVs) generating an abundance of newly developing illnesses substantially affect livestock and public health [1]. As a vector, ticks disperse pathogens responsible for causing cardinal diseases among cattle and human beings [2]. Around 900 tick species were recently identified [3]. The life-threatening pathogenic viruses are conveyed to animals by infected ticks' bites, especially in the family Ixodidae [4]. Ticks are notorious vectors for around 38 pathogenic viral species transmitted in the animal kingdom [5]. Worldwide, US\$ 7 billion is lost annually, and nearly 80% of farm animals are in high-risk because of ticks and tick-borne diseases (TTBDs). They are very serious BSVs that spread several arboviruses above 80% in lives-stock and have a positive infection of



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

DNA barcoding of waterbirds: A novel technique in environmental conservation biology

Jeganathan Pandiyan^a, Arun Zachariah^b, Bathrachalam Chandramohan^c, Shahid Mahboob^{d,*}, Khalid A. Al-Ghanim^d, Marcello Nicoletti^e, Zaib-Un-Nisa^f, **Marimuthu Govindarajan**^{g,h,*}^aDepartment of Zoology and Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai, Tamil Nadu, India^bDepartment of Forests and Wildlife, Government of Kerala, Sultan Battery, Wayanad, India^cSciGenom Research Foundation, Cheruthuruthy, Kerala, India^dDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^eDepartment of Environmental Biology, Sapienza University of Rome, Rome 00185, Italy^fDepartment of Environmental Sciences and Engineering, Government College University, Faisalabad, Pakistan^gUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^hDepartment of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 20 September 2021

Revised 31 October 2021

Accepted 30 November 2021

Available online 4 December 2021

Keywords:

Waterbirds

DNA barcoding- mt DNA

Phylogenetics

Cytochrome c oxidase I

Conservation

ABSTRACT

Cytochrome c oxidase I (COI) is one of the mitochondrial genes, an excellent marker used for the DNA barcoding of various organisms. The COI was isolated from twelve different species of waterbirds using the Gene Elute DNA miniprep Kit. The relationship among the waterbirds was assessed by making a phylogenetic tree with the software MEGA-X. The phylogenetic tree's dendrogram showed two main branches in which seven species of water birds aligned one group with four subgroups and the remaining five species aligned with two subgroups based on their similar COI sequences. The Little egret showed 96% similarity with the Cattle egret, Purple heron and Oriental Darter produced 94% similarity with the Grey heron, pond heron shared 90% similarity with the Black-crowned night heron, Asian openbill shared 89% with Pheasant-tailed Jacana and the Common coot 94% similarity with White-breasted waterhen. The pond heron and the Black-crowned night heron showed zero % pairwise distance, but the Cattle egret, Little egret, Oriental Darter, Purple heron and Grey heron showed < 0.29%. However, the other seven species of water birds showed > 12% of the pairwise distance. Twenty-one conserved haplotypes have been shown in their COI sequences based on the multiple sequences alignment. DNA barcoding identifies the species with their genetic property rather than based on their ecology and behaviour.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Generally, the taxonomists identify the avian species based on their field characters and behavior, in which several masked and similar morphological species have shown distinct species, based on the DNA barcoding techniques (Hebert et al., 2004). Neverthe-

less, millions of species are yet to be described and identified because expertise in avian taxonomy amongst zoologists is dwindling globally. DNA barcoding is a practical and modern tool for taxonomists who are endowed with preparing inventory and management recommendations of the vast and altering earth biodiversity.

Indeed Hebert et al. (2004) published an article on the barcoding of birds in Central American birds using the COI gene. He has made a breakthrough in taxonomy, wherein mitochondrial DNA, rather than nuclear DNA, formed another most attractive and remarkable protocol in the molecular taxonomy of species by using a novel gene called Cytochrome c Oxidase I (COI). Globally, after Hebert's publication in 2004, several studies have been made and documented the species diversity based on COI, for example, Eurasian birds (Zink et al., 2008), Korean birds (Yoo et al., 2006), Kingfishers (Moyle et al., 2007), Palearctic birds (Kerr et al., 2009a), Scandinavian birds (Johnsen et al., 2010), Marine biodiversity

* Corresponding authors at: Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India.

E-mail addresses: mushahid@ksu.edu.sa (S. Mahboob), drgovind1979@gmail.com (M. Govindarajan).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Changes in the contour of karyology and histoarchitecture of the primary respiratory organ in the fish *Oreochromis mossambicus* (Peters, 1852) inhabiting the polluted estuarine ecosystem

Subramaniam Punitha^a, Rajamanickam Krishnamurthy^a, Kuppusamy Elumalai^a,
Shahid Mahboob^b, Khalid A. Al-Ghanim^b, Zubair Ahmed^b, Ahmed Mustafa^c,
Marimuthu Govindarajan^{d,e,*}

^a Department of Advanced Zoology & Biotechnology, Government Arts College (Autonomous), Chennai, 600035, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Department of Biology, Purdue University Fort Wayne, Fort Wayne, IN, USA

^d Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Wetland ecosystem
Heavy metal toxicity
Contaminations
Oreochromis mossambicus

ABSTRACT

The wetland ecosystem (WE) is subject to pollution by many anthropogenic activities, including domestic and industrial effluents. These effluents may contain toxic heavy metals that can interact within the aquatic ecosystem and have a capacity to disturb the metabolic activities, histological profile, and genetic structure and functions in aquatic species inhabiting the environment. The present study observed the karyological and histological alterations in gills of the freshwater fish, Mozambique tilapia, *Oreochromis mossambicus* in two different sublethal concentrations (1% and 3%) of heavy metals in 7, 15, and 30 days of experimental periods. The heavy metals induced various structural damages such as ring chromosome, sister chromatid exchange, acrocentric association region, condensed chromosomal morphology, heterochromatin region, and nucleolar organizer region in the chromosomes of *O. mossambicus* treated with 1% and 3% sublethal concentrations of water sample collected from Pallikaranai wetland ecosystem. Gills exposed to 1% and 3% effluent exhibited several variations in the respiratory surfaces of gill arches or lamellae in the light and scanning microscopical study. The gills exposed to 1% concentration for 30 days showed marked necrosis, and the secondary lamellae showed the lamellar membrane's dissolution. Exposure of gills to raw effluent in the field condition was observed in the presence of Cd, Pb, Cr, Cu, and Zn. Thus, this present study shows the environmental deterioration by heavy metal pollution on the structure of the gills in tilapia.

1. Introduction

The aquatic and terrestrial regions are highly polluted by various anthropogenic activities, including domestic and industrial effluents, untreated dangerous chemicals, agricultural runoffs, and others (Maurya et al., 2019; Reznia et al., 2021). Major cities in India produce thousands of tons of effluents daily that get dumped in ecologically sensitive areas like streams, ponds, lakes, riverbanks, wetlands, and seashores (Jayaprakash et al., 2010). During the past five decades, ninety percent of Pallikaranai wetland was lost due to dumping, inhabiting, and city development (Vencatesan et al., 2014). Chennai city

produces 3500 tons of waste materials every day, and about 55–60% of waste is dumped around the Pallikaranai wetland ecosystem. These waste materials contain exceedingly toxic heavy metals like Chromium (Cr), Cadmium (Cd), Copper (Cu), Zinc (Zn), Arsenic (As), Mercury (Hg), Nickel (Ni), lead (Pb) (Punitha et al., 2018), that can interact with aquatic residents of the region and disturb their metabolic activities, histology and genetic structure and function in aquatic animals. As heavy metals cannot degrade or be destroyed, they severely damage the ecosystem due to the toxicity persistence, accumulation ability, and bio-magnification in the food chain (Georgieva et al., 2016).

Different fish species are widely used for different toxicity studies for

* Corresponding author. Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India.
E-mail address: drgovind1979@gmail.com (M. Govindarajan).



Research Article

Facile synthesis of silver nanoparticles using the *Simarouba glauca* leaf extract and their impact on biological outcomes: A novel perspective for nano-drug development

Sivaselvi Duraisamy^a, Natesan Vijayakumar^{a,*}, Jayaprakash Rajendran^b, Amalan Venkatesan^a, Balachandran Kartha^c, Senthilkumar Palani Kandasamy^d, Marcello Nicoletti^e, Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f, Marimuthu Govindarajan^{g,h,**}

^a Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India

^b Department of Chemistry, School of Arts and Science, Aarupadai Veedu Campus, Vinayaka Mission's Research Foundation (Deemed to be University), Paiyanoor-603 104, Tamil Nadu, India

^c Department of Chemistry, Vivekanandha College of Engineering for Women, Namakkal, Tamil Nadu, India

^d Department of Microbiology, Faculty of Science, Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India

^e Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^g Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^h Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India



ARTICLE INFO

Keywords:

Simarouba glauca
Nano green synthesis
Anti-microbial
Reverse mutation
Anti-cancer activity
Drug delivery

ABSTRACT

Nowadays, novel green metal-based nanoparticles are receiving global research interest for their potential utilization in several fields, including the formation of nanoparticles for medical treatment. This work reports a synthesis of silver nanoparticles (AgNPs) using *Simarouba glauca* (Sg). The obtained Sg-AgNPs were characterized by UV-visible, FTIR, XRD and SEM. Furthermore, the biological properties of synthesized Sg-AgNPs were investigated for reverse mutation, antimicrobial properties, biofilm analysis, hemolytic activities, DNA fragmentation and MTT assay on HeLa cells. The results revealed a biological efficacy of Sg-AgNPs between 25 and 400 µg/mL concentrations. They showed excellent permeability and a promising drug delivery medium.

1. Introduction

As evidenced by the new vaccine against malaria containing a saponin derived from the bark of *Quillaja saponaria*, plants are a fundamental source of new medical drugs, including treatments for HIV and cancer [1,2]. Besides the activity, the form and the adequate size are key aspects in the delivery and possibility of interacting at the molecular level [3,4]. Due to their size-based properties, nanomaterials have been proposed in different applications [5]. *Simarouba glauca* DC. (Sg), commonly known as the Paradise tree, has shown a potential aid in the treatment of different types of cancer [6–8]. Previously, the Sg chemical constituents were determined and their biological activities investigated [9–12]. In particular, Sg leaf extract contains active cytotoxic

polyphenols, such as tannins, gallic acid, protocatechuic acid and caffeic acid dimethyl ether [13]. The Sg extracts were tested for various biological applications in cancer and microbial inhibition [14]. This study reports the utilization of Sg for bio-catalytic reduction reaction of silver nitrate to silver nanoparticles, whose properties were investigated.

Due to their reduced size, novel metal nanoparticles (NPs) expose suitable biological properties [15–18], such as anti-viral, anti-cancer, antibacterial, antifungal and larvicidal applications [19–23]. The facile preparation of Sg mediated green-synthesized (Sg-AgNPs) is described. The obtained nanoparticles were tested against different microorganisms, biofilm analysis, hemolytic activity, DNA fragmentation, and MTT assay on HeLa cells as part of the thorough screening.

Sg-AgNPs were obtained using microwave energy in the presence of

* Corresponding author. Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalai Nagar 608002, Tamil Nadu, India.

** Corresponding author. Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India.

E-mail addresses: nvkbiochem@yahoo.co.in (N. Vijayakumar), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.jddst.2022.103160>

Received 21 August 2021; Received in revised form 30 January 2022; Accepted 4 February 2022

Available online 7 February 2022

1773-2247/© 2022 Elsevier B.V. All rights reserved.



Original article

Green synthesis, characterization and biological activity of *Solanum trilobatum*-mediated silver nanoparticles

Sengani Manimegalai^{a,*}, Vijayarangan Devi Rajeswari^b, Ranganathan Parameswari^c, Marcello Nicoletti^d, Saud Alarifi^e, **Marimuthu Govindarajan^{f,g}**^aDepartment of Biotechnology, Faculty of Science and Humanities, SRM Institute of Science and Technology, Ramapuram, Chennai 87, Tamil Nadu, India^bDepartment of Bio-Medical Sciences, School of BioSciences and Technology, Vellore Institute of Technology (VIT), Vellore 632 014, Tamil Nadu, India^cDepartment of Biochemistry, Auxilium College, Gandhi Nagar, Vellore 06, India^dDepartment of Environmental Biology, Sapienza University of Rome, Rome 00185, Italy^eDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^fUnit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^gUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 4 October 2021

Revised 10 November 2021

Accepted 17 November 2021

Available online 24 November 2021

Keywords:

Green synthesis

Solanum trilobatum

Inflammation

Antioxidant

Nanoparticle

Cyclooxygenase

ABSTRACT

Biologically inspired synthesis of nanoparticles was found to be more attractive in metal nanoparticle synthesis. The present study reported an *in-situ* biogenic synthesis of silver nanoparticles (AgNPs) using *Solanum trilobatum* aqueous leaf extract. On this basis, the aqueous leaf extract of *S. trilobatum* acted as a reducing agent and stabilizing agent to synthesize highly stable AgNPs at ambient temperature. Eventually, the synthesized and stabilized AgNPs surface plasmon resonance was near 430 nm through a UV–visible (UV–vis) spectrophotometer. Here, the stability of the silver colloids monitored through zeta potential and mean particle size was evaluated through diffraction light scattering (DLF). Further, the average particle size was found to be 27.6 nm and spherical, confirmed with transmission electron microscopy (TEM). Also, colloidal AgNPs and aqueous extract are found to be rich sources of antioxidants and exhibit higher free radical scavenging ability. Thus, efficient inhibition with COX1 and COX2 enzymes and the protective effect with human red blood cell (HRBC) membrane stability showed significant results. These features are promising, suggesting the possibility of the AgNPs to be useful to disease-modifying for treating inflammatory disorders and associated complications.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Regulation of oxidative stress manifests potential therapeutic targets in various cancers and neurological disorders (Barnham et al., 2004). Oxidative stress is correlated with reactive oxygen species (ROS) fluctuations in macromolecules, like lipids, proteins, carbohydrates, and nucleic acids (Rahal et al., 2014). The peroxisomal oxidation in mitochondrial fatty acids can generate ROS, which consumes excessive oxygen. Among the consequences, the accumulation of free radicals and oxidative phosphorylation found in

the mitochondrial respiratory chain down-regulates oxygen metabolism (Ježek and Hlavatá, 2005). Nowadays, the field of neurobiology includes neurogenesis neurotrophins, corticosteroids, inflammatory cytokines, mitochondrial energy generation, and oxidative stress to explain the concepts of bipolar disorder (Berk et al., 2011). Additionally, ROS provokes the mitochondrial permeability abnormalities and induces signal catalyzing to prognoses autophagy, apoptosis, and necrosis (Montaigne et al., 2012). In addition, dysregulation of apoptosis results in inflammations and associated neurodegenerative disorders, arthritis, and types of cancer (Hwang and Kim, 2015). The traditional herbs are polyphenols rich and influential in regulating inflammation and associated symptoms (Singh et al., 2011). Conventionally, these phytochemicals had profound regulation with transcription factors, nuclear factor kappa-light-chain-enhancer of activated B cells, Tumour Necrosis Factor-alpha, Interleukin -1 β , c-Jun N-terminal kinase, Interleukin -6, Mitogen-activated protein kinase 1, Interleukin 1 beta, Mitogen-activated protein kinases and COX-2, which

* Corresponding author.

E-mail address: smanimegalaimsc@gmail.com (S. Manimegalai).

Peer review under responsibility of King Saud University.

<https://doi.org/10.1016/j.sjbs.2021.11.048>

1319-562X/© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Original article

Green synthesis of zinc oxide nanoparticles using *Anoectochilus elatus*, and their biomedical applications

Natesan Vijayakumar^a, Venkataesan Kumari Bhuvaneshwari^a, Gandhimathi Kaliyamoorthi Ayyadurai^b, Rajendran Jayaprakash^c, Kasi Gopinath^{d,*}, Marcello Nicoletti^e, Saud Alarifi^f, **Marimuthu Govindarajan^{g,h}**^aDepartment of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalaiagar 608002, Tamilnadu, India^bDepartment of Chemistry, Sri Sairam Engineering College, Sai Leo Nagar, West Tambaram, Chennai 600044, Tamilnadu, India^cDepartment of Chemistry, School of Arts and Science, Aarupadai Veedu Institute of Technology, Vinayaka Mission's Research Foundation (Deemed to be University), Paiyanoor 603104, Tamilnadu, India^dDepartment of Packaging, Yonsei University, 1 Yonseidae-gil, Wonju-si, Gangwon-do 26493, South Korea^eDepartment of Environmental Biology, Sapienza University of Rome, Rome 00185, Italy^fDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^gUnit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalaiagar-608 002, Tamilnadu, India^hUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamilnadu, India

ARTICLE INFO

Article history:

Received 29 September 2021

Revised 14 November 2021

Accepted 28 November 2021

Available online 3 December 2021

Keywords:

Anoectochilus elatus

Nanotechnology

Zinc Oxide nanoparticles

Antimicrobial activity

Anti-inflammatory activity

ABSTRACT

Zinc and its derivatives requirement increased to enhance human immunity against the different pandemics, including covid-19. Green synthesis is an emerging field of research. Zinc oxide (ZnO) nanoparticles have been prepared from *Anoectochilus elatus* and characterized using absorption, vibrational and electron microscope analysis. They were carried for antibacterial, inflammatory control tendency, and potential antioxidant activities. The brine shrimp lethal assay tested the biologically derived nanomaterial toxicity and the lethal concentration (LC₅₀) is 599.79 µg/ml. The inhibition against the important disease-causing pathogens was measured against four-gram negative, gram-positive bacteria and two fungus pathogens. The nanomaterial exposed inhibition zone for gram-positive bacteria between 17 mm and 25 mm. The inhibition zone against gram-negative bacteria exists between 19 mm and 24 mm. The anti-inflammatory activity was assessed by inhibition of protein denaturation and protease inhibitory activity using nanomaterial. The antioxidant activity was examined using four assays for the therapeutic activities. The average size range of 60–80 nm nanoparticles has prepared and exposed the good biological activity between 50 µg/ml and 100 µg/ml. The comparative results of anti-inflammatory and antioxidant assay results with standards such as Aspirin and vitamin C exposed that two to three times higher concentrations are required for the fifty percent of inhibitions. The prepared low-cost nanoparticle has exhibited excellent biological activity without any side effects and may enhance immunity.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: gloriousnanogopi@gmail.com (K. Gopinath).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

Green synthesis has received more attention for various applications because of the low-cost, eco-friendly, safe nanoparticles (NPs) preparation (Govindarajan et al., 2016a,b; Govindarajan and Benelli, 2017; Balalakshmi et al., 2017; Suganya et al., 2017; Balasooriya et al., 2017). Due to the size and shape of the nanomaterials, they have shown various applications in medicinal fields (Narayanan and Sakhivel, 2011). Synthesis of new molecules using the biosynthetic technique is a growing field and the method using bio-extracts has expected more responsiveness than chemical and physical methods for the nanomaterial preparation (Iravani, 2011).

<https://doi.org/10.1016/j.sjbs.2021.11.065>

1319-562X/© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Available online at www.sciencedirect.com

jmr&t
Journal of Materials Research and Technology
journal homepage: www.elsevier.com/locate/jmrt



Original Article

Synthesis of Cu-MOF/CeO₂ nanocomposite and their evaluation of hydrogen production and cytotoxic activity



Shanmugasundaram Kumaraguru ^a, Ravi Nivetha ^b, Kasi Gopinath ^c,
Elumalai Sundaravadivel ^{a,**}, Bader O. Almutairi ^d, Mikhlid H. Almutairi ^d,
Shahid Mahboob ^d, M.R. Kavipriya ^e, Marcello Nicoletti ^f,
Marimuthu Govindarajan ^{g,h,*}

^a Department of Chemistry, SRM Institute of Science and Technology, Kattankulathur, 603203, Chengalpattu District, Tamilnadu, India

^b Center for Nanotechnology Research, Vellore Institute of Technology, Vellore, 632014, Tamilnadu, India

^c Department of Packaging, Yonsei University, 1 Yonseidaegil, Wonju-si, Gangwon-do 26493, South Korea

^d Department of Zoology, College of Science, King Saud University, P.O. Box 2455, 11451, Riyadh, Saudi Arabia

^e Department of Botany, School of Life Sciences, Bharathidasan University, Tiruchirappalli, 620024, Tamil Nadu, India

^f Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

^g Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^h Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 16 November 2021

Accepted 5 March 2022

Available online 18 March 2022

Keywords:

Nanomaterials

Energy

Electrocatalysis

Cytotoxicity

Biocompatibility

ABSTRACT

In this investigation, new composites of cerium oxide were incorporated in the copper-based metal–organic framework using trimesic acid by the precipitation method. Synthesized Cu-MOF/CeO₂ nanocomposite is characterized by numerous analytical methods to interpret their structure, morphology, and thermal behaviour. Surface micrograph analysis revealed that spherical particles and cerium oxide nanoparticles are embedded in the Cu-MOF matrix. The prepared nanocomposite shows a lesser onset potential and overpotential with a high current density (18.6 mA cm⁻²). The Cu-MOF/CeO₂ composite exhibits a small Tafel slope of 54.6 mV dec⁻¹ and suggests that the hydrogen evolution reaction follows the Heyrovsky mechanism. On the other hand, five different MOF/CeO₂ nanocomposite concentrations were tested on a human osteosarcoma cell line (MG63). The 50% cell mortality was observed at 97.9 µg/ml and it proved less cytotoxicity effect with better biocompatibility. Thus, overall results showed that the nanocomposite as an

* Corresponding author.

** Corresponding author.

E-mail addresses: sundarae@srmist.edu.in (E. Sundaravadivel), drgovind1979@gmail.com (M. Govindarajan).<https://doi.org/10.1016/j.jmrt.2022.03.028>2238-7854/© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



Toxicity evaluation of polypropylene microplastic on marine microcrustacean *Artemia salina*: An analysis of implications and vulnerability

Jeyaraj Jeyavani^a, Ashokkumar Sibiyi^a, Sundaresan Bhavaniramy^a, Shahid Mahboob^b, Khalid A. Al-Ghanim^b, Zaib-un Nisa^c, Mian Nadeem Riaz^d, Marcello Nicoletti^e, Marimuthu Govindarajan^{f,g}, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Science Campus 6th Floor, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Department of Environmental Science and Engineering, Government College University, Faisalabad, Pakistan

^d Texas A&M University, College Station, TX, USA

^e Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy


^f Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^g Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

HIGHLIGHTS

- Various life stages of *Artemia* easily intake polypropylene microplastics (11–44 μm).
- Microplastics were accumulated in their tract, triggering the redox-hemostasis.
- Microplastic consumption disturbs the antioxidant biomarkers and neurotransmitter activity.
- Mortality was observed in nauplii & meta-nauplii; in juveniles, only behaviour changes.
- Morphological analysis of the juvenile intestinal epithelial layer was damaged.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling Editor: Venkatramanan Senapathi

Keywords:
Microplastics
Pollution
Brine shrimp
Biomarkers
Histology

ABSTRACT

Polypropylene microplastic particles are one of the predominant pollutants in marine ecosystems and their toxic effects are unknown in aquatic biota. The study aims to prepare the spherical shaped polypropylene microplastics (size range 11.86 μm –44.62 μm) and assess their toxic effects (1, 25, 50, 75 and 100 $\mu\text{g}/\text{mL}$) in various life stages (nauplii, metanauplii and juvenile) of marine microcrustacean *Artemia salina* within 48 h. In addition, microplastics ingestion by *Artemia* nauplii was proved by FTIR analysis. The results revealed, microplastics accumulation in their tract leads to change in their homeostasis, as followed increase in the oxidative burst causes mortality in nauplii (LC₅₀ 40.947 $\mu\text{g}/\text{mL}$) and meta nauplii (LC₅₀ 51.954 $\mu\text{g}/\text{mL}$). In juvenile, swimming behaviour was changed. Moreover, microplastic consumption disturbs the antioxidant biomarkers such as superoxide dismutase (SOD), catalase (CAT), reduced glutathione (GSH), glutathione -S- Transferase (GST) and

* Corresponding author.

E-mail addresses: vaseeharanb@gmail.com, vaseeharanb@alagappauniversity.ac.in (B. Vaseeharan).

<https://doi.org/10.1016/j.chemosphere.2022.133990>

Received 3 December 2021; Received in revised form 21 January 2022; Accepted 11 February 2022

Available online 18 February 2022

0045-6535/© 2022 Elsevier Ltd. All rights reserved.



Synthesis and physicochemical characteristics of Ag-doped hydroxyapatite nanoparticles, and their potential biomedical applications

Saleth Sebastiammal^{a,i,*}, Arul Sigamani Lesly Fathima^{a,i}, Saud Alarifi^b, Shahid Mahboob^b, Johnson Henry^c, M.R. Kavipriya^d, Marimuthu Govindarajan^{e,f,**}, Marcello Nicoletti^g, Baskaralingam Vaseeharan^h

^a Research Department of Physics, Holy Cross College (Autonomous), Nagercoil, 629004, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Department of Physics, Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli, 627012, Tamil Nadu, India

^d Department of Botany, School of Life Sciences, Bharathidasan University, Tiruchirappalli, 620024, Tamil Nadu, India

^e Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608002, Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612001, Tamil Nadu, India

^g Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

^h Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Science Campus, Karaikudi, 630004, Tamil Nadu, India

ⁱ Affiliated to Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli, 627012, Tamil Nadu, India

ARTICLE INFO

Keywords:

Green nanotechnology
Ag-nanostructure
Biomaterials
HeLa cell Lines

ABSTRACT

In this present scenario, hydroxyapatite (HAp) nanostructures were synthesized through green routes for biomedical applications, particularly remediation towards human pathogens and cancer cells. The present study aims at forming non-toxic and eco-friendly silver (Ag⁺) doped HAp using Polyethylene glycol (PEG), Cetyl Trimethyl ammonium bromide (CTAB) and curcumin. Ag⁺ doped HAp nanoparticles (NPs) were prepared by the sol-gel method with a cube and rod-like morphology. Ag-HApNPs showed a sharp and well-defined diffraction peak, which possesses the hexagonal crystalline structure with space group P6₃/m. The Fourier-transform infrared spectroscopy and Raman spectra confirmed the formation of Ag-HApNPs, and the bandgap values were obtained using UV-DRS analysis. The Ag-HApNPs with PEG, CTAB and curcumin might be fabricated materials were examined against antibacterial, antifungal, antioxidant, and cytotoxic activities, which provided exemplary biomedical applications. Overall, Ag-HApNPs can be used as potential drug delivery and perspectives to control multidrug-resistant pathogens.

1. Introduction

Recently, the application of nanomaterials in biomedical aspects (Govindarajan et al., 2016; Govindarajan and Benelli, 2017; Ishwarya et al., 2017; Chi et al., 2022; Das et al., 2022) considerably increased by their novelty of physicochemical properties and it has been intensively analyzed due to their peculiar properties and high human osseous tissue similarity (Turkoz et al., 2013; Suganya et al., 2017; Shen et al., 2017; Balalakshmi et al., 2017). A revolution has arisen in using bioceramics to enhance human life quality during the last fifty years. The biocompatibility of calcium phosphate-based bioceramics, such as hydroxyapatite

(HAp), is the key factor for dental and orthopedic applications (Fakharzadeh et al., 2017). Hydroxyapatite is a significant biocompatible and osteoconductive substance with finite antibacterial resistance (Khalili et al., 2017). Properties of HAp can be enhanced by substituting some cations, including Ag⁺, Sr²⁺, Mg²⁺, Zn²⁺, Ce³⁺, La³⁺, Y³⁺ and Gd³⁺ into hydroxyapatite lattice (Kalita et al., 2004; Pietak et al., 2007; Bang et al., 2011; Suganthi et al., 2011; Webster et al., 2004; Ahymah Joshy et al., 2011; Yasukawa et al., 2012a,b; Yang et al., 2008) which leads to enhance the biological properties. Many techniques were adopted for the synthesis of monovalent and divalent doped HAp, which includes co-precipitation (Paluszkiwicz et al., 2010; Dubnika et al.,

* Corresponding author. Research Department of Physics, Holy Cross College (Autonomous), Nagercoil, 629004, Tamil Nadu, India.

** Corresponding author. Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608002, Tamil Nadu, India.

E-mail addresses: sebastiscc@gmail.com (S. Sebastiammal), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.envres.2022.112979>

Received 30 December 2021; Received in revised form 15 February 2022; Accepted 17 February 2022

Available online 23 February 2022

0013-9351/© 2022 Elsevier Inc. All rights reserved.



Biomimetically synthesized *Physalis minima* fruit extract-based zinc oxide nanoparticles as eco-friendly biomaterials for biological applications

Mariappan Yazhiniprabha^a, Sasikumar Banu^a, Ramachandran Ishwarya^a,
Viswanathan Vinotha^a, Marimuthu Govindarajan^{b,c}, Mohammad Ahmad Wadaan^d,
Shahid Mahboob^d, Marcello Nicoletti^e, Baskaralingam Vaseeharan^{a,*}

^a Nanobioscience and Nanopharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^c Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^d Department of Zoology, College of Science, King Saud University, P.O. Box 2455, Riyadh, 11451, Saudi Arabia

^e Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

ARTICLE INFO

Keywords:

Physalis minima
Green synthesis
Antibacterial activity
Biocompatibility

ABSTRACT

In the present study, the fruit extract of *Physalis minima* was used to synthesize zinc oxide nanoparticles (*Pm*-ZnO NPs). The first validation of the *Pm*-ZnO NPs production was obtained using a UV-Vis spectrophotometer at 327 nm. *Pm*-ZnO NPs crystalline nature was determined as hexagonal, relating to a joint corporation of powder diffraction standards (JCPDS) 75-1526 examined in XRD analysis. FTIR spectroscopy was used to identify functional groups associated with *Pm*-ZnO NPs, i.e., O-H and C=N. *Pm*-ZnO NPs were measured to be 50-150 nm in size and spherical in form using a TEM. Arrested augmentation of Gram-positive (*Enterococci faecalis*) and Gram-negative bacteria (*Pseudomonas aeruginosa*) populations was recorded at a 50 µg mL⁻¹ concentration of *Pm*-ZnO NPs, where the zones of inhibition were measured as 31.4 ± 2.8 mm & 33.5 ± 0.8 mm, respectively. *Pm*-ZnO NPs inhibited the growth of bacteria *E. faecalis* and *P. aeruginosa* at a minimum concentration of 30 µg mL⁻¹. Inverted microscopic images revealed that *Pm*-ZnO NPs diminished bacterial biofilm thickness at 50 and 100 µg mL⁻¹. The biocompatibility of *Pm*-ZnO NPs was examined by potential antioxidant properties (DPPH assay). Biototoxicity values of *Pm*-ZnO NPs were determined as LC₅₀ - 76.5 (68.7-86.4) µg mL⁻¹ and LC₉₀ - 153.4 (132.4-189.6) µg mL⁻¹ on 48 h treatment with *Artemia salina* nauplii and damages in the mid-gut region were examined under a stereomicroscope. *Pm*-ZnO NPs exhibited photocatalytic activity at 48 h duration against methylene blue (MB) dye. Concludingly, all the reported findings suggest the utilization of these generated nanoparticles as nanomedicine against microbial infection.

1. Introduction

Human health is at risk, due to emerging of new alerts involving infectious microbial diseases [1]. Among several microbial infections, the disease caused by multi-drug resistant (MDR) bacteria is considered the most challenging task by medical practitioners and researchers focused on finding a permanent solution to cure those infections. Resistance is the consequence of the overuse of various antibiotics in clinical medicines that provoked the generation of MDR strains of bacterial pathogens [2]. *Enterococcus faecalis* is a Gram-positive facultative anaerobic bacterium that colonises the human mouth cavity and

gastrointestinal system [3]. *E. faecalis* resistant to vancomycin is commonly a urinary tract infectious pathogen that infects the endocardium, bloodstream, biliary tract, abdomen and burns wounds [4-6]. Similarly, *Pseudomonas aeruginosa*, a Gram-negative bacterium, is an opportunistic infectious disease in immunocompromised people and causes nosocomial infections that are resistant to various medications, such as cephalosporins and penicillin carbapenems, cephamycins and cefepime [7-10]. Hence, an urge to design a bactericide against these two dangerous MDR pathogens is essential.

The interdisciplinary field "Nanopharmacology" incorporates nanotechnology and medicine to assist disease diagnosis, drug delivery,

* Corresponding author.



E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jddst.2022.103475>

Received 9 February 2022; Received in revised form 18 May 2022; Accepted 27 May 2022

Available online 30 May 2022

1773-2247/© 2022 Elsevier B.V. All rights reserved.



Swift synthesis of zinc oxide nanoparticles using unripe fruit extract of *Pergularia daemia*: An enhanced and eco-friendly control agent against Zika virus vector *Aedes aegypti*

Ramachandran Ishwarya^{a,b}, Rengarajan Jayakumar^c, Tamilmani Govindan^b,
 Marimuthu Govindarajan^{d,e}, Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f,
 Marcello Nicoletti^g, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, Tamil Nadu 630004, India

^b Mandapam Regional Centre, ICAR - Central Marine Fisheries Research Institute, Mandapam, Tamil Nadu, India

^c ICAR - Central Institute of Brackishwater Aquaculture (CIBA), DARE, Ministry of Agriculture, Government of India, #75, Santhome High Road, Raja Annamalaiapuram, MRC Nagar, Chennai, Tamil Nadu 600 028, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu 608 002, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu 612 001, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^g Department of Environmental Biology, Sapienza University of Rome, Rome 00185, Italy

ARTICLE INFO

Keywords:

Green synthesis
Pergularia daemia
 ZnONPs
Aedes aegypti
 Larvicidal activity

ABSTRACT

In this study *Pergularia daemia* unripe fruits were used to synthesize zinc oxide nanoparticles (*Pd*-ZnONPs). UV-vis Spectroscopy detected the production of ZnONPs. XRD, FTIR, SEM, and TEM studies were used to characterize the synthesized *Pd*-ZnONPs. *Aedes aegypti* (*Ae. aegypti*) third instar larvae were analyzed to diverse concentrations of *Pd*-unripe fruit extract and *Pd*-ZnONPs for 24 hours to assess the larvicidal effect. Mortality was also detected in *Ae. aegypti* larvae under laboratory conditions, with corresponding LC₅₀ and LC₉₀ values of 11.11 and 21.20 µg/ml respectively. As a result of this study, the levels of total proteins, esterases, acetylcholine esterase, and phosphatase enzymes in the third instar larvae of *Ae. aegypti* were significantly lower than the control. These findings suggest that *Pd*-ZnONPs could be used to suppress mosquito larval populations.

1. Introduction

Along with its ecological and physiological flexibility, the *Ae. aegypti* mosquito is now acknowledged as the world's most common invasive mosquito, spreading its range through consignments of bamboos, wet shrubbery and water containers, the worldwide commerce of used tires (Benelli and Govindarajan, 2017; Govindarajan, 2017; Fahimnisha et al., 2020). Dengue control requires a reduction in *Ae. aegypti* populations (WHO, 2012). Dengue fever is the majority universal mosquito-borne viral disease, disturbing nearly a quarter of the world population and spreading rapidly (WHO, 2016). Mosquito populations have developed resistance to synthetic chemical pesticides, which can have unintended consequences for beneficial non-target creatures (Melo-Santos et al., 2010; Govindarajan, 2011a,b; Govindarajan and Benelli, 2016, 2017).

Due to their cheap cost and limited cross-resistance, natural insecticides have been studied for their larvicidal effects on *Ae. aegypti* resistant lines (Govindarajan et al., 2005; Braga and Valle, 2007; Sim et al., 2007). Insects' acetylcholinesterase and detoxifying enzymes have been tested for inhibitory effects on green plants' larvicidal function to find synergists for commonly used pesticides (Larson et al., 2010; Maheswaran and Ignacimuthu, 2012). Due to their eco-friendly qualities and great biodegradability they have recently become a viable alternative for vector control via plant secondary metabolites (Mathivanan et al., 2010; Govindarajan and Sivakumar, 2012; Benelli, 2015; Govindarajan et al., 2016a). *Pergularia daemia* (*P. daemia*) (Forssk.) Chiov. is a milky plant of the Asclepiadaceae family (Pankaj, 2003). *P. daemia*, also called as "Veliparuthi" in Tamil, is used to cure infantile diarrhea and malarial intermittent fevers (Kirtikar and Basu, 1999; Nadkarani, 1976). Phytochemicals like cardenolides, alkaloids, and saponins have been

* Corresponding author.

E-mail address: vaseeharanb@alagappauniversity.ac.in (B. Vaseeharan).

<https://doi.org/10.1016/j.actatropica.2022.106489>

Received 27 March 2022; Received in revised form 22 April 2022; Accepted 24 April 2022

Available online 27 April 2022

0001-706X/© 2022 Elsevier B.V. All rights reserved.



Azadirachta indica-wrapped copper oxide nanoparticles as a novel functional material in cardiomyocyte cells: An ecotoxicity assessment on the embryonic development of *Danio rerio*

Santhoshkumar Jayakodi^a, Rajeshkumar Shanmugam^{b,*}, Bader O. Almutairi^c, Mikhliid H. Almutairi^c, Shahid Mahboob^c, M.R. Kavipriya^d, Ramesh Gandusekar^e, Marcello Nicoletti^f, Marimuthu Govindarajan^{b,g,h,**}

^a Department of Biotechnology, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Science (SIMATS), Chennai, 602105, TN, India

^b Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science (SIMATS), Chennai, 600077, TN, India

^c Department of Zoology, College of Science, King Saud University, P.O. Box 2455, Riyadh, 11451, Riyadh, Saudi Arabia

^d Department of Botany, School of Life Sciences, Bharathidasan University, Tiruchirappalli, 620024, Tamil Nadu, India

^e Department of Regenerative Medicine & Immune Regulation, Medical University of Bialystok (MUB), Poland

^f Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

^g Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^h Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

ARTICLE INFO

Keywords:

Neem
CuO nanoparticles
H9c2 cell
Apoptosis
Zebrafish embryos

ABSTRACT

This research reports on the production of copper oxide nanoparticles (CuO NPs) through the green synthesis method using *Azadirachta indica* (Ai) flower extract. Synthesized Ai-CuO NPs are characterized by Zeta Potential, TGA, SEM and TEM analysis. The Ai-CuO NPs gave a maximum peak at 270 nm. As per XRD studies, the Ai-CuO NPs obtained were crystalline. FTIR spectrum Ai-CuO NPs showed the presence of functional groups like the O–H group, aromatic group, etc. TEM and SEM assist in investigating the size and morphology of the Ai-CuO NPs, which were spherical and varied in size between 10.11 nm and 17.54 nm. EDAX showed that Ai-CuO NPs were pure with no impurities. The synthesized Ai-CuO NPs were then analyzed for their cytotoxicity at various concentrations (5, 10, 20, 30, 40 and 50 µg/mL) against H9c2 cardiomyocyte cells using MTT assay. DOX-induced H9c2 cell damage of apoptosis and ROS. The nanoparticle formed by Ai-CuO was cured with different concentrations (5, 10 and 20 µg/mL). In zebrafish, 48 hpf and 72 hpf were measured at 75 µM to reduce dysfunction and mortality during organ development. These results can have a beneficial impact on eco-toxicological effects.

1. Introduction

Nanoparticles are ultra-fine inorganic particles existing in the natural world surrounding an interfacial layer created by human activities (Batista et al., 2015; Benelli and Govindarajan, 2017; Balalakshmi et al., 2017). They are defined as solid particles with a size in the range of 10–100 nm that shows particulate dispersions (Mohanraj and Chen., 2006; Govindarajan and Benelli, 2017; Ishwarya et al., 2017; Fahim-munisha et al., 2020). Biotechnology scientists have shown proven biology and material science (De et al., 2008; Suganya et al., 2017; Thyaga et al., 2018; Vinotha et al., 2019). The applications of

nanoparticles, mainly in medical nanotechnology, provide enormous opportunities (Gao et al., 2009). This may eventually include protein and peptide delivery, drug delivery, nanoparticle targeting, medical surgeries and endo visualizations, whereas the similar term biomedical nanotechnology is used in the biomedical sector (Ahmed et al., 2012). Since these nanoparticles provide a large surface area to the volume ratio, they also exemplify magnetic and optic roles in biology and sensing and catalysts. Compared to any other bulk materials, they can be considered as potentials regarding in antimicrobial and anticancer drugs (Prakash et al., 2018). Some applications include semiconductor doping such as antimicrobial agents, chemical sensors and anticancer (Karthika

* Corresponding author.

** Corresponding author. Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India.

E-mail addresses: ssrajeshkumar@hotmail.com (R. Shanmugam), drgovind1979@gmail.com, dr.m.govindarajan@gcw.ac.in (M. Govindarajan).

<https://doi.org/10.1016/j.envres.2022.113153>

Received 28 February 2022; Received in revised form 15 March 2022; Accepted 18 March 2022

Available online 24 March 2022

0013-9351/© 2022 Elsevier Inc. All rights reserved.



Comparative toxicity of silver nanoparticles and silver nitrate in freshwater fish *Oreochromis mossambicus*: A multi-biomarker approach

Ashokkumar Sibiya^a, Narayanan Gopi^a, Jeyaraj Jeyavani^a, Shahid Mahboob^b,
Khalid A. Al-Ghanim^b, Salma Sultana^c, Ahmed Mustafa^d, Marimuthu Govindarajan^{e,f},
Baskaralingam Vaseeharan^{a,*}

^a Nano biosciences and Nanopharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Science Campus 6th Floor, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^c Department of Zoology, Government College University, Faisalabad 38000, Pakistan

^d Department of Biological Sciences, Purdue University Fort Wayne, IN 46805, USA

^e Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Edited by Martin Grosell

Keywords:

Oreochromis mossambicus
Comparative toxicity
AgNPs
AgNO₃
Oxidative stress
Neurotoxicity

ABSTRACT

Silver nanoparticles (AgNPs) in the aquatic environment affect ecological repercussions and have fatal impacts on aquatic animals. The current study examined and correlated the toxicity of silver nitrate (AgNO₃) and silver nanoparticles (AgNPs) to the Mozambique tilapia, *Oreochromis mossambicus*. The comparative toxicity studies were done by exposing *O. mossambicus* to various doses of AgNO₃ and AgNPs (0, 25, 50, 75, and 100 µg/L) over a 7-day subacute exposure period. AAS analysis was used to detect Ag accumulation, while the histological examination established gill tissue damage. Oxidative stress affects lipid peroxidation (LPO) and protein carbonyl activity (PCA) in the gill tissue. Antioxidant parameters such as glutathione-S-transferase (GST), glutathione peroxidase (GPx), superoxide dismutase (SOD), catalase activity (CAT), and non-enzymatic antioxidants such as metallothionein (MT) and reduced glutathione. The serum in the blood was used to determine non-specific immunological characteristics such as lysozyme (LYZ), myeloperoxidase (MPO), and respiratory burst activity (RBA). The neurotoxic impact of acetylcholine esterase activity (AChE) was investigated in brain tissues. The findings demonstrated that larger concentrations of AgNO₃ than AgNPs improved enzymatic antioxidant activities in the gill tissue. Histological examination of fish gills demonstrated that both AgNPs and AgNO₃ induced telangiectasia and epithelial cell hyperplasia. By increasing the concentration of AgNPs and AgNO₃, the present research demonstrated that silver accumulation leads to inefficient oxidative stress and altered enzymatic and non-enzymatic parameters, leading to cellular damage.

1. Introduction

Engineered nanomaterials are increasing due to their many uses in biology and biomedicine, health care technology, home appliances, pharmaceutical, textile, cosmetic, and electronics sectors (Hana et al., 2020; Fabrega et al., 2011; Benelli and Govindarajan, 2017; Ishwarya et al., 2017). The buildup of nanomaterials in freshwater habitats due to human activities leads to the contamination of freshwater ecosystems with nanomaterials (Shah and Mraz, 2019; Jeyavani et al., 2022; Paulpandian et al., 2022; Rashidian et al., 2021; Sibiya et al., 2022). Because

of the heavy usage of the above-said materials, the impact of the nanomaterial contamination on the aquatic organism is of great alarm as they are highly toxic, tenacious, non-decomposable and vastly bio-accumulative (Jafarabadi et al., 2020).

The hazardous nature of nanoparticles (NPs) in the aquatic ecosystem can be well evaluated by considering their bioavailability and toxic nature to the aquatic organisms (Nowack and Bucheli, 2007). NPs and their ionic forms may interact at an elevated concentration which possibly causes antagonistic effects on aquatic organisms (Griffitt et al., 2007; Xiang et al., 2020). Nevertheless, the ecological consequences of

* Corresponding author.

E-mail address: vaseeharanb@alagappauniversity.ac.in (B. Vaseeharan).

<https://doi.org/10.1016/j.cbpc.2022.109391>

Received 26 January 2022; Received in revised form 18 May 2022; Accepted 29 May 2022

Available online 2 June 2022

1532-0456/© 2022 Elsevier Inc. All rights reserved.



Citrullus lanatus-encased zinc oxide nanoparticles as potential anti-diabetic, anti-inflammatory and antibacterial agents: A new strategy towards biocompatible nano-drugs

Ramachandran Ishwarya^{a,b}, Jeyaraj Jeyavani^a, Rengarajan Jayakumar^b, Saud Alarifi^c,
Marimuthu Govindarajan^{d,e}, Marcello Nicoletti^f, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Mandapam Regional Centre, ICAR - Central Marine Fisheries Research Institute, Mandapam, Tamil Nadu, India

^c Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^f Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

ARTICLE INFO

Keywords:

Drug development
Diabetic
Citrullus lanatus
seed Extract

ABSTRACT

This research aimed to develop a rapid, cheap, and environmentally friendly method for synthesizing zinc oxide nanoparticles (ZnONPs) from *Citrullus lanatus* (Cl) seed extract and the Cl-ZnONPs were analyzed using several techniques to determine their properties. These included UV-vis spectroscopy, XRD, FTIR, SEM, and TEM. Cl-ZnONPs have sizes between 10 and 50 nm and are spherical in form. Compared to Cl seed extract, Cl-ZnONPs displayed significantly higher levels of antioxidant activity. Comparatively to α -amylase, Cl-ZnONPs showed much stronger inhibition of β -glucosidase and DPPIV (anti-diabetic activity). Cl-ZnONPs and Cl-seed extract were tested for their ability to reduce inflammation, demonstrating potent in vitro anti-inflammatory properties (membrane stabilization, proteinase inhibitory, and lipoxygenase). In addition, the Cl-ZnONPs' antibacterial activity revealed that they had a significantly greater effect on inhibiting the growth of Gram-negative bacteria such as *Escherichia coli* and *Pseudomonas aeruginosa*. Synthesized Cl-ZnONPs are promising solutions for treating inflammatory and diabetic concerns, as suggested by the outcomes. In conclusion, the biosynthesized Cl-ZnONPs shown in this study provide a synthetic substance replacement that is safe and effective in various biomedical and pharmaceutical applications, including those requiring antioxidant, antibacterial, and anti-inflammatory properties.

1. Introduction

Biomedical nanomaterials have recently received much interest because of their well-known biological and biomedical uses. A decade ago, more than 30,000 t of ZnO NPs were produced annually, and it is the most prominent metal oxide nanoparticle. It is widely used in various sectors because of its unique physical and chemical characteristics [1,2] and extra effortlessly immersed through the body when the particle size is tiny. As a consequence, nano-ZnO is widely applied in food. Furthermore, the US Food and Drug Administration (FDA) has categorized ZnO as a "GRAS" (generally recognized as safe) material [3]. ZnO NPs are relatively affordable and less toxic than other metal oxide NPs, and have a broad spectrum of application uses, including in

medical and pharmaceutical fields [4-6].

Diabetes is a metabolic disorder distinguished through lack of secretion or action of endogenous insulin resulting in the elevation of blood glucose and several other problems and it is one of the most significant public troubles. According to World Health Organization (WHO) estimates, diabetes will reach 300 million or more by 2025 [7]. Moreover, after an injury or infection, inflammation is a multifactorial phenomenon that restores cellular homeostasis and the tissue microenvironment. Nonsteroidal anti-inflammatory drugs have various side effects like stomach ulcers, bleeding, and cardiovascular strokes, limiting their use in treating chronic inflammation diseases [8,9]. This emphasizes the importance of activity-based phytopharmacological evaluation of herbal medications. More awareness has been paying attention to

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jics.2022.100703>

Received 20 June 2022; Received in revised form 19 August 2022; Accepted 28 August 2022

Available online 2 September 2022

0019-4522/© 2022 Indian Chemical Society. Published by Elsevier B.V. All rights reserved.



Article

Terminalia chebula-Assisted Silver Nanoparticles: Biological Potential, Synthesis, Characterization, and Ecotoxicity

Munusamy Tharani¹, Shanmugam Rajeshkumar^{1,*}, Khalid A. Al-Ghanim², Marcello Nicoletti³, Nadezhda Sachivkina^{4,*} and Marimuthu Govindarajan^{5,6}

¹ Nanobiomedicine Lab, Department of Pharmacology, Saveetha Dental College and Hospitals, SIMATS, Saveetha University, Chennai 600 077, Tamil Nadu, India

² Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

³ Department of Environmental Biology, Sapienza University of Rome, 00185 Rome, Italy

⁴ Department of Microbiology V.S. Kiktenko, Institute of Medicine, Peoples Friendship University of Russia Named after Patrice Lumumba (RUDN University), Moscow 117198, Russia

⁵ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

⁶ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

* Correspondence: rajeshkumars.sdc@saveetha.com (S.R.); sachivkina@yandex.ru (N.S.)

Abstract: In the current research, an aqueous extract of *Terminalia chebula* fruit was used to produce silver nanoparticles (Ag NPs) in a sustainable manner. UV-visible spectrophotometry, transmission electron microscopy (TEM), and scanning electron microscopy (SEM) were used to characterize the synthesized nanoparticles. Synthesized Ag NPs were detected since their greatest absorption peak was seen at 460 nm. The synthesized Ag NPs were spherical and had an average size of about 50 nm, with agglomerated structures, as shown via SEM and TEM analyses. The biological activities of the synthesized Ag NPs were evaluated in terms of their antibacterial and antioxidant properties, as well as protein leakage and time-kill kinetics assays. The results suggest that the green synthesized Ag NPs possess significant antibacterial and antioxidant activities, making them a promising candidate for therapeutic applications. Furthermore, the study also evaluated the potential toxicological effects of the Ag NPs using zebrafish embryos as a model organism. The findings indicate that the synthesized Ag NPs did not induce any significant toxic effects on zebrafish embryos, further supporting their potential as therapeutic agents. In conclusion, the environmentally friendly production of Ag NPs using the extract from *T. chebula* is a promising strategy for discovering novel therapeutic agents with prospective uses in biomedicine.

Keywords: *Terminalia chebula*; silver nanoparticles; mechanism; embryonic toxicology; antibacterial; antioxidant agent



Citation: Tharani, M.; Rajeshkumar, S.; Al-Ghanim, K.A.; Nicoletti, M.; Sachivkina, N.; Govindarajan, M. *Terminalia chebula*-Assisted Silver Nanoparticles: Biological Potential, Synthesis, Characterization, and Ecotoxicity. *Biomedicines* **2023**, *11*, 1472. <https://doi.org/10.3390/biomedicines11051472>

Academic Editor: Matteo Calvaresi

Received: 22 April 2023

Revised: 10 May 2023

Accepted: 16 May 2023

Published: 18 May 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The distinct characteristics of nanomaterials have garnered increased interest from scientists, thereby facilitating diverse biomedical implementations [1]. The fabrication of nanoparticles with diverse structures can be accomplished through physical, biological, and chemical methodologies. However, physical and chemical methods often result in low yields, high costs, high energy consumption, and environmental damage. Biological synthesis, which employs plants and microbes as reducing agents, is one of the three synthesis approaches [2,3]. Because of their superior thermal, strong optoelectronic, catalytic, and surface volume ratio properties, metal nanoparticles play an essential role in the healthcare sectors. This is due to the fact that they have great physicochemical features. Silver, gold, and platinum are characterized as the “noble metal nanoparticles” among all of the other metal nanoparticle types [4,5]. Silver nanoparticles have been paid special



Article

Probiotic-Bacteria (*Lactobacillus fermentum*)-Wrapped Zinc Oxide Nanoparticles: Biosynthesis, Characterization, and Antibacterial Activity

Rajeshkumar Shanmugam ^{1,*}, Tharani Munusamy ¹, Santhoshkumar Jayakodi ², Khalid A. Al-Ghanim ³, Marcello Nicoletti ⁴, Nadezhda Sachivkina ⁵ and Marimuthu Govindarajan ^{6,7,*}

- ¹ Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Science (SIMATS), Chennai 600077, Tamil Nadu, India; tharaninano@gmail.com
 - ² Department of Biotechnology, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Science (SIMATS), Chennai 602105, Tamil Nadu, India; santhoshkumarj.sse@saveetha.com
 - ³ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia; kghanim@ksu.edu.sa
 - ⁴ Department of Environmental Biology, Foundation in Unam Sapientiam, Sapienza University of Rome, 00185 Rome, Italy; marcello.nicoletti@uniroma1.it
 - ⁵ Department of Microbiology V.S. Kiktenko, Institute of Medicine, Peoples Friendship University of Russia Named after Patrice Lumumba (RUDN University), Moscow 117198, Russia; sachivkina@yandex.ru
 - ⁶ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608002, Tamil Nadu, India
 - ⁷ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India
- * Correspondence: rajeshkumars.sdc@saveetha.com (R.S.); drgovind1979@gmail.com or dr.m.govindarajan@gcwk.ac.in (M.G.)



Citation: Shanmugam, R.; Munusamy, T.; Jayakodi, S.; Al-Ghanim, K.A.; Nicoletti, M.; Sachivkina, N.; Govindarajan, M. Probiotic-Bacteria (*Lactobacillus fermentum*)-Wrapped Zinc Oxide Nanoparticles: Biosynthesis, Characterization, and Antibacterial Activity. *Fermentation* **2023**, *9*, 413. <https://doi.org/10.3390/fermentation9050413>

Academic Editors: Yi-huang Hsueh and Nikos G. Chorianopoulos

Received: 25 March 2023

Revised: 20 April 2023

Accepted: 24 April 2023

Published: 26 April 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Recently, fabricated nanoparticles (NPs), which can efficiently penetrate biological systems, have found increased usage in the health and hygiene industries. Microbial enzymes and proteins have recently shown their potential to act as reducing agents for the production of NPs, thereby providing an alternative to physical and chemical methods. Not only is this approach efficient and cost-effective, but it also produces a minimal ecological footprint. In this study, zinc oxide nanoparticles (ZnO NPs) were synthesized using probiotic bacteria (*Lactobacillus fermentum*) as the reducing and capping agent. Several analytical methods, including Fourier transform infrared spectroscopy (FT-IR), scanning electron microscopy (SEM), X-ray diffraction analysis (XRD), ultraviolet–visible spectroscopy (UV–Vis), and atomic force microscopy (AFM), were used to analyze the produced ZnO NPs. The SEM analysis confirmed the spherical form of the nanoparticles and estimated their average size to be between 100 and 120 nm. FT-IR analysis verified that the ZnO NPs’ surfaces contained many functional groups. X-ray diffraction examination evidenced that the biogenically produced nanoparticles were crystalline. AFM analysis revealed that the nanoparticles’ size was about 90–100 nm. The maximum absorption peak, determined via a UV–visible spectrophotometer, was 510 nm. The synthesized ZnO NPs’ antimicrobial activity against various bacterial strains was tested, and the highest level of antimicrobial activity was noted against a *Vibrio harveyi* strain. The maximum concentration, namely, 20 mM of ZnO NPs, showed the highest antimicrobial activity. These observations indicate that the synthesized ZnO NPs possess remarkable antimicrobial potency. This method is an efficient, environmentally friendly, cost-effective approach for producing ZnO NPs that are useful for various biomedical applications.

Keywords: green synthesis; nanoparticles; fish pathogen; bio-medicinal application; *Vibrio harveyi*; *Lactobacillus fermentum*



Process and techno-economic analysis of bioethanol production from residual biomass of marine macroalgae *Ulva lactuca*



Kalavathy Gengiah^{a,1}, Naveenkumar Rajendran^{b,c,1}, Khalid A. Al-Ghanim^d,
Marimuthu Govindarajan^{e,f}, Baskar Gurunathan^{a,*}

^a Department of Biotechnology, St. Joseph's College of Engineering, Chennai 600119, Tamil Nadu, India

^b Biological Systems Engineering, University of Wisconsin–Madison, Madison, WI 53706, United States

^c Forest Products Laboratory, USDA Forest Service, Madison, WI 53726, United States

^d Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia


^e Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

HIGHLIGHTS

- Pretreatment of algal residue was optimized to improve the availability of carbohydrates.
- Enzymatic hydrolysis and fermentation of algal biomass were optimized.
- Bioethanol produced under optimized conditions was characterized using GC-MS
- Techno-economic analysis of bioethanol production was studied.
- The minimum selling price of bioethanol was 0.47 \$/kg.

GRAPHICAL ABSTRACT



ARTICLE INFO

Editor: Huu Hao Ngo

Keywords:

Residual biomass

Ulva lactuca

Bioethanol

Fermentation

Techno-economic analysis

ABSTRACT

In the present work, the residual biomass of the green seaweed *Ulva lactuca* was chosen as feedstock to undergo separate hydrolysis and fermentation process to produce bioethanol. The hydrolysis process was optimized for cellulase, biomass, temperature, and time conditions. The maximum yield of fermentable sugars was 13.48 mg/mL. The recovered hydrolysate was subjected to fermentation using *Saccharomyces cerevisiae*. The bioethanol produced was subjected to gas chromatography coupled mass spectrometry analysis to determine the presence of ethanol. The technical performance and economic feasibility of the bioethanol production from *U. lactuca* were evaluated using the lab-scale data obtained for optimized conditions. The plant capacity was 10 MT/day of bioethanol production. The plant's capital investment and annual operating cost were 3.18 M\$ and 0.86 M\$ respectively. The total annual revenue of the plant was 1.41 M\$. The minimum selling price of bioethanol was 0.47 \$/kg. The ROI, payback period, IRR and NPV of the plant were 16.99 %, 5.89 years, 11.57 % and 291,000 \$ respectively. The utilization of residual biomass for biofuels helps to develop an economic and environmentally sustainable plant.

1. Introduction

The economic development of many countries depends on fossil fuels consumption in transportation sector, industrial industries, and many other uses. In addition to the limited availability of resources, these fuels

* Corresponding author.

E-mail address: basg2004@gmail.com (B. Gurunathan).

¹ Contributed Equally.

Article

Green Synthesis of Copper Oxide Nanoparticles Using *Sesbania grandiflora* Leaf Extract and Their Evaluation of Anti-Diabetic, Cytotoxic, Anti-Microbial, and Anti-Inflammatory Properties in an In-Vitro Approach

Kanagavalli Ramasubbu ¹, Siddharth Padmanabhan ¹, Khalid A. Al-Ghanim ², Marcello Nicoletti ³,
Marimuthu Govindarajan ^{4,5,*}, Nadezhda Sachivkina ⁶ and Vijayarangan Devi Rajeswari ^{1,*}

¹ Department of Biomedical Sciences, School of Biosciences and Technology, Vellore Institute of Technology, Vellore 632014, TN, India; kanaga0497@gmail.com (K.R.); siddharthpadmanabhan50@gmail.com (S.P.)

² Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia; kghanim@ksu.edu.sa

³ Department of Environmental Biology, Sapienza University of Rome, 00185 Rome, Italy; marcello.nicoletti@uniroma1.it

⁴ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608002, TN, India

⁵ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, TN, India

⁶ Department of Microbiology V.S. Kiktenko, Peoples Friendship University of Russia (RUDN University), Moscow 117198, Russia; sachivkina@yandex.ru

* Correspondence: drgovind1979@gmail.com (M.G.); vdevirajeswari@vit.ac.in (V.D.R.)



Citation: Ramasubbu, K.; Padmanabhan, S.; Al-Ghanim, K.A.; Nicoletti, M.; Govindarajan, M.; Sachivkina, N.; Rajeswari, V.D. Green Synthesis of Copper Oxide Nanoparticles Using *Sesbania grandiflora* Leaf Extract and Their Evaluation of Anti-Diabetic, Cytotoxic, Anti-Microbial, and Anti-Inflammatory Properties in an In-Vitro Approach. *Fermentation* **2023**, *9*, 332. <https://doi.org/10.3390/fermentation9040332>

Academic Editor: Ronnie G. Willaert

Received: 5 March 2023

Revised: 20 March 2023

Accepted: 23 March 2023

Published: 27 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Green methods of synthesizing nanoparticles are safer than chemical and physical methods, as well as being eco-friendly and cost-efficient. In this study, we use copper oxide nanoparticles (CuO NPs) fabricated with *Sesbania grandiflora* (Sg) (Hummingbird tree) leaves to test the effectiveness of green synthesizing methods. The attained Sg-CuO NPs physical and optical nature is characterized by UV-Vis spectroscopy Differential Reflectance Spectroscopy (UV-Vis DRS), Fourier Transform Infra-Red spectroscopy (FTIR), X-ray Diffraction spectroscopy (XRD), Scanning Electron Microscope (SEM), and Energy Dispersive X-ray Analysis (EDAX). UV-Vis spectrum for Sg-CuO NPs revealed a peak at 410 nm. SEM images showed the aggregation of needle-shaped particles, at a size of 33 nm. The amylase and glucosidase enzymes were inhibited by the Sg-CuO NPs up to 76.7% and 72.1%, respectively, indicating a possible antihyperglycemic effect. Fabricated Sg-CuO NPs disclosed the excellent inhibition of DPPH-free radicle formation (89.7%) and repressed protein degradation (81.3%). The results showed that Sg-CuO NPs display good anti-bacterial activity against the gram-negative (*Escherichia coli* and *Pseudomonas aeruginosa*) and gram-positive (*Staphylococcus aureus*). Cytotoxicity of the Sg-CuO NPs was determined using an IC₅₀ of 37 µg/mL. Sg-CuO NPs have shown promising anti-diabetic, anti-oxidant, protein degradation-inhibiting, and anti-microbial properties. Our findings have shown that synthesized Sg-CuO NPs have biological activities that may be utilized to treat bacterial infections linked to hyperglycemia.





Keywords: nanotechnology; green synthesis; medicinal plant; antihyperglycemic; bioactivity; anticancer

1. Introduction

Nanotechnology focuses on synthesizing nanomaterial, and has diverse application, such as biomedicine, delivery of nutrients and drugs, imaging techniques, etc. Multiple methods are available for nanoparticle synthesizing, including chemical processes, sol-gel, laser ablation, electro biosynthesis, biological, and green synthesis. In the green synthesis method of nanoparticles, instead of chemicals, plant-mediated products are likely to play the role of reducing agents; green reducing agents lessen the toxic effects of chemicals in the

Article

Insecticidal Potential of *Matricaria chamomilla*'s Essential Oil and Its Components (E)- β -Farnesene, Germacrene D, and α -Bisabolol Oxide A against Agricultural Pests, Malaria, and Zika Virus Vectors

Khalid A. Al-Ghanim ^{1,*}, Kaliyamoorthy Krishnappa ², Jeganathan Pandiyan ², Marcello Nicoletti ³, Baskar Gurunathan ⁴ and Marimuthu Govindarajan ^{5,6,*}

¹ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

² Department of Zoology and Wildlife Biology, A.V.C. College, Mannampandal, Mayiladuthurai 609 305, Tamil Nadu, India

³ Department of Environmental Biology, Foundation in Unam Sapientiam, Sapienza University of Rome, 00185 Rome, Italy; marcello.nicoletti@uniroma1.it

⁴ Department of Biotechnology, St. Joseph's College of Engineering, Chennai 600 119, Tamil Nadu, India

⁵ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

⁶ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

* Correspondence: kghanim@ksu.edu.sa (K.A.A.-G.); drgovind1979@gmail.com or dr.m.govindarajan@gcw.ac.in (M.G.)



Citation: Al-Ghanim, K.A.; Krishnappa, K.; Pandiyan, J.; Nicoletti, M.; Gurunathan, B.; Govindarajan, M. Insecticidal Potential of *Matricaria chamomilla*'s Essential Oil and Its Components (E)- β -Farnesene, Germacrene D, and α -Bisabolol Oxide A against Agricultural Pests, Malaria, and Zika Virus Vectors. *Agriculture* **2023**, *13*, 779. <https://doi.org/10.3390/agriculture13040779>

Academic Editor: Azucena González Coloma

Received: 20 February 2023

Revised: 24 March 2023

Accepted: 27 March 2023

Published: 28 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Agricultural and medical insect pests are damaging edible crops, spreading diseases, and harming non-target fauna and flora. Prominent polyphagous insect pests harass farmers in the agronomy sector, causing uncountable revenue corrosion. Ecofriendly phytopesticides can avoid the consequences of the bulk usage of synthetic chemicals. In this study, the toxic effect on third-instar larvae of four different insect species (*Spodoptera litura*, *Helicoverpa armigera*, *Aedes vittatus*, and *Anopheles subpictus*) and the bio-toxicity on non-target fauna (NTF) (*Stegodyphus sarasinorum* and *Gambusia affinis*) were evaluated using *Matricaria chamomilla* (Mc) essential oil (EO) and its major phytoconstituents (Mc-MPCs). GC-MS analysis of the studied *M. chamomilla* EO gathered 39 constituents, with (E)- β -Farnesene (24.3%), Germacrene D (9.4%), and α -Bisabolol oxide A (10.2%) accounting for the major constituents. Remarkable larval death was seen in *H. armigera* and *Ae. vittatus*. In addition, (E)- β -Farnesene, Germacrene D, and α -Bisabolol oxide A exhibited a relevant maximum toxic effect on the target pest's third-instar larvae. The bio-toxicity of *M. chamomilla* EO and Mc-MPCs was tested on terrestrial and aquatic NTF. The LC₅₀ values for *S. sarasinorum* and *G. affinis* ranged from 922.65 to 1750.49 μ g/mL. *M. chamomilla* EO and its MPCs evidenced prospective phytopesticidal efficiency on selected agricultural and medical insect pests.



Keywords: phytoconstituents; pesticide; insecticide; larvae; non-target fauna; ecofriendly

1. Introduction

Globally, agri-pests play a pivotal role in the deterioration of several high-value agri-products. Due to human activities, vector proliferation in tropical and subtropical environments has dramatically increased [1,2]. Among arthropods, mosquitoes are massive blood-sucking vectors that cause several public health problems. The effects of blood-sucking vectors on the human population have increased yearly [3–6] in terms of death and morbidity. Vectors are a major cause of significant, worrying economic problems in many parts of the world, as well as are responsible for transmitting a wide variety of infectious illnesses to humans and other species that rely on humans for their vascular system [7]. *Spodoptera litura* (Fab.) (*Cotton leafworm*) and *Helicoverpa armigera* Hubner (*Cotton bollworm*)

Article

Exopolysaccharides-Mediated ZnO Nanoparticles for the Treatment of Aquatic Diseases in Freshwater Fish *Oreochromis mossambicus*

Muthukumar Abinaya¹, Sathappan Shanthi¹, Jesudasan Palmy² , Khalid A. Al-Ghanim³, **Marimuthu Govindarajan**^{4,5}  and Baskaralingam Vaseeharan^{1,*}

- ¹ Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th Floor, Burma Colony, Karaikudi 630004, Tamil Nadu, India; govindamalmuthukumar06@gmail.com (M.A.); shanthisathappan@gmail.com (S.S.)
 - ² Poultry Production and Product Safety Research Unit, ARS, USDA, Center of Excellence for Poultry Science, University of Arkansas, 1260 W Maple St., Fayetteville, AR 72701, USA; palmy;jesudhasan@usda.gov
 - ³ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia; kghanim@ksu.edu.sa
 - ⁴ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608002, Tamil Nadu, India; drgovind1979@gmail.com
 - ⁵ Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India
- * Correspondence: vaseeharanb@gmail.com

Abstract: Bacterial fish disease outbreaks are a key concern for aquaculture. Complementary feed additives such as immunostimulants can serve as an ideal solution for disease prevention. Herein, we scrutinized the efficacy of exopolysaccharides (EPSs) from probiotic *Bacillus licheniformis* and EPS-mediated zinc oxide nanoparticles (EPS-ZnO NPs) for a diet to evaluate growth parameters, antioxidant enzyme activities, and immune stimulation together with disease resistance against *Aeromonas hydrophila* and *Vibrio parahaemolyticus* in Mozambique tilapia *Oreochromis mossambicus*. Fish were separated into seven groups, with six experimental groups fed with EPS and EPS-ZnO NPs at 2, 5, and 10 mg/g and a control fed a basal diet. The fish ingesting feed supplemented with EPS and EPS-ZnO NPs at 10 mg/g showed improved growth performance. Cellular and humoral-immunological parameters were tested in serum and mucus after 15 and 30 days of feeding. These parameters were substantially enriched with a 10 mg/g diet ($p < 0.05$) of EPS and EPS-ZnO NPs in comparison with the control. Furthermore, the EPS and EPS-ZnO NP supplemental diet actively enhanced the antioxidant response (glutathione peroxidase, superoxide dismutase, and catalase). In addition, the supplemental diet of EPS and EPS-ZnO NPs lowered the death rate and improved the disease resistance of *O. mossambicus* following assessment with *A. hydrophila* and *V. parahaemolyticus* at 50 μ L. Hence, the overall results suggest that the supplemental diet of EPS and EPS-ZnO NPs might be used to ensure aquaculture feed additives.

Keywords: *Bacillus licheniformis*; zinc nanoparticle; immune parameters; antioxidant response; growth performance; aquatic pathogens



Citation: Abinaya, M.; Shanthi, S.; Palmy, J.; Al-Ghanim, K.A.; Govindarajan, M.; Vaseeharan, B. Exopolysaccharides-Mediated ZnO Nanoparticles for the Treatment of Aquatic Diseases in Freshwater Fish *Oreochromis mossambicus*. *Toxics* **2023**, *11*, 313. <https://doi.org/10.3390/toxics11040313>

Academic Editor: Antonia Concetta Elia

Received: 18 February 2023

Revised: 22 March 2023

Accepted: 24 March 2023


Published: 27 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Aquaculture is the most important agriculture industry, and it has grown far faster than other food sectors to satisfy fish market requirements. Around 15% of the protein ingested by three billion people comes from fish. In emerging nations such as South Asia, Southeast Asia, China, and Japan, fish is considered a primary nutrition source, creating a massive demand for intensive farming. With the expansion of industry and increased yields, intensive culture practices have been adopted that lead to environmental stress and disease prevalence, which are considered essential challenges for the sector [1–4]. Bacterial



Dual strategy for bioconversion of elephant grass biomass into fermentable sugars using *Trichoderma reesei* towards bioethanol production

Jayaraj Iyyappan^a, Ravichandran Pravin^b, Khalid A. Al-Ghanim^c, **Marimuthu Govindarajan^{d,e}**, Marcello Nicoletti^f, Gurunathan Baskar^{b,*}

^a Department of Biotechnology, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences (SIMATS), Thandalam, Chennai 602107, India

^b Department of Biotechnology, St. Joseph's College of Engineering, Chennai 600119, Tamil Nadu, India

^c Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^d Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar-608 002, Tamil Nadu, India


^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

^f Department of Environmental Biology, Sapienza University of Rome, Rome 00185, Italy

HIGHLIGHTS

- Lignolytic enzymes were produced by recombinant *Trichoderma reesei*.
- Bidelignification assisted with NiO nanoparticles has removed 54% of lignin.
- Hydrolytic enzymes were produced by *Trichoderma reesei* with enhancer.
- Dual strategy has supported bioconversion of elephant grass into ethanol.

GRAPHICAL ABSTRACT



ARTICLE INFO

Keywords:

Bioethanol
Elephant grass
Bidelignification
Hydrolytic activity
Nanoparticle

ABSTRACT

In this study, bidelignification and enzymatic hydrolysis of elephant grass were performed by recombinant and native strain of *Trichoderma reesei*, respectively. Initially, r*T. reesei* displaying Lip8H and MnP1 gene was used for bidelignification with NiO nanoparticles. Saccharification was performed by combining hydrolytic enzyme produced with NiO nanoparticles. Elephant grass hydrolysate was used for bioethanol production using *Kluyveromyces marxianus*. Maximum lignolytic enzyme production was obtained with 15 µg/L of NiO nanoparticles and initial pH of 5 at 32 °C. Subsequently, about 54% of lignin degradation was achieved after 192 h. Hydrolytic enzymes showed elevated enzyme activity and resulted in 84.52 ± 3.5 g/L of total reducing sugar at 15 µg/mL NiO NPs. About 14.65 ± 1.75 g/L of ethanol was produced using *K. marxianus* after 24 h. Thus, dual strategy employed for conversion of elephant grass biomass into fermentable sugar and subsequent biofuel production could become potential platform for commercialization.

* Corresponding author.

E-mail address: basg2004@gmail.com (G. Baskar).

<https://doi.org/10.1016/j.biortech.2023.128804>

Received 23 January 2023; Received in revised form 22 February 2023; Accepted 24 February 2023

Available online 26 February 2023

0960-8524/© 2023 Elsevier Ltd. All rights reserved.



Investigations on evaluation of marine macroalgae *Dictyota bartayresiana* oil for industrial scale production of biodiesel through technoeconomic analysis

Pravin Ravichandran^a, Naveenkumar Rajendran^{b,c}, Khalid A. Al-Ghanim^d,
Marimuthu Govindarajan^{e,f}, Baskar Gurunathan^{a,*}

^a Department of Biotechnology, St. Joseph's College of Engineering, Chennai 600119, Tamil Nadu, India

^b Biological Systems Engineering, University of Wisconsin–Madison, Madison, WI 53706, USA

^c Forest Products Laboratory, USDA Forest Service, Madison, WI 53726, USA

^d Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia


^e Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

HIGHLIGHTS

- Biodiesel production from marine macroalgae *Dictyota bartayresiana* oil was studied.
- CaO nanocatalyst was synthesized using waste collected from building demolition site.
- The reaction parameters for efficient production of biodiesel was optimized.
- Investigation on techno-economic aspects was studied for biodiesel production.
- Sensitivity analysis was used to study the uncertainty of biodiesel production process.

GRAPHICAL ABSTRACT



ARTICLE INFO

Keywords:

Dictyota bartayresiana
Marine macroalgae
Biodiesel
Response surface methodology
Process optimization
Techno-economic analysis

ABSTRACT

The investigation on utilizing macroalgae for industrial scale biodiesel production is an imperative action needed for commercialization. In the present research work, the biooil from marine macroalgae *Dictyota bartayresiana* was used for biodiesel production using calcium oxide nanocatalyst synthesized using waste collected from building demolition site. The optimization results obtained were the calcination temperature 573 °C, concentration of catalyst 5.62%, methanol to oil molar ratio 14.36:1, temperature 55.7 °C and time 67.57 min for the transesterification with the biodiesel yield of 89.6%. The techno-economic aspects of biodiesel production were investigated for 20 MT/batch. The return on investment and internal rate of return from the biodiesel production plant was found to be 25.39% and 31.13% respectively. The plant payback period was about 3.94 years with a positive NPV value of about 14,053,000 \$/yr. Thus, *Dictyota bartayresiana* biomass can be efficiently used for the sustainable production of biodiesel.

* Corresponding author.

E-mail address: basg2004@gmail.com (B. Gurunathan).

<https://doi.org/10.1016/j.biortech.2023.128769>

Received 14 January 2023; Received in revised form 17 February 2023; Accepted 18 February 2023

Available online 24 February 2023

0960-8524/© 2023 Elsevier Ltd. All rights reserved.

Article

Biochemical, Genotoxic and Histological Implications of Polypropylene Microplastics on Freshwater Fish *Oreochromis mossambicus*: An Aquatic Eco-Toxicological Assessment

Jeyaraj Jeyavani ¹, Ashokkumar Sibiya ¹, Thambusamy Stalin ², Ganesan Vigneshkumar ², Khalid A. Al-Ghanim ³, Mian Nadeem Riaz ⁴, **Marimuthu Govindarajan** ^{5,6}  and Baskaralingam Vaseeharan ^{1,*}

¹ Biomaterials and Biotechnology in Animal Health Lab., Department of Animal Health and Management, Alagappa University, Karaikudi 630003, Tamil Nadu, India

² Department of Industrial Chemistry, Alagappa University, Karaikudi 630003, Tamil Nadu, India

³ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

⁴ Texas A&M University, College Station, TX 77843, USA

⁵ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608002, Tamil Nadu, India; drgovind1979@gmail.com or dr.m.govindarajan@gcwk.ac.in

⁶ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

* Correspondence: vaseeharanb@gmail.com



Citation: Jeyavani, J.; Sibiya, A.; Stalin, T.; Vigneshkumar, G.; Al-Ghanim, K.A.; Riaz, M.N.; Govindarajan, M.; Vaseeharan, B. Biochemical, Genotoxic and Histological Implications of Polypropylene Microplastics on Freshwater Fish *Oreochromis mossambicus*: An Aquatic Eco-Toxicological Assessment. *Toxics* **2023**, *11*, 282. <https://doi.org/10.3390/toxics11030282>

Academic Editor: Jordan Jun Chul Park

Received: 9 February 2023

Revised: 14 March 2023

Accepted: 16 March 2023

Published: 19 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: In recent years, polypropylene microplastic has persisted in freshwater ecosystems and biota, forming ever-growing threats. This research aimed to prepare polypropylene microplastics and evaluate their toxicity to the filter feeder *Oreochromis mossambicus*. In this research, fish were given a dietary supplement of polypropylene microplastics at 100, 500, and 1000 mg/kg for acute (96 h) and sub-acute (14 days) durations to assess toxic effects on liver tissues. FTIR results revealed the presence of polypropylene microplastic in their digestion matter. The ingestion of microplastics in *O. mossambicus* led to fluctuations in homeostasis, an upsurge in reactive oxygen species (ROS) levels, an alteration in antioxidant parameters, including superoxide dismutase (SOD), catalase (CAT), glutathione-S-transferase (GST), and glutathione peroxidase (GPx); a promotion in the oxidation of lipid molecules; and a denaturation in the neurotransmitter enzyme acetylcholinesterase (AChE). Our data indicated that sustained exposure to microplastics (14 days) produced a more severe threat than acute exposure (96 h). In addition, higher apoptosis, DNA damage (genotoxicity), and histological changes were found in the liver tissues of the sub-acute (14 days) microplastics-treated groups. This research indicated that the constant ingestion of polypropylene microplastics is detrimental to freshwater environments and leads to ecological threats.




Keywords: polypropylene microplastics; freshwater ecosystem; ever-growing threats; liver tissues; food; genotoxicity

1. Introduction

Microplastics may pose a hazard to a variety of ecosystems, making them a potential contaminant. Due to their diminutive size, they are readily diffused and carried throughout all aquatic environments, including sediment, the surface, and other zones such as the littoral, limnetic, profundal, euphotic, and benthic [1–3], and even in aquatic creatures [4]. In general, biota accidentally ingests microplastics, via oral, and gill, which cause obstruction or irritation of the digestive system due to their accumulation. Accumulated microplastics were transported into different internal organs via circulatory body fluids. Which triggers the dysfunction or alteration in endocrine and metabolic pathways, leads to oxidative stress, cell necrosis, finally inducing cell apoptosis, and aids the organism's death [5,6]. Microplastics and their impacts were reported in algae, zooplankton, fish, marine reptiles, sea birds, and aquatic mammals [7–11]. Among these, fish is a crucial predator and is

Article

Fishing Cat Scats as a Biomonitoring Tool for Toxic Heavy Metal Contamination in Aquatic Ecosystems

Thirupati Lakshmi Harika ¹, Khalid A. Al-Ghanim ², Mian Nadeem Riaz ³, Kaliyamoorthy Krishnappa ¹ , Jeganathan Pandiyan ¹  and **Marimuthu Govindarajan** ^{4,5,*} 

¹ Department of Zoology and Wildlife Biology, A.V.C. College, Mannampandal, Mayiladuthurai 609 305, Tamil Nadu, India

² Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

³ 2476 TAMU, Texas A&M University, College Station, TX 77843, USA

⁴ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

⁵ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

* Correspondence: drgovind1979@gmail.com

Abstract: Mangrove forest is one of the productive ecosystems that provide essential habitats for various fauna as breeding and feeding drives. However, heavy metal pollution in the mangrove forest has led to severe health problems for several aquatic species. Biomonitoring of metals using a nondestructive method is an emerging technique. Scats of the fishing cat (*Prionailurus viverrinus*) were collected from five locations in the Godavari estuary mangrove habitats, Coringa Wildlife Sanctuary, Andhra Pradesh, India, to determine the level of various metals. An opportunistic method was applied to collect scats in the mangrove forest. Six scat samples were collected from each of the sampling sites. The following prey species, such as crabs, fishes, birds, rodents, plants, plastics, and unidentifiable prey matters, were found in the scats. Select metals, such as chromium (Cr), copper (Cu), and lead (Pb) were analyzed from the scats of the fishing cat since they intensively influence the physiology and behavior of top predators. The concentration of Cu in fishing cat scats was higher than the other two metals assessed. Metals showed statistically substantial variation across locations ($p < 0.05$). According to the current study, heavy metals may significantly threaten the fishing cat in the Coringa Wildlife Sanctuary. The fishing cat is a vulnerable species in accordance with the ICUN categories. Due to pollution and other human pressures, the fishing cat may soon be categorized as a threatened or endangered species; the research advises that authorities should prioritize the protection of the vulnerable species of the fishing cat from the Coringa Wildlife Sanctuary, Andhra Pradesh, India.

Keywords: heavy metal; toxicity; scats; prey species; mangrove forests; carnivores; conservation



Citation: Harika, T.L.; Al-Ghanim, K.A.; Riaz, M.N.; Krishnappa, K.; Pandiyan, J.; Govindarajan, M. Fishing Cat Scats as a Biomonitoring Tool for Toxic Heavy Metal Contamination in Aquatic Ecosystems. *Toxics* **2023**, *11*, 173. <https://doi.org/10.3390/toxics11020173>

Academic Editor: Luis Alberto Henríquez-Hernández

Received: 18 January 2023

Revised: 31 January 2023

Accepted: 10 February 2023

Published: 13 February 2023






Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Metals are a class of non-biodegradable contaminants that cause extensive ecological damage by accumulating in aquatic ecosystems and posing substantial health risks to various organisms [1]. Evaluating heavy metals in the aquatic environment is crucial to comprehend the toxicity of metals and their effect on aquatic life. Countless animal species rely on mangrove forests for sustenance and shelter, making them an essential aquatic environment. Mangrove forests have been shown to play an important role in the lives of many animal species [2]. The eastern coast of southern India is home to one of India's biggest mangrove forests, the Godavari forest. The Godavari mangrove forest has major ecological and biological functions in its estuary locations [3]. Due to its importance as a marine ecosystem, the Godavari mangroves have been declared as a "Coringa Wildlife Sanctuary."

Article

Green Synthesis, Characterization and Bioactivity of *Mangifera indica* Seed-Wrapped Zinc Oxide Nanoparticles

Shanmugam Rajeshkumar ^{1,*}, Royapuram Parthasarathy Parameswari ¹, Dayalan Sandhiya ²,
Khalid A. Al-Ghanim ³, Marcello Nicoletti ⁴ and Marimuthu Govindarajan ^{5,6,*}

¹ Department of Pharmacology, Saveetha Dental College and Hospitals, Saveetha University, SIMATS, Chennai 600077, TN, India

² Department of Medicinal and Applied Chemistry, Kaohsiung Medical University, Kaohsiung City 80708, Taiwan

³ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

⁴ Department of Environmental Biology, Sapienza University of Rome, 00185 Rome, Italy

⁵ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608002, TN, India

⁶ Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, TN, India

* Correspondence: ssrajeshkumar@hotmail.com (S.R.); drgovind1979@gmail.com (M.G.)

Abstract: In the realm of nanoparticles, metal-based nanoparticles have traditionally been regarded as the pioneering category. Compared to other nanoparticles, zinc oxide nanoparticles have several advantages, including optical and biological properties, which provide them a significant competitive advantage in clinical and biological applications. In the current investigation, we used an aqueous *Mangifera indica* seed extract to synthesize nanoparticles of zinc oxide (ZnO NPs). UV-Vis spectroscopy, Fourier transform infrared spectroscopy analysis, atomic force spectroscopy, X-ray diffraction, scanning electron microscopy, and transmission electron microscopy were used to characterize the synthesized ZnO NPs. The nanoparticles were assessed for their potential to inhibit bacterial growth and protect cells from free radical damage. According to the current study's findings, zinc oxide nanoparticles that had been modified with the aid of mango seeds were very efficient in preventing the development of the tested bacteria and were also powerful antioxidants.

Keywords: nanotechnology; ZnO NPs; biofabrication; antioxidant; antibacterial activity



Citation: Rajeshkumar, S.; Parameswari, R.P.; Sandhiya, D.; Al-Ghanim, K.A.; Nicoletti, M.; Govindarajan, M. Green Synthesis, Characterization and Bioactivity of *Mangifera indica* Seed-Wrapped Zinc Oxide Nanoparticles. *Molecules* **2023**, *28*, 2818. <https://doi.org/10.3390/molecules28062818>

Academic Editor: Maria Camilla Bergonzi

Received: 29 January 2023

Revised: 16 March 2023

Accepted: 18 March 2023

Published: 21 March 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Nanotechnology has been gaining attention recently as a potential platform for future growth in several fields. Nanotechnology has drawn significant attention in the healthcare, engineering and food industries by offering novel prospects in the respective fields. In particular, theranostics, a cutting-edge combination system of therapeutics and diagnostics, utilize nanotechnology principles for target-specific drug delivery and enhanced bioavailability of active pharmaceutical ingredients [1,2]. The field of nanotechnology deals with various synthesis methods, particle size reformations and structural variations of nanoparticles. Nanoparticles are nanosized materials ranging in size <100 nm with high thermal stability, high surface-to-volume ratio, high electrical, mechanical, optical as well as magnetic properties [3].

In the last decade, the use of nanoparticles has been the most significant archetype advancement in engineering, medicine and technology [4]. Nanoparticles may be classified as organic and inorganic nanoparticles. While metals and metal-derived oxide nanoparticles come under the inorganic nanoparticles classification, organic nanoparticles include solid lipid nanoparticles, polymeric nanoparticles, lipid-based nanocarriers, liposomes and carbon-based nanomaterials [5]. Metal nanoparticles are promise for site-specific drug administration, clinical diagnostics, bio-imaging, dental implants, and biomedicine due

Research Article

Mahendrakumar Mani, Aruna Sharmili Sundararaj, Khalid A. Al-Ghanim, Shiny Punalur John, Kuppusamy Elumalai*, Marcello Nicoletti, and **Marimuthu Govindarajan***

Rapid synthesis of copper nanoparticles using *Nepeta cataria* leaves: An eco-friendly management of disease-causing vectors and bacterial pathogens

<https://doi.org/10.1515/gps-2023-0022>

received February 06, 2023; accepted May 07, 2023

Abstract: Insecticides kill mosquitoes but damage other animals including humans. Eco-friendly metal nanoparticles may be a plant-based pesticide for vector control. Here, the copper nanoparticles (Cu NPs) synthesized from *Nepeta cataria* leaves have been investigated for their antibacterial and larvicidal properties. Fourier transforms infrared spectroscopy demonstrated that biochemicals reduced and stabilized nanoparticles by shifting peaks from 1,049 to 1,492 cm^{-1} , and ultraviolet–visible spectrometry proved that produced Cu NPs had a peak at 550 nm. Transmission electron microscopic and scanning electron microscope showed that the particles are spherical and 23–29 nm in size. X-ray diffraction analysis

shows that Cu NPs are crystalline. At a 100 $\mu\text{g}\cdot\text{mL}^{-1}$ concentration, Cu NPs exhibited a higher percentage of inhibition in the order of *Escherichia coli* \gg *Enterococcus faecalis* \gg *Staphylococcus aureus*. The lethal concentration (LC_{50}) of the Cu NPs against the larvae of *Aedes aegypti*, *Anopheles stephensi*, and *Culex quinquefasciatus* was determined to be 60.63, 56.58, and 54.32 $\mu\text{g}\cdot\text{mL}^{-1}$, respectively. This ground-breaking study describes the biological production of Cu NPs utilizing *N. cataria* leaf extract for the first time. Based on these findings, the bio-synthesized Cu NPs and the aqueous extract of *N. cataria* may provide a potential alternative method for managing these vector populations.

Keywords: biosynthesis, copper nanoparticles, larvicidal efficacy, microbial pathogens, vector control

* Corresponding author: Kuppusamy Elumalai, Department of Advanced Zoology and Biotechnology, Government Arts College for Men (Autonomous), Nandanam, Chennai, 600035, Tamil Nadu, India, e-mail: professorelumalai@gmail.com

* Corresponding author: Marimuthu Govindarajan, Department of Zoology, Unit of Vector Control, Phytochemistry and Nanotechnology, Annamalai University, Annamalainagar 608002, Chidambaram, Tamil Nadu, India; Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

Mahendrakumar Mani: Department of Biotechnology, Guru Nanak College (Autonomous), Velachery, Chennai, 600042, Tamil Nadu, India

Aruna Sharmili Sundararaj: Department of Biotechnology, Stella Maris College (Autonomous), Chennai, 600086, Tamil Nadu, India

Khalid A. Al-Ghanim: Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

Shiny Punalur John: Department of Chemical Engineering, Indian Institute of Technology - Madras, Chennai, Tamil Nadu 600036, India

Marcello Nicoletti: Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

1 Introduction

Nanotechnology deals with controlling and manipulating matter at the molecular level between 1 and 100 nm. This field leads to the development of discrete properties collated with those related to atoms or molecules or bulk behaviour [1]. In the expanding field of nanotechnology, physical, chemical, and even ecologically friendly biological processes might all be employed to create nanomaterials. Chemical methods of generating nanoparticles (NPs) need the unrestricted use of harmful compounds, polluting the environment [2]. The biologically mediated processes can be conceded using bacteria, fungi, yeast, actinomycetes, and viruses [3–5]. Green synthesis based on the extraction of plant extracts has been reported in the literature [6–8]. Plant-based approaches are one of the most cost-effective, safe, and environmentally friendly ways to produce NPs [9]. The plant-mediated biosynthesized NPs are also steadier and can be synthesized faster than other synthesis methods [10].



Response of hepatic biochemical parameters and neurotoxicity to carbamazepine and ibuprofen in *Oreochromis mossambicus*

Ashokkumar Sibiya · Jeyaraj Jeyavani · Manoharan Saravanan · Mohammed F. Albeshr · Marcello Nicoletti · **Marimuthu Govindarajan** · Baskaralingam Vaseeharan

Received: 2 September 2022 / Accepted: 26 January 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract Globally, the prevalence and pollution of pharmaceutical drugs in aquatic environments have been steadily increasing. This study sought to evaluate the effects of 14 days of exposure to environmental-relevant doses (ibuprofen 0.5, 5, and 50 µg/L, and carbamazepine 0.005, 1, and 10 µg/L) of the non-steroidal anti-inflammatory drugs ibuprofen and carbamazepine in the freshwater fish *Oreochromis mossambicus*. The results showed a significant ($P < 0.05$)

decrease in *O. mossambicus* superoxide dismutase, catalase, biotransformation enzymes, glutathione-S-transferase, glutathione peroxidase, oxidative stress lipid peroxidation, protein carbonyl activity, cellular damage metallothionine, reduced glutathione, immunological activities, and respiratory burst activity. Consequently, the acquired data revealed that *O. mossambicus* treated with ibuprofen and carbamazepine shows more significant alterations in metabolic depression, biochemical parameters, and oxidative stress. In addition, increased neurotoxic effects were observed in ibuprofen and carbamazepine treated *O. mossambicus*.

A. Sibiya · J. Jeyavani · M. Saravanan · B. Vaseeharan (✉)
Nano Biosciences and Nanopharmacology Division,
Biomaterials and Biotechnology in Animal Health Lab,
Department of Animal Health and Management, Science
Campus 6th Floor, Alagappa University, Karaikudi,
630004 Tamil Nadu, India
e-mail: vaseeharanb@gmail.com; vaseeharanb@
alagappauniversity.ac.in

M. F. Albeshr
Department of Zoology, College of Science, King Saud
University, Riyadh 11451, Saudi Arabia

M. Nicoletti
Department of Environmental Biology, Sapienza
University of Rome, 00185 Rome, Italy

M. Govindarajan
Unit of Mycology and Parasitology, Department
of Zoology, Annamalai University, Annamalaiagar,
608 002 Tamil Nadu, India

M. Govindarajan
Unit of Natural Products and Nanotechnology,
Department of Zoology, Government College for Women
(Autonomous), Kumbakonam, 612 001 Tamil Nadu, India



Keywords Drugs · *Oreochromis mossambicus* ·
Oxidative stress · Neurotoxicity

Introduction

Human, veterinary, agricultural, and aquaculture sectors, as well as human health as a whole, have benefited from the pharmaceutical industry's creation of a vast array of drugs. (Saravanan et al. 2012). The aquatic environment is regularly exposed to a range of medications (parental compounds) and their by-products (metabolites) because of the widespread usage and large-scale manufacture of pharmaceuticals (parental compounds) (metabolites). These drugs may enter the local aquatic environment through pharmaceutical companies, hospital effluent, and domestic

Article

Synthesis of Tungsten Oxide Nanoflakes and Their Antibacterial and Photocatalytic Properties

Sarinthip Thanakkasaranee ¹, Gopinath Kasi ^{1,*}, Sivasamy Kadiravan ², Ayyakannu Arumugam ³, Khalid A. Al-Ghanim ⁴, Mian Nadeem Riaz ⁵ and Marimuthu Govindarajan ^{6,7,*}

¹ Division of Packaging Technology, School of Agro-Industry, Faculty of Agro-Industry, Chiang Mai University, Chiang Mai 50100, Thailand

² Department of Chemistry, Swami Dayananda College of Arts and Science, Manjakkudi, Tiruvarur 612 610, Affiliated to Bharathidasan University, Tiruchirappalli 620 024, Tamil Nadu, India

³ Department of Botany, Alagappa University, Karaikudi 630 003, Tamil Nadu, India

⁴ Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

⁵ 2476 TAMU, Texas A&M University, College Station, TX 77843, USA

⁶ Unit of Mycology and Parasitology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India

⁷ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

* Correspondence: gloriousnanogopi@gmail.com (G.K.); drgovind1979@gmail.com (M.G.)

Abstract: This current work revealed a single-step fabrication of tungsten oxide nanoflakes (WO₃ NFs) with the help of *Terminalia arjuna* bark extract. Bioactive phytoconstituents of *T. arjuna* bark extract were involved in the nucleation process and promoted the material crystalline growth in a particular direction. The as-prepared sample thermal decomposition was analyzed by TG/DTG. The as-prepared sample was annealed at 300 °C for 2 h, and the annealed sample was characterized by UV-Vis-DRS, FTIR, Raman, XRD, SEM, EDX, and TEM. Synthesized WO₃ samples showed a monoclinic phase of the flake-like structure with lengths of 25~230 nm and diameters of 25~120 nm. The WO₃ NFs were evaluated against *S. aureus* and *E. coli*. Over 3 mg concentrations of WO₃ NFs outperform the positive control in antibacterial activity. The pseudo-first-order kinetics of the WO₃ NFs enhanced the photocatalytic performance of methylene blue (MB). These results prove that WO₃ NFs have sustainable performance in antibacterial and MB degradation applications.

Keywords: *Terminalia arjuna*; bark extract; nano-flakes; reactive oxygen species; photodegradation



Citation: Thanakkasaranee, S.; Kasi, G.; Kadiravan, S.; Arumugam, A.; Al-Ghanim, K.A.; Riaz, M.N.; Govindarajan, M. Synthesis of Tungsten Oxide Nanoflakes and Their Antibacterial and Photocatalytic Properties. *Fermentation* **2023**, *9*, 54. <https://doi.org/10.3390/fermentation9010054>

Academic Editor: Nadezhda Sachivkina

Received: 28 October 2022

Revised: 2 January 2023

Accepted: 3 January 2023

Published: 9 January 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Bacterial infections have recently been identified as a major global health threat [1]. For the most part, organic antibiotics have been utilized to treat human bacterial diseases, resulting in bacterial pathogens becoming resistant to antibiotics. However, novel classes of antibiotics slow their further development [2]. On the other hand, the prime source of all living organisms for sustainable development needs is water. The discharges of industrial effluents, especially from the textile industry, such as azo dye, phenyl matter, and toxic organic substances resistant to degradation and decomposition by biological treatment, cause severe water pollution. Chemical markers and biological staining procedures employ methylene blue (MB), a cationic dye. Due to the high chromatic nature and poor biodegradability, it has been complicated to process degradability. More importantly, it causes various harmful effects on human health care, such as diarrhea, vomiting, severe headache, and methemoglobinemia-like syndrome. In addition, it significantly affects the waterbody phytoplankton photosynthesis and microorganism metabolic activity in the water environment [3–6]. Photocatalyst is one of the novel methods to prevent water pollution. Two-dimensional (2D) nanomaterials are efficient catalysts for solving bacterial pathogens, mosquito vectors, and environmental remediation problems. Nowadays,



Synthesis of zinc oxide nanoparticles using *Vigna mungo* seed husk extract: An enhanced antibacterial, anticancer activity and eco-friendly bio-toxicity assessment on algae and zooplankton

Ramachandran Ishwarya^{a,b}, Govindan Tamilmani^b, Rengarajan Jayakumar^c,
 Mohammed F. Albeshr^d, Shahid Mahboob^d, Duaa Shahid^e, Mian Nadeem Riaz^f,
 Marimuthu Govindarajan^{g,h}, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Mandapam Regional Centre, ICAR - Central Marine Fisheries Research Institute, Mandapam, 623 520, Tamil Nadu, India

^c ICAR - Central Institute of Brackishwater Aquaculture (CIBA), #75, Santhome High Road, MRC Nagar, Chennai, 600 028, Tamil Nadu, India

^d Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^e Hult International Business School, Cambridge, MA, 02141, USA

^f Texas A&M University, College Station, TX, USA

^g Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^h Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Vigna mungo
 Green synthesis
 Nanodrug: microalgae
 Copepod
 Breast cancer cell line

ABSTRACT

An ecological synthesis of zinc oxide nanoparticles (ZnONPs) from plant extracts has several applications in contemporary research owing to its enhanced pharmacological efficacy, lower toxicity, and cost-effectiveness. This study synthesized ZnONPs using the seed husk extracts of *Vigna mungo* (*Vm*) (Black gram). UV, XRD, FTIR, SEM, and TEM were used to analyze well-dispersed *Vm*-ZnONPs. Moreover, we examined the bactericidal efficacy of *Vm*-ZnONPs against aquatic Gram-negative pathogens. In addition, the breast cancer cell line MDA-MB-231 was utilized to conduct conventional apoptosis tests, fluorescent dye assays, and cytotoxicity of cancer cell line data, which revealed that *Vm*-ZnONPs had good anticancer activity against MDA-MB-231. Furthermore, the growth rate and biotoxicity of *Vm*-ZnONPs were investigated on three marine microalgae: *Chaetoceros calcitrans*, *Isochrysis galbana*, and *Nannochloropsis oculata*. The concentration of *Vm*-ZnONPs significantly improved the growth rate of three algae species. At 96 h, *I. galbana* grew at a faster rate than *N. oculata* and *C. calcitrans* in response to *Vm*-ZnONPs. Also, the effects of biotoxicity on *Vm*-ZnONPs were validated using two animal models, including zooplankton rotifers and copepods. *Brachionus plicatilis* and *Parvocalanus crassirostris* treated with *Vm*-ZnONPs exhibited low toxicity, with LC₅₀ values of 62.87 and 71.73 µg/mL, respectively. The *Vm*-ZnONPs were highly identified by zooplankton rotifers (*B. plicatilis*), copepods (*P. crassirostris*), and no morphological or anatomical alterations verified their safety to the variety of fauna. This is the first research to examine the impact of *Vm*-ZnONPs on the marine ecosystem biotoxicity of three phytoplanktons and two zooplanktons.

1. Introduction

Biowaste materials have been recognized as a global ecological problem, and they have prompted the development of innovative pathways to create renewable, low-cost, and extensive therapeutic and aesthetic uses. *Vigna mungo* (Black gram) is the most important pulse crop in India; 25% of by-products (agro waste) are derived by grinding

black gram into dhal [1]. This seed husk is one of these agro-industrial by-products, and it is utilized as livestock feed; it has no economic value. It was discovered that the seed husk of *V. mungo* contains flavonoids, glycoflavones, flavonol glycosides, and compounds such as vitexin, leucodelphinidin, and delphinidin 3-glucoside [2]. In addition, the seed husk contains saponins, namely acetylsoyasaponin and soyasaponin [3].

ZnONPs have recently attracted much interest because of their

* Corresponding author.


E-mail addresses: vaseeharanb@gmail.com, vaseeharanb@alagappauniversity.ac.in (B. Vaseeharan).

<https://doi.org/10.1016/j.jddst.2022.104002>

Received 30 September 2022; Received in revised form 5 November 2022; Accepted 17 November 2022

Available online 18 November 2022

1773-2247/© 2022 Elsevier B.V. All rights reserved.



Acaricidal efficacy of silver nanoformulations of *Commiphora molmol* and *Zingiber officinale* against the camel Tick, *Hyalomma dromedarii* (Ixodida: Ixodidae)

M. Nabil^{a,*}, Hanem F. Khater^b, Abdelfattah Selim^c, Mohamed M. Baz^d,
Marimuthu Govindarajan^e, Hanan A.A. Taie^f, S. Negm^a

^a Department of Basic Engineering Sciences, Faculty of Engineering (Shoubra), Benha University, Cairo, Egypt

^b Department of Parasitology, Faculty of Veterinary Medicine, Benha University, Toukh 13736, Qalyubia, Egypt

^c Department of Animal Medicine (Infectious Diseases), Faculty of Veterinary Medicine, Benha University, Egypt

^d Department of Entomology, Faculty of Science, Benha University, Qalyubiya 13518, Egypt

^e Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^f Plant Biochemistry Department, Agricultural and Biology Research Institute, National Research Centre, 33 El-Bohouth St. (Former El-Tahrir St.), Dokki, Cairo 12622, Egypt

ARTICLE INFO

Keywords:

Nanoparticles
Myrrh
Resistance
Polyphenol
Flavonoid
Tannin

ABSTRACT

Ticks are blood-feeding ectoparasites transmitting many dangerous pathogens to humans and animals, leading to great economic losses. Emerging acaricide resistance has urged the use of eco-friendly acaricides. The novel *In vitro* application of the aqueous extracts of myrrh, *Commiphora molmol* (AME) and ginger, *Zingiber officinale* (AGE) and their corresponding silver nanoparticles (AgNPs), synthesized by subjecting myrrh and ginger extracts to laser ablation (NME and NGE, respectively), was evaluated against the camel tick, *Hyalomma dromedarii* by adult immersion bioassays. Moreover, spot-on treatments of infested cattle was done. AgNPs were characterized by Transmission Electron Microscopy and UV-visible spectrophotometry. The phytochemical profile showed that the used extracts contained phenols, gallic acid, flavonoids, and tannins. Fifteen days post-treatment (PT), 96 and 84.01% mortalities were observed PT with AME and AGE, respectively; 100% mortality was reached 7 and 9 days PT with 12% of NME and NGE, respectively. Three days PT, the LC₅₀ values of AME, AGE, NME, and NGE were 10.37, 12.81, 2.38, and 4.12%, respectively. The corresponding LT₅₀ values PT with 4% were 5.6 and 6.73, 2.25, and 3.56 days, respectively. Three days PT, AME, AGE, NME, and NGE reduced cattle-tick infestations by 54.45, 45.73, 100, and 100%, respectively. Ticks showed resistance against Deltamethrin (Butox®). This study demonstrated the novel acaricidal effect of myrrh and ginger and their silver nanoformulations through laser ablation, which increased the speed and efficacy of the aqueous extracts against *H. dromedarii*. Consequently, they could be produced as efficient eco-friendly pesticides after revealing their ecotoxicological profile.

1. Introduction

Ticks are important ectoparasites that transmit pathogens of medical and veterinary importance, which cause serious health issues to humans and domestic animals [1,2] leading to great economic losses [3]. Ticks are vectors of severe viral, bacterial, and parasitic diseases [4–9]. In addition to spreading diseases, ticks have direct effects on the health of livestock through their bites, which can be painful, cause blood loss, damage the skin, and make animals lose their appetite, which slows their growth [10,11]. Therefore, it is necessary to fight these pests to reduce the damage caused by them and limit their spread through the

application or use of the appropriate control strategies. Using of chemical pesticides in the control of ticks led contaminated meat and dairy products, polluted environment, and development of acaricide-resistant strains of ticks. Consequently, using eco-friendly acaricides is a potential solution to the resistance problem, which is what we seek in this current work [12,13].

Biorational pesticides are safe, cost-effective, and biodegradable [12–14]. Besides their antiparasitic effect [15,16], botanicals have ovicidal, adulticidal [17]; larvicidal [18,19], repellent, and deterrent effects [20]. Pesticidal activity of *Commiphora molmol* and *Araucaria heterophylla* methanol and hexane extracts against ectoparasites

* Corresponding author.

<https://doi.org/10.1016/j.inoche.2022.110229>

Received 18 July 2022; Received in revised form 9 November 2022; Accepted 16 November 2022

Available online 20 November 2022

1387-7003/© 2022 Elsevier B.V. All rights reserved.



Ingestion and impacts of water-borne polypropylene microplastics on *Daphnia similis*

Jeyaraj Jeyavani¹ · Ashokkumar Sibiya¹ · Narayanan Gopi¹ · Shahid Mahboob² · Khalid A. Al-Ghanim² · Fahad Al-Misned² · Zubair Ahmed² · Mian Nadeem Riaz³ · Balasubramanian Palaniappan⁴ · Marimuthu Govindarajan^{5,6} · Baskaralingam Vaseeharan¹

Received: 21 September 2021 / Accepted: 9 September 2022 / Published online: 22 September 2022
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

Abstract

Polypropylene microplastics are the leading contaminant in aquatic environments, although research on their toxicity remains scarce. The proposed research focuses on the harmful consequences of acute exposure to polypropylene microplastics in *Daphnia similis*. This work converts widely available polypropylene bags into microplastics using xylene. FTIR findings demonstrated the lack of xylene residue in the produced polypropylene microplastic particles, which were spherical and ranged in size from 11.86 to 44.62 μm (FE-SEM). The results indicate that acute exposure to polypropylene microplastics causes immobility in *D. similis*. Ingestion of microplastics enhances the generation of reactive oxygen species (ROS), as shown by biochemical studies. Due to the production of free radicals in *D. similis*, the antioxidant enzymes superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione-S-transferase (GST) and a non-antioxidant enzyme of reduced glutathione (GSH) and also oxidative stress effects in lipid (lipid peroxidation — LPO), protein (carbonyl protein — CP) were increased. Additionally, the amount of the neurotransmitter enzyme acetylcholinesterase (AChE) activity was decreased. These findings indicate that the accumulation of polypropylene microplastics in the bodies of filter-feeding organisms should aggravate toxicity in the freshwater environment.

Keywords Polypropylene microplastics · *Daphnia similis* · Acute exposure · Aquatic environments · Biochemical parameters

Responsible Editor: Bruno Nunes

✉ Baskaralingam Vaseeharan
vaseeharanb@gmail.com;
vaseeharanb@alagappauniversity.ac.in

¹ Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi 630004, Tamil Nadu, India

² Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

³ Texas A&M University, College Station, TX, USA

⁴ Department of Clinical and Translational Science, Marshall University Joan C. Edwards School of Medicine, 1700 Byrd Biotech Science Center, Huntington, WV 25755, USA


⁵ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, 608 002, Annamalai Nagar, Tamil Nadu, India

⁶ Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

Introduction

In the middle of the nineteenth century, the manufacturing and usage of plastic were raised annually owing to its durability, adaptability, strength and insulating properties. For instance, in 2018, the global plastic manufacturing rate climbed to 359 million tonnes (Plastic Europe 2019). According to Bilal and Iqbal (2020), plastics were divided into micro (100–5 mm) and nano-sized (100 nm) particles by the action of physical (photo radiation, weathering conditions), chemical and natural abrasion. Eighty per cent of microplastics were ultimately derived from the terrestrial environment (Jambeck et al. 2015; Dusi and Coors 2016), while 20% was derived from ocean-based sources (Wright et al. 2013; Karbalaeei et al. 2018). However, its ineffective disposal methods persist in the environment, such as landfilling, incineration, recycling and pyrolysis (Panda et al. 2010).

In general, freshwater is a critical impact on the transfer of microplastic debris from land to the ocean, which



Contents lists available at ScienceDirect

Fish and Shellfish Immunology

journal homepage: www.elsevier.com/locate/fsi

Molecular interaction analysis of β -1, 3 glucan binding protein with *Bacillus licheniformis* and evaluation of its immunostimulant property in *Oreochromis mossambicus*

Mahalingam Anjugam^a, Arokiadhas Iswarya^a, Ashokkumar Sibiya^{a,b}, Chandrabose Selvaraj^c, Sanjeev Kumar Singh^c, Marimuthu Govindarajan^{d,e}, Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f, Jeyachandran Sivakamavalli^g, Baskaralingam Vaseeharan^{a,*}

^a Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma colony, Karaikudi, 630004, Tamil Nadu, India

^b Centre for Animal Science Research and Extension Service, Foundation for Innovative Research in Science and Technology, Kelavannanvilai, NGO Colony Road, Nagercoil, 629002, Tamil Nadu, India

^c Department of Bioinformatics, Alagappa University, Science Block, 4th floor, Burma colony, Karaikudi, 630004, Tamil Nadu, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^g Department of Fisheries, Chonnam National University, Yeosu, South Korea

ARTICLE INFO

Keywords:

Beta 1
3 glucan binding protein
Bacillus licheniformis
Synbiotics
Immune responses
Antioxidant

ABSTRACT

Analyzing the health benefits of any two immunostimulants (synbiotics) in combined form and information on their interactions gain more visibility in the usage of synbiotics in aquafarms. With this intention, the current work explores the immunostimulant effect and structural interaction of synbiotic (β -1, 3 glucan binding protein from marine crab, *Portunus pelagicus* (Pp β -GBP) and *Bacillus licheniformis*) on *Oreochromis mossambicus*. The experimental diet was prepared with Pp β -GBP and *B. licheniformis*, and nourished to the fingerlings of *O. mossambicus* for 30 days. After the experimental trial, a higher growth rate and immune reactions (lysozyme, protease, myeloperoxidase and alkaline phosphatase activity) were noticed in the fish nourished with synbiotic (*B. licheniformis* and Pp β -GBP) enriched diet. Moreover, the synbiotic enriched diet elevated the antioxidant responses like glutathione peroxidase (GSH-Px) and catalase (CAT) activity in the experimental diet-nourished fish. At the end of the feed trial, synbiotic diet nourished fish shows an increased survival rate during *Aeromonas hydrophila* infection, reflecting the disease resistance potential of experimental fish. Also, the interaction between Pp β -GBP and *Bacillus licheniformis* was analyzed through computational approaches. The results evidenced that, Pp β -GBP interacts with the *B. licheniformis* through sugar-based ligand, β -glucan through a hydrogen bond with a good docking score. Thus, the synbiotic diet would be an effective immunostimulant to strengthen the fish immune system for better productivity.

1. Introduction

Currently, disease appearance in aquaculture and the development of antibiotic-resistant genes in pathogenic microorganisms become a significant hindrance towards aquaculture production. Fish culture under a restricted environment (intensive farming) creates stress conditions that weaken the immune system and makes the animal more

prone to infectious disease-causing microbes. An innovative feeding method is one of the left practices employed in aqua farming by feed manufacturers and farmers to attain better growth in cultured fish [1]. Immunostimulants are harmless and beneficial bio-agents administered to farming animals (intensive or extensive) to prevent the entry of infectious microbes and improve the growth rate of the cultured organisms. Supplementation of immunostimulants such as probiotics,

* Corresponding author. Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma colony, Karaikudi, 630004, Tamil Nadu, India.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.fsi.2021.12.044>

Received 15 July 2021; Received in revised form 16 December 2021; Accepted 23 December 2021

Available online 28 December 2021

1050-4648/© 2021 Elsevier Ltd. All rights reserved.



TOXIC EFFECT OF LAMBDA CYHALOTHRIN AND PRETILACHLOR ON BIOMASS AND HISTOLOGY OF TESTIS OF *Lampito mauritii* (KINBERG)

G. INDHUMATHI ^{a*}, V. KAVITHA ^{a*#} AND R. ANANDHAN ^{a#}

^aDepartment of Zoology, Annamalai University, Annamalai Nagar - 608 002, Tamil Nadu, India.

AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.56557/UPJOZ/2022/v43i173157

Editor(s):

(1) Dr. Juan Carlos Troiano, University of Buenos Aires, Argentina.

Reviewers:

(1) Sangeetha Naveen, University of Malaya, Malaysia.

(2) Cenk Kılıç, Turkiye.

Received: 05 July 2022

Accepted: 09 September 2022

Published: 17 September 2022

Original Research Article

ABSTRACT

Lambda cyhalothrin and pretilachlor is widely used to control agricultural pest but also affect non target organisms like earthworms. The present laboratory study was conducted to determine the effect of Lambda cyhalothrin and Pretilachlor on biomass and histology of testis of *Lampito mauritii*. Lower and higher sub-lethal concentrations of Lambda cyhalothrin (T1 and T2) and Pretilachlor (T3 and T4) was mixed with soil substrate. Biomass was observed once in 10 days up to 90 days. 10 non clitellate *L.mauritii* was introduced into each treatment. Soil substrate without lambda cyhalothrin and pretilachlor served as control. The present study was revealed that lambda cyhalothrin and pretilachlor was highly affected the growth of *L. mauritii*. For histopathology study, four adult *L. mauritii* were introduced in to C, T1, T2, T3 and T4. After 5th, 15th and 30th day, testis was dissected out. The results reported that histology of testis was highly damaged on 5th and 15th day of experiment than 30th day.

Keywords: Earthworm; *Lampito mauritii*; lambda cyhalothrin; pretilachlor; biomass; testis.

1. INTRODUCTION

Our nation father GandhiG told that agriculture is a backbone of India. The growing human population is expected to be 9.1 billion by 2050 (UN [1]). It needs more food production. So, chemical fertilizers and

pesticides are used in the agricultural field to avoid increasing demand for food which leads to soil contamination. India is the largest manufacturer of pesticides. More than 128 pesticides are registered in India Laxmi [2]. They are not only destroy pest and boostup plant growth also affect farmer's friend as

*Research Scholar;

#Assistant Professor;

*Corresponding author: Email: kavisamikshaa@gmail.com;

A Study on Phytochemical Analysis and Anti Stress Activity of Ficus Carica Fruits

Dr. C. Jayanthi¹, M. Jaya², M. Nirmala³, L. Nandhini⁴

Department of Chemistry, Government College for Women (Autonomous), Kumbakonam,
Tamil Nadu, India

ABSTRACT

The research work was aimed to identify the presence of phytochemicals qualitatively and quantitatively and to investigate the anti stress activity of ficus carica fruits. To assess the anti-stress activity of Ficus carica extract against the oxidative stress induced by hydrogen peroxide in RBC by using the parameters MDA, CAT, GPx, this study rationalizes Ficus carica fruit extract play an active role in providing anti-stress activity.

INTRODUCTION

Anxiety and depression are the most frequent psychiatric conditions commonly found in today's day-to-day life. A number of the population suffers from these conditions sometimes during their life (Selvi *et al.*, 2012). Homeostasis regulates the physiological actions in the body and depends on the stress and antioxidant levels in the cells. Stress is involved in the major portion of alterations of physiological actions, leading to pathogenesis (Kilari *et al.*, 2015). Stress is a common phenomenon that is experienced by every individual. When stress becomes extreme, it is harmful for the body, and hence, needs to be treated. Stress is involved in the pathogenesis of a variety of diseases including hypertension, peptic ulcer, immune depression, reproductive dysfunction, and behavior disorder (Tiwari *et al.*, 2014). Overload of stress increases free radicals, produces damage to neuronal receptors and a variety of tissues.

Free radical scavenging agents may have a great potential in ameliorating these disease disorders. Stress basically is a reaction of mind and body against change in the homeostasis. The productive stress is called eustress while harmful stress is called distress. Stress triggers a wide range of body changes called general adaptation syndrome (GAS). The stimuli, which produce GAS, are called stressors and range from physical to psychological factors including cold, heat, infection, toxins, and major personal disappointment (Lee and Rhee, 2017).

MATERIALS AND METHODS

Collection of Plant Materials

The fruit powder of *Ficus carica* were purchased in March 2023 from Siddha Medicinal shop, Thanjavur, Thanjavur district, Tamil Nadu, India.

Preparation of Plant Extract:

100 gram of the powder of *Ficus carica* fruit were transferred in to different conical flask (250ml). The conical flask containing 50ml of different solution (ethanol and water). The conical flask *Ficus carica* fruit were shake it well for 30 minutes by free hand. After 24 hrs, the extracts were filtered using what man filter paper No.1 and filtrate used for further analysis.

Phytochemical Screening

Chemical tests were carried out on the extract using standard procedures to identify the constituents as described by Sofowara (1993), Trease and Evans (1989) and Harborne (1973 and 1984).

Test for Tannins

About 1ml of sample is boiled in 20 ml of water in a test tube and then filtered. A few drops of 0.1% ferric chloride is added and observed for brownish green or a blue-black colouration.

Test for Saponin

About 2 ml of sample is boiled in 20 ml of distilled water in a water bath and filtered. 10ml of the filtrate is mixed with 10 ml of distilled water and shaken vigorously for a stable persistent froth. The frothing is mixed with 3 drops of olive oil and shaken vigorously, then observed for the formation of emulsion.



ANTI-CORROSIVE STUDY OF ADATHODA VASICA EXTRACT IN 1M HCL ON THE CORROSION OF MILD STEEL

S. Jasmine Mary¹, A. Vijayalakshmi², A. Parvathi Priya³

Article History: Received: 12.02.2023

Revised: 27.03.2023

Accepted: 12.05.2023

Abstract

In the current era, natural products extracts are used for corrosion inhibition activity. This method is an ecovative and non-toxic technique. Many researchers do their research on inhibition of corrosion on this green method. The efficiency of corrosion inhibition carried out by weight loss method. In the present study, the leaves extract of *Adathoda vasica* was studied as corrosion inhibitors by weight loss method. The variation of corrosion rate, percentage inhibition efficiency and surface coverage with increasing concentration were measure in this work.

Keywords: Keywords: *Adathoda vasica*; corrosion inhibitor; weight-loss method; Anti-corrosive study

¹Chemistry dept, Government of for Women, Kumbakonam, Tamil Nadu, India

^{2,3}Chemistry dept, R.M.K. Engineering College, Thiruvallur Dt, Tamil Nadu, India

Email: 1jasminephd2011@gmail.com, 2avl.sh@rmkec.ac.in, 3app.sh@rmkec.ac.in

DOI: 10.31838/ecb/2023.12.s2.258

1. Introduction

Corrosion is an inevitable natural process of deterioration of metals/materials by chemical or electrochemical process with their environment. It is one of the sources of waste [1]. Corrosion is called a billion-dollar thief because every year several billion loss occur due to this corrosion. As per International Zinc Association at 2021, India loses 5-7% [2] of Gross Domestic Product (GDP),

Corrosion is a natural, spontaneous and it is thermodynamically stable process favoured by nature. So, nobody prevents or stop corrosion. But, the rate of corrosion can be controlled (i.e) just like postponing the death of material. There are many methods to control corrosion and an usage of inhibitors in small quantities is a common process. Suppose chromates were used as corrosion inhibitors, environmental scientists pointed out the health hazards caused by chromium (VI) ions. Hence corrosion research scientists go for eco-friendly, innovative, non-toxic natural products extracts as corrosion inhibitors.

2. Experimental procedure:

2.1 Materials:

For the weight loss measurements, a mild steel rod (C 0.2% by wt) with approximate size of 2.5×2.5×0.1cm were used. Before that a pre surface treatment of dipped in pickling solution for the removal of rust and then it is polished on emery paper. After that the rod was washed and degreased by using alcohol and acetone.

Step Aerobics Is The Best

Dr.R. VIDHYA SREE
(B.P.E.S., M.P.E.S., M.Phil., Ph.D., MSc Yoga)
Director of Physical Education
Govt.College for Women,Kumbakonam

Abstract

Aerobic dance has its foundation in dance-inspired movements. It is a cardiovascular workout set to music in a group exercise setting. Aerobics is a form of physical exercise that combines rhythmic aerobic exercise with stretching and strength training routines with the goal of improving all elements of fitness (flexibility, muscular strength, and cardio-vascular fitness). It is usually performed to music and may be practiced in a group setting led by an instructor (fitness professional), although it can be done solo and without musical accompaniment. With the goal of preventing illness and promoting physical fitness, practitioners perform various routines comprising a number of different dance-like exercises. Formal aerobics classes are divided into different levels of intensity and complexity. A well-balanced aerobics class will have five components: warm-up (5–10 minutes), cardio vascular conditioning (25–30 minutes), muscular strength and conditioning (10–15 minutes), cool-down (5–8 minutes) and stretching and flexibility (5–8 minutes). Aerobic exercise is any physical activity that makes us sweat, causes to breathe harder, and gets our heart beating faster than at rest. It strengthens our heart and lungs and trains our cardiovascular system to manage and deliver oxygen more quickly and efficiently throughout our body. Aerobic exercise uses our large muscle groups, is rhythmic in nature, and can be maintained continuously for at least 10 minutes. Aerobic dancing involves any kind of dance put to music and can include everything from Zumba to hip-hop dancing. A typical dance class usually begins with a 5 to 10-minute warm-up, followed by a 20- to 30-minute aerobic routine, and ends with a 5- to 10-minute cool down.

Introduction:

Aerobic means "requiring oxygen." Aerobic metabolism occurs during low-intensity, long-duration exercises. One example would be jogging. Anaerobic means "without oxygen." Anaerobic metabolism in muscle tissue occurs during intense physical activities like sprinting or weight lifting. During aerobic exercise, we breathe faster and deeper than when our heart rate is at rest. We're maximizing the amount of oxygen in the blood. Our heart rate goes up, increasing blood flow to the muscles and back to the lungs. During anaerobic exercise, our body requires immediate energy. Our body relies on stored energy sources, rather than oxygen, to fuel itself. That includes breaking down glucose. Aerobic exercise is exercise that requires oxygen to full the work due to the length of activity. Generally speaking, any activities greater than 60 seconds in duration require oxygen to continue to burn carbohydrates or fat as fuel source. Anaerobic exercise would be any activities that do not require oxygen to continue to use fuel sources because they are short in duration generally less than 60 seconds. Examples of aerobic exercise would be jogging, running a marathon, or hiking. Examples of anaerobic exercise would include weight lifting, throwing a baseball, or a 100-meter sprint.

Aerobic dance is a choreographed, repetitive movement routine set to music. A typical aerobics program begins with 5 to 10 minutes of warm-ups and stretching, peaks with 20 to 30 minutes of target heart range dance, can include 20 minutes of a muscle stretching floor program known as body sculpting, and ends with 5 to 10 minutes of cool down and more stretching. Aerobic exercise is any physical activity that makes you sweat, causes you to breathe harder, and gets your heart beating faster than at rest. It strengthens your heart and lungs and trains your cardiovascular system to manage and deliver oxygen more quickly and efficiently throughout your body. Aerobic exercise uses your large muscle groups, is rhythmic in nature, and can be maintained continuously for at least 10 minutes.



Data-driven multi-valley dark solitons of multi-component Manakov Model using Physics-Informed Neural Networks

Meiyazhagan Jaganathan ^a, Tamil Arasan Bakthavatchalam ^{a,*}, Murugesan Vadivel ^a, Selvakumar Murugan ^a, Gopinath Balu ^a, Malaikannan Sankarasubbu ^a, Radha Ramaswamy ^b, Vijayalakshmi Sethuraman ^c, Boris A. Malomed ^{d,e}

^a Saama AI Research Lab, Chennai 600032, India

^b Centre for Nonlinear Science (CeNSc), Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, India

^c Department of Physics, Presidency College (Autonomous), University Of Madras, Chennai 600005, India

^d Department of Physical Electronics, School of Electrical Engineering, Faculty of Engineering, and Center for Light-Matter Interaction, Tel Aviv University, P.O.B. 39040, Ramat Aviv, Tel Aviv, Israel

^e Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica, Chile

ARTICLE INFO

Keywords:

Physics informed neural networks
Deep Learning
Multi-valley dark solitons
Nonlinear Schrödinger equation
Manakov model
Bose–Einstein condensates

ABSTRACT

In this paper, we employ a Deep Learning technique, namely Physics-Informed Neural Network for solving multi-component Manakov models. In particular, we consider three and four-coupled nonlinear Schrödinger equations. We show that in the former case, one of the components and two others can carry, respectively, a double valley dark soliton and asymmetric bright solitons. In the four-component Manakov model, all four components are predicted by PINN with high accuracy in which the first component is shown to be a triple-valley dark soliton. We compare the results with the exact solutions and bring out the ability of Deep Learning in solving coupled systems of nonlinear partial differential equations with high accuracy. The performance of the PINN in approximating the solutions is estimated by taking the squared difference between the exact and predicted squared magnitude of the solutions. We have also computed the L^2 -norm errors for real, imaginary and absolute-value components of the solutions to emphasize the accuracy of PINN.

1. Introduction

Most of the dynamical systems in nature are inherently nonlinear being described by nonlinear differential equations, either continuous or discrete. Finding their exact or approximate solutions is crucial to the understanding of most of the natural phenomena around us. They can be broadly classified into integrable and nonintegrable systems. While integrable systems can be analytically solved using celebrated techniques like Inverse Scattering Transform [1], Hirota method [2], Darboux transformation method [3] etc., nonintegrable systems can only be investigated in the general case by means of numerical simulations [4,5]. With the rapid advancement in computing power combined with the vast availability of data, the use of Machine Learning (ML) and Deep Learning (DL) techniques to investigate relevant problems in physics [6] has skyrocketed in recent times. In fact, the vast availability of data in the domain of both integrable and nonintegrable nonlinear dynamical systems has helped us to experiment with ML and DL methods to penetrate deep into the dynamics of natural phenomena.

In particular, the chimera states are predicted using ML [7] while recurrent neural networks are used to predict chaotic attractors [8–10]. Further, ML and DL algorithms have been used in the study of extreme events [11–14] as well as in the identification of ground states of Bose–Einstein Condensates (BECs) [15,16]. ML and DL methods have also been used to solve diverse systems of ordinary and partial differential equations (ODEs and PDEs) [17]. Raissi et al. [18] introduced a DL method called Physics-Informed Neural Networks (PINN) for solving problems involving linear and nonlinear PDEs by incorporating the physical information about the system into the Artificial Neural Network (ANN). The PINN model and its variants have been used in various fields, including bio-engineering [19–21], fluid dynamics [22,23], stochastic differential equations [24], nano-optics and metamaterials [25,26], design of photonic devices [27] and also in disease spreading dynamics [28–30].

In particular, PINNs have been widely used for solving nonlinear PDEs namely, Nonlinear Schrödinger (NLS) equation. For example, the

* Corresponding author.

E-mail address: tamilarasanbakthavatchalam@gmail.com (T.A. Bakthavatchalam).

<https://doi.org/10.1016/j.chaos.2023.113509>

Received 2 January 2023; Received in revised form 21 March 2023; Accepted 25 April 2023

Available online 15 May 2023

0960-0779/© 2023 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

RESEARCH



Thermal quantum correlations and teleportation in a graphene sheet

S. Bhuvaneswari¹ · R. Muthuganesan² · R. Radha¹

Received: 24 January 2023 / Accepted: 1 April 2023

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2023

Abstract

The characterization of quantum resources in dynamical systems is one of the most important problems to be addressed in quantum information theory. In this article, we investigate the behaviors of quantum correlations and teleportation technique in a graphene sheet comprising of disordered electrons in a two-dimensional honeycomb lattice. We use three different measures of quantum correlations such as entanglement, measurement-induced nonlocality and uncertainty-induced nonlocality. We study the ground state properties of the graphene sheet from the perspective of quantum correlations. At thermal equilibrium, we show that the band parameter strengthens the quantum correlations whereas the scattering strength weakens the correlations. Finally, the impact of the system's parameters on the teleportation technique is also expounded.

1 Introduction

Nonlocality is a fundamental feature of quantum systems which is known to be at the heart of the development of modern quantum technologies. Since the entanglement resources are crucial for various quantum information processing tasks such as dense coding, teleportation, remote state preparation and key generation [1], their characterization and quantification have attracted considerable attention ever since the inception of EPR paradox [2, 3] and Bell nonlocality [4, 5]. Further, the entanglement also quantifies correlations of nonlocal distributions between spatially separated particles. In the context of quantum information science, the characterization of any quantum system from the perspective of nonlocality is a fundamental and formidable task. The quantification of the nonlocal aspects present in pure states is more straightforward and entanglement provides the complete spectrum of the nonlocality. On the other hand, there are certain open challenges in the understanding of nonlocality of mixed states and multiqubit systems. In the realm of Bell nonlocality, it is widely accepted that the

entanglement captures only a portion of nonlocal character of the system. To support this, certain tasks were demonstrated in the environment of zero entanglement i.e., other quantum correlations were also detected in some naturally unentangled bipartite systems [6, 7]. To bring out the complete spectrum of nonlocality, researchers have identified various measures from the perspective of the measurements. Nonclassical signatures of quantum systems can be quantified by correlation measures such as quantum discord [8, 9], measurement-induced nonlocality (MIN) [10], measurement-induced disturbance (MID) [11] and uncertainty-induced nonlocality (UIN) [12]. The correlation measures with different notions have unique advantages in the context of information processing task. In particular, MIN and UIN are considered to be useful resources for quantum communication [13, 14] and quantum metrology [15–17] respectively.

The rapid developments which has taken place in the domain of quantum information processing has virtually shifted the attention of the semiconductor industry towards the fabrication of nanomaterials which work in the realm of quantum regime. The graphene is one of the most important nanomaterials with a relatively high decoherence time [18]. In view of the implementation of QIP, the behaviors of quantum resources in the graphene system are worthy of investigation. Graphene is a two-dimensional novel nanomaterial with carbon atoms hexagonally positioned on a honeycomb lattice and is predicted to show interesting magnetic phenomena like intrinsic ferromagnetism [19, 20], nontrivial electronic properties [21] etc. The electronic excitations in graphene arise due to chiral and massless Dirac fermions

✉ R. Radha
vittal.cnls@gmail.com

¹ Centre for Nonlinear Science (CeNSc), PG and Research Department of Physics, Government College for Women (A), Kumbakonam, Tamil Nadu, India

² Department of Physics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Břehov 7, 115 19 Praha 1-Star Mřsto, Czech Republic



Fabrication of an effectual, stable and reusable Mg-doped CdAl₂O₄ nanoparticles for photodegradation of toxic pollutants under visible light illumination

S. Mahalaxmi^{a,b}, G. Rajesh^{a,b}, P. Senthil Kumar^{a,b,*}, S. Akilandeswari^c, M. Arul Joshua^{a,b}, V. Uma Shankar^{a,b}, M. Ramya^{a,b}, K. Thirumalai^d, Gayathri Rangasamy^{e,f}

^a Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India

^b Centre of Excellence in Water Research (CEWAR), Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603 110, Tamil Nadu, India

^c PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^d Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India


^e School of Engineering, Lebanese American University, Byblos, Lebanon

^f Department of Sustainable Engineering, Institute of Biotechnology, Saveetha School of Engineering, SIMATS, Chennai, 602105, India

HIGHLIGHTS

- Preparation of a novel CdAl₂O₄ and Mg-doped CdAl₂O₄ nanoparticles under a co-precipitation approach.
- Mg-doped CdAl₂O₄ products slightly increase and reduce the energy gap compared to pure.
- The PL emission spectra of the pure CdAl₂O₄ and Mg-doped CdAl₂O₄ nanoparticles exhibited some defect or oxygen vacancies.
- ESR and Reactive species tests suggested that h⁺ and ·OH⁻ played significant roles in the BB and BG photocatalytic process.
- The Mg (0.075 M) doped CdAl₂O₄ catalyst exhibit substantially high photocatalytic capability.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling Editor: Kuan Shiong Khoo

Keywords:

Mg-doped CdAl₂O₄,
Photoluminescence
ESR
Scavenger
Photodegradation

ABSTRACT

The water contamination caused by discharging extensive organic dyes stuff into water bodies is one of the utmost significant concerns disturbing the environment and human life. CdAl₂O₄ spinel materials have been excellent in the elimination of emerging pollutants by the photocatalysis route. These materials, when altered through methods namely doping with Mg ions, have benefits over CdAl₂O₄, especially reduced energy gap and light absorbed in the visible region. The XRD established the creation of space group R $\bar{3}$ with no other phase step being found. The photoluminescence outcomes indicated that Mg-doped CdAl₂O₄ nanoparticles had the preventing e⁻-h⁺ recombination possibility, which was favorable for the photocatalytic process. The Mg (0.075 M)-doped CdAl₂O₄ catalyst had higher photocatalytic performance with 94 and 96% removal of two azo (BB and BG)

* Corresponding author. Centre of Excellence in Water Research (CEWAR), Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, Chennai.

E-mail address: senthilkumar@ssn.edu.in (P. Senthil Kumar).

<https://doi.org/10.1016/j.chemosphere.2023.138178>

Received 8 October 2022; Received in revised form 20 January 2023; Accepted 17 February 2023

Available online 22 February 2023

0045-6535/© 2023 Elsevier Ltd. All rights reserved.



Strategies for ameliorating the photodegradation efficiency of Mn-doped CdAl₂O₄ nanoparticles for the toxic dyes under visible light illumination

G. Rajesh^{a,b}, P. Senthil Kumar^{a,b,**}, S. Akilandeswari^c, Gayathri Rangasamy^{d,f,*}, S. Lohita^{a,b}, V. Uma Shankar^{a,b}, M. Ramya^{a,b}, K. Nirmala^{a,b}, K. Thirumalai^e

^a Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India

^b Centre of Excellence in Water Research (CEWAR), Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603 110, Tamil Nadu, India

^c PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^d School of Engineering, Lebanese American University, Byblos, Lebanon


^e Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India

^f Department of Sustainable Engineering, Institute of Biotechnology, Saveetha School of Engineering, SIMATS, Chennai, 602105, India

HIGHLIGHTS

- Mn-doped CdAl₂O₄ nanoparticles were fabricated for the first time by a chemical co-precipitation route.
- UV-DRS, ESR and PL investigation affirm the existence of Vo or lattice defects in the present product.
- The recyclable Mn-doped CdAl₂O₄ nanoparticles exhibited an improved dye degradation properties.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling editor: Kuan Shiong Khoo

Keywords:

Mn-doped CdAl₂O₄
COD
Scavenger species
ESR
Photodegradation

ABSTRACT

Worldwide environmental issues have been escalating with the growth of the global economy and become a vital problem. To solve the problems, we require an eco-friendly and sustainable binary catalyst for the degradation of Azo dye pollutants. In this work, magnetically reusable, multifunctional novel Mn-doped CdAl₂O₄ nanoparticles were effectively fabricated by the co-precipitation approach. It was utilized for the degradation of two Azo dyes, exhibiting 96 and 98% Mn (0.050 M)-doped CdAl₂O₄ removal rates under visible light illumination, and presenting improved photocatalytic capability than that of pure and other dopants. More notably, the Mn (0.050 M)-doped CdAl₂O₄ catalyst was recycled using centrifuges without major loss and displays almost similar photodegradation behaviors for six successive runs. According to the ESR measurements, outcome and quenching tests affirmed that 'OH⁻ and h⁺ radicals were better reactive species responsible for Azo dyes removal. A possible photodegradation reaction mechanism underlying the elimination of Azo dyes by Mn (0.050 M)-doped CdAl₂O₄ catalyst is also proposed. Elaborated analyzes by variable reaction parameters such as the role of reactive species and catalyst dosage, pH, COD and irradiation time in the degradation route was also discussed. We assume that our outcomes will provide novel insights into using a highly effectual Mn (0.050 M)-doped CdAl₂O₄ catalyst, with possible applications in the treatment of both industrial and domestic wastewater.

* Corresponding author. School of Engineering, Lebanese American University, Byblos, Lebanon.

** Corresponding author. Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India.

E-mail addresses: senthilkumar@ssn.edu.in (P. Senthil Kumar), granga1983@gmail.com (G. Rangasamy).

<https://doi.org/10.1016/j.chemosphere.2023.138069>

Received 29 September 2022; Received in revised form 23 December 2022; Accepted 5 February 2023

Available online 8 February 2023

0045-6535/© 2023 Elsevier Ltd. All rights reserved.



A synergistic consequence of catalyst dosage, pH solution and reactive species of Fe-doped CdAl₂O₄ nanoparticles on the degradation of toxic environmental pollutants

G. Rajesh^{a,b}, P. Senthil Kumar^{a,b,**}, S. Akilandeswari^c, Gayathri Rangasamy^{d,f,*},
Aindrila Mandal^{a,b}, V. Uma Shankar^{a,b}, M. Ramya^{a,b}, K. Nirmala^{a,b}, K. Thirumalai^e

^a Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India

^b Centre of Excellence in Water Research (CEWAR), Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603 110, Tamil Nadu, India

^c PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^d School of Engineering, Lebanese American University, Byblos, Lebanon


^e Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India

^f Department of Sustainable Engineering, Institute of Biotechnology, Saveetha School of Engineering, SIMATS, Chennai, 602105, India

HIGHLIGHTS

- A novel pure and Fe-doped CdAl₂O₄ nanoparticles fabricated by chemical coprecipitation approach.
- Fe (0.075 M) doped CdAl₂O₄ catalyst removed 94 and 96% for BB and BG dyes under visible illumination.
- ·OH⁻ and h⁺ radicals were the major scavenging species in the degradation of the two Azo dyes.
- Fe (0.075 M) doped CdAl₂O₄ catalyst showed 78 and 84% degradation efficiencies in six consecutive runs.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling Editor: Kuan Shiong Khoo

Keywords:

Fe-doped CdAl₂O₄
Photoluminescence
ESR
Scavenging species
Photodegradation

ABSTRACT

Industrial wastewater treatment techniques are one of the biggest challenges of the scientific community that necessitate an increased consciousness to address water scarcity worldwide. Herein, an eco-friendly and cost-effective process was demonstrated to cope with tannery, textile and pharmaceutical dye wastes through the co-precipitation of highly reusable Fe-doped CdAl₂O₄ samples. The XRD studies exposed the space group R $\bar{3}$ with no secondary phase step being found for all samples. The outcomes of optical absorbance spectra demonstrate that Fe doping diminished the energy gap from 3.66 to 1.67 eV. HR-TEM images of existing spherical particles and some of the particles' rod-like structures with little agglomeration were found for Fe (0.075 M) doped CdAl₂O₄ nanoparticles. The PL emission outcomes show that Fe doping effectively prevented the charge carrier's recombination in CdAl₂O₄ during photocatalysis. All Fe-doped CdAl₂O₄ samples demonstrated higher photodegradation behaviors towards the effectual degradation of both dye solutions as compared to pure CdAl₂O₄ samples. Particularly, Fe (0.075 M)-doped CdAl₂O₄ samples exhibited improved

* Corresponding author. Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India.

** Corresponding author. School of Engineering, Lebanese American University, Byblos, Lebanon.

E-mail addresses: senthilkumar@ssn.edu.in (P.S. Kumar), granga1983@gmail.com (G. Rangasamy).

<https://doi.org/10.1016/j.chemosphere.2023.137919>

Received 30 September 2022; Received in revised form 2 January 2023; Accepted 18 January 2023

Available online 23 January 2023

0045-6535/© 2023 Elsevier Ltd. All rights reserved.



Preparation and characterization of a novel cobalt-substitution cadmium aluminate spinel for the photodegradation of azo dye pollutants

G. Rajesh^{a,b}, P. Senthil Kumar^{a,b,**}, S. Akilandeswari^c, Gayathri Rangasamy^{e,f,*}, S. Lohita^{a,b}, V. Uma Shankar^{a,b}, M. Ramya^{a,b}, K. Thirumalai^d

^a Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India

^b Centre of Excellence in Water Research (CEWAR), Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603 110, Tamil Nadu, India

^c PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^d Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India


^e School of Engineering, Lebanese American University, Byblos, Lebanon

^f Department of Sustainable Engineering, Institute of Biotechnology, Saveetha School of Engineering, SIMATS, Chennai, 602105, India

HIGHLIGHTS

- A novel Co (0.025–0.075 M) doped CdAl₂O₄ was fabricated by chemical co-precipitation approach.
- The ESR and PL confirms the existence of a substantial number of V_O or defects state in the spinel product.
- The active species h⁺ and ·OH⁻ acted the predominant role in the photodegradation process.
- The novel photocatalyst shows outstanding performance in degrading pollutants in water.

GRAPHICAL ABSTRACT



ARTICLE INFO

Handling Editor: Kuan Shiong Khoo

Keywords:

Co-doped CdAl₂O₄
Scavenger species
ESR
Photodegradation
COD

ABSTRACT

Modern-year organic contaminants have been highly observed in ecosystems since they are not removed entirely and remain dangerous. Semiconductor binary oxide photocatalysts have been well accredited as capable technology for ecological contaminants degradation in the existence of visible irradiation. In this research, novel Co ions doped CdAl₂O₄ materials were fabricated by a facile co-precipitation approach. The fabricated pure and Co-doped CdAl₂O₄ exhibited the typical peaks of CdAl₂O₄ with the E_g of 3.66, 3.24, 2.57, and 2.41 eV respectively. The HR-TEM microstructures revealed that the Co (0.075 M) doped CdAl₂O₄ has rod-like morphology, and some places are spherical with particle sizes reaching 21 nm. The PL peaks of the Co (0.075 M)-CdAl₂O₄ are much lesser than that of the other dopant and pure CdAl₂O₄, representing much more effectual separation of generated e⁻ and h⁺ at the interface which in fact outcomes in superior expected photodegradation behaviours. The Co (0.075 M)-CdAl₂O₄ catalyst demonstrated the highest performances of 92 and 94% toward the degradation of both dyes, respectively, owing to the lowest e⁻ and h⁺ recombination rate. The Co (0.075 M) doped CdAl₂O₄ photocatalyst revealed outstanding reusability and stability under visible irradiation, retaining the performance of about 83 and 86% after the fifth consecutive run of BB and BG elimination. A probable photodegradation

* Corresponding author.

** Corresponding author. Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India.


E-mail addresses: senthilkumar@ssn.edu.in (P. Senthil Kumar), granga1983@gmail.com (G. Rangasamy).

<https://doi.org/10.1016/j.chemosphere.2023.138232>

Received 15 October 2022; Received in revised form 5 February 2023; Accepted 22 February 2023

Available online 23 February 2023

0045-6535/© 2023 Elsevier Ltd. All rights reserved.



Fabrication of a novel Ni-doped CdAl₂O₄ nanoparticles and applications in photo-oxidation processes under visible light illumination

G. Rajesh^{a,b}, P. Senthil Kumar^{a,b,*}, Gayathri Rangasamy^{c,d}, S. Akilandeswari^e,
Aindrila Mandal^{a,b}, V. Uma Shankar^{a,b}, M. Ramya^{a,b}, K. Nirmala^{a,b}, K. Thirumalai^f

^a Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603110, Tamil Nadu, India

^b Centre of Excellence in Water Research (CEWAR), Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam, 603 110, Tamil Nadu, India

^c School of Engineering, Lebanese American University, Byblos, Lebanon

^d Department of Sustainable Engineering, Institute of Biotechnology, Saveetha School of Engineering, SIMATS, Chennai, 602105, India

^e PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^f Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India

ARTICLE INFO

Keywords:

Ni-doped CdAl₂O₄
Scavenger species
ESR
Photoluminescence
Photodegradation

ABSTRACT

The existence of azo dyes (BB and BG) in textile wastewater is a major problem owing to their possible effect on human health and the environment. In this work, we examine the photodegradation performance under the visible sources of BB and BG are Azo dyes extensively applied in the food and textile industries. An initial screening of dissimilar Ni-doped CdAl₂O₄ nanoparticles was determined to classify the most outstanding candidate for BB and BG removal. The Ni (0.075 M) doped CdAl₂O₄ nanoparticles after 90 min illuminations, 95 and 96% of BB and BG removal were noticed in pH =5 (acidic) and 5 mg catalyst dosage conditions. The recyclability of the Ni (0.075 M) doped CdAl₂O₄ catalyst was tested under optimized circumstances. The outcomes demonstrate that Ni (0.075 M) doped CdAl₂O₄ products exhibit substantially high stability with above 85 and 90% removal after the sixth catalytic run. The scavenging (free radicals) experiments revealed that 'OH⁻ and h⁺ radicals play a central role in BB and BG degradation. Overall, the outcomes obtained propose that the suggested route may denote an appropriate technique for the elimination of BB and BG from textile wastes. This work shows the new pathway to improve more efficient materials for the elimination of industrial effluents.

1. Introduction

Water contamination has become a significant issue in the current era of climate change, industrialization, and urbanization. The primary ecological contaminants commonly encountered are pesticides, metal ions, aromatic compounds, pharmaceuticals, and organic/inorganic dyes [1]. Among them, inorganic/organic dye solutions are extensively applied in various industries like plastics, paint, tanneries, textile, and paper, which causes an everyday release of vast quantities of coloured wastes [2]. Organic/inorganic dyestuffs have lower biodegradability and are resistant to chemical reagents and heat. Furthermore, a maximum of them are regarded as carcinogenic components and mutagenic.

Their existence in water bodies can also disturb the aquatic ecosystem [3,4], so the elimination of these contaminants from water

bodies with sustainable and effective green approaches has become an essential task for researchers [5,6]. Dissimilar methods have been applied for dye elimination from water effluents, such as membrane filtration [7], reverse osmosis [8], adsorption [9], coagulation [10], ozone treatment [11], and photocatalytic degradation [12]. The Photocatalysis route is significantly probable in the remediation of the removal of inorganic/organic dyestuffs as it can completely mineralize inorganic/organic dyestuffs molecules into CO₂ and H₂O [13,14].

For this intent, dissimilar oxides materials like MnAl₂O₄ [15], CoAl₂O₄ [16], ZnAl₂O₄ [17], CaAl₂O₄ [18], MgAl₂O₄ [19], SrAl₂O₄ [20] and so are frequently applied. Generally, these mixed metal oxides materials have fascinated much attention owing to their numerous properties like, photocatalytic [16], magnetic [21], gas sensing [15], antibacterial activity [22], electrochemical performance [23], or applied in Li-ion batteries [24]. Tangcharoen et al., [25] reported the

* Corresponding author: P. Senthil Kumar, Department of Chemical Engineering, Sri Sivasubramaniya Nadar College of Engineering, Kalavakkam - 603110, Tamil Nadu, India.

E-mail address: senthilkumar@ssn.edu.in (P. Senthil Kumar).

<https://doi.org/10.1016/j.mcat.2022.112835>

Received 13 September 2022; Received in revised form 19 November 2022; Accepted 23 November 2022

Available online 30 November 2022

2468-8231/© 2022 Elsevier B.V. All rights reserved.



Unburned Carbon from Bagasse Fly Ash to Produce Activated Carbon in a Single-Stage Chemical Process

V. Hariharan¹, S. Renuka², P. Sutharsan³, K. Dhanaraj^{*4}

¹Department of Physics, Rajalakshmi Institute of Technology, Chennai, India.

²Department of Physics, Govt. College for Women (A), Kumbakonam - 612 001, Tamilnadu, India.

³Department of Physics, Sir Theagaraya College, Chennai, India.

⁴Department of Physics, Arunai Engineering College, Tiruvannamalai, India.

ABSTRACT: Bagasse ash waste is generated in sugar mills from the use of bagasse as a fuel. This waste frequently noticed, its way to landfilled places where it is disposed. The dumping of biomass wastes caused in numerous agricultural processes is usually an environmental problem as odour and pollutes the soil. Recycling these leftovers for the manufacturing of activated carbon, an adsorbent with numerous uses, including the removal of pollutants, was one option for such a problem. In this work, activated carbon was made from Bagasse ash using a chemical activation process at 800 °C. Fourier Transform-Infrared (FT-IR) spectroscopy, Scanning Electron Microscopy (SEM) and X-ray diffraction (XRD) were used to characterize the sample. The functional groups are confirmed by FT-IR and the existence of hydrocarbons. The SEM and XRD studies show that the generated activated carbon has a low amount of inorganic elements when compared to the precursor. These experimental results suggested that Bagasse ash might be used as a starter in the activated carbon manufacturing method, making it a cost effective resource.

KEYWORDS: Activated carbon, Bagasse fly ash, FT-IR, SEM, XRD.

1. INTRODUCTION

Bagasse is the non-homogeneous fibre residue that remains while crushing sugarcane stalks to get juice. Sugarcane bagasse has become one of the most prolific agricultural wastes, with a yearly output of more than 540 million metric tons [1]. It is estimated that sugar factory produces 3 tons of wet bagasse for 10 tons of sugarcane crushed, which indicates each sugar manufacturing country produces significant amount of bagasse. Even though, it is a byproduct, it has many applications like fuel and paper production. The bagasse is made up of pith fiber, an attempt has been made to produce and understand characteristic of activated carbon from bagasse. As number of raw material may be used to produce activated carbon, bagasse is chosen for the current study because over the recent years production of activated carbon in low cost has prompted a growing interest since it play a vital role in both waste management and pollution control [2-5]. Activated carbon has a long history of usage in the removal of chemical species from aqueous medium generated by the medicinal, agriculture, food and industrial applications [6]. The absorption properties of activated carbon mainly depend on pore volume, porosity and the functional groups presence [2]. The current study, the surface characteristics and the internal pore structure was studied using scanning electron microscope. The presence of functional group and crystalline formation during activation process is analyzed using FT-IR and XRD respectively.

2. MATERIALS AND METHODS

Sugarcane bagasse ash (SBA) used in this work collected after electric power generation in EDI parry sugarcane industry, Nellikuppam, Cuddalore disiret, Tamil Nadu, India. All of the chemicals utilized in this investigation were of analytical reagent (AR) quality. Bagasse ash was dried in a 110 °C oven for 6 hours. Subsequent, it was crushed with a micro hammer blade grinder and mesh sieve to a grain size of 10 or 32 meshes (2.0 mm or 500µm). For the characterization and synthesis of activated carbons, bagasse ash with only a particle size of 500 µm was utilized.

2.1. Single-stage chemical activation and carbonization process

In a glass beaker, 10 g raw material was combined with the chemical reagent at 1:4 ratios. Distilled water was included in an amount equal to ten times the total weight of the combination. The liquid was then mixed and heated to homogenise before being infused at 85 °C until a thick homogeneous powder was formed.



Unburned Carbon from Bagasse Fly Ash to Produce Activated Carbon in a Single-Stage Chemical Process

V. Hariharan¹, S. Renuka², P. Sutharsan³, K. Dhanaraj^{*4}

¹Department of Physics, Rajalakshmi Institute of Technology, Chennai, India.

²Department of Physics, Govt. College for Women (A), Kumbakonam - 612 001, Tamilnadu, India.

³Department of Physics, Sir Theagaraya College, Chennai, India.

⁴Department of Physics, Arunai Engineering College, Tiruvannamalai, India.

ABSTRACT: Bagasse ash waste is generated in sugar mills from the use of bagasse as a fuel. This waste frequently noticed, its way to landfilled places where it is disposed. The dumping of biomass wastes caused in numerous agricultural processes is usually an environmental problem as odour and pollutes the soil. Recycling these leftovers for the manufacturing of activated carbon, an adsorbent with numerous uses, including the removal of pollutants, was one option for such a problem. In this work, activated carbon was made from Bagasse ash using a chemical activation process at 800 °C. Fourier Transform-Infrared (FT-IR) spectroscopy, Scanning Electron Microscopy (SEM) and X-ray diffraction (XRD) were used to characterize the sample. The functional groups are confirmed by FT-IR and the existence of hydrocarbons. The SEM and XRD studies show that the generated activated carbon has a low amount of inorganic elements when compared to the precursor. These experimental results suggested that Bagasse ash might be used as a starter in the activated carbon manufacturing method, making it a cost effective resource.

KEYWORDS: Activated carbon, Bagasse fly ash, FT-IR, SEM, XRD.

1. INTRODUCTION

Bagasse is the non-homogeneous fibre residue that remains while crushing sugarcane stalks to get juice. Sugarcane bagasse has become one of the most prolific agricultural wastes, with a yearly output of more than 540 million metric tons [1]. It is estimated that sugar factory produces 3 tons of wet bagasse for 10 tons of sugarcane crushed, which indicates each sugar manufacturing country produces significant amount of bagasse. Even though, it is a byproduct, it has many applications like fuel and paper production. The bagasse is made up of pith fiber, an attempt has been made to produce and understand characteristic of activated carbon from bagasse. As number of raw material may be used to produce activated carbon, bagasse is chosen for the current study because over the recent years production of activated carbon in low cost has prompted a growing interest since it play a vital role in both waste management and pollution control [2-5]. Activated carbon has a long history of usage in the removal of chemical species from aqueous medium generated by the medicinal, agriculture, food and industrial applications [6]. The absorption properties of activated carbon mainly depend on pore volume, porosity and the functional groups presence [2]. The current study, the surface characteristics and the internal pore structure was studied using scanning electron microscope. The presence of functional group and crystalline formation during activation process is analyzed using FT-IR and XRD respectively.

2. MATERIALS AND METHODS

Sugarcane bagasse ash (SBA) used in this work collected after electric power generation in EDI parry sugarcane industry, Nellikuppam, Cuddalore disiret, Tamil Nadu, India. All of the chemicals utilized in this investigation were of analytical reagent (AR) quality. Bagasse ash was dried in a 110 °C oven for 6 hours. Subsequent, it was crushed with a micro hammer blade grinder and mesh sieve to a grain size of 10 or 32 meshes (2.0 mm or 500µm). For the characterization and synthesis of activated carbons, bagasse ash with only a particle size of 500 µm was utilized.

2.1. Single-stage chemical activation and carbonization process

In a glass beaker, 10 g raw material was combined with the chemical reagent at 1:4 ratios. Distilled water was included in an amount equal to ten times the total weight of the combination. The liquid was then mixed and heated to homogenise before being infused at 85 °C until a thick homogeneous powder was formed.

HOSTED BY



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

Synthesis and antimicrobial activity of silver nanoparticles: Incorporated *couroupita guianensis* flower petal extract for biomedical applications



S. Logambal^a, T. Thilagavathi^b, M. Chandrasekar^c, C. Inmozhi^d, Philippe Belle Ebanda Kedi^e,
F.A. Bassyouni^f, R. Uthrakumar^{a,*}, Azhaguchamy Muthukumar^g, Suresh Naveenkumar^g,
K. Kaviyarasu^{h,1,*}

^a Department of Physics, Government Arts College (Autonomous), Salem 636007, Tamil Nadu, India

^b Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

^c Department of Physics, TAG. Govt. Arts College, Tindivanam 604307, Tamil Nadu, India

^d Department of Physics, Government Arts College for Women, Salem 636008, Tamil Nadu, India

^e Department of Animal Biology and Physiology, Faculty of Science, University of Douala, P.O. Box 24157, Douala, Cameroon

^f Chemistry of Natural and Microbial Products Department, National Research Center, 12622 Cairo, Egypt

^g Department of Biotechnology, Kalasalingam Academy of Research and Education, Krishnankoil, Tamil Nadu, India

^h UNESCO-UNISA Africa Chair in Nanosciences/Nanotechnology Laboratories, College of Graduate Studies, University of South Africa (UNISA), Muckleneuk Ridge, PO Box 392, Pretoria, South Africa

¹ Nanosciences African Network (NANOAFNET), Materials Research Group (MRG), iThemba LABS-National Research Foundation (NRF), 1 Old Faure Road, 7129, PO Box 722, Somerset West, Western Cape Province, South Africa

ARTICLE INFO

Article history:

Received 11 June 2022

Revised 21 October 2022

Accepted 14 November 2022

Available online 17 November 2022

Keywords:

Couroupita guianensis flower petals

Silver nanoparticles

X-ray

DSL analysis

Antimicrobial activity

Photocatalytic activity

ABSTRACT

Recently, numerous plant-based synthesis techniques have been used to develop metal nanoparticles. The current study uses the medicinal plant extract of *couroupita guianensis* (CG) petals to create biogenic silver nanoparticles in an environmentally friendly manner. Different techniques, including ultraviolet-visible spectroscopy (UV-vis), Fourier Transformation Infrared spectroscopy (FTIR), X-ray diffraction (XRD), and Dynamic Light Scattering (DLS) analysis, are used to evaluate synthesized silver nanoparticles. From the XRD results confirm the emergence of nanosilver crystalline arrangement with the characteristic peaks at the glancing angles of 38.04°, 44.22°, 64.40°, and 77.37°. UV-vis spectroscopy displays the spectral absorption at $\lambda^{\text{max}} = 282$ nm and shows the formation of silver nanoparticles. Images of produced Ag NPs taken with a scanning electron microscope (SEM) show the creation of flower-shaped particles. The functional behavior of flavones, triterpenoids, and polyphenols belonging to *couroupita guianensis* has been observed by ensuring their selective absorptions in FTIR spectral analysis silver nanoparticles had a substantial antibacterial effect on Gram-positive (*B. subtilis*) and Gram-negative (*Escherichia coli*) bacteria in general. It is found to become effective when symbiotic with the extract of *couroupita guianensis* flower petals for enhancing their antibacterial properties. This composite product gives a new and cost-effective formulation with more therapeutic possibilities. The observed results wide open the avenues of research possibilities with a lot of future scopes. The photocatalytic degradation efficiency of CG-Ag NPs on methylene blue (MB) dye was evaluated under visible light irradiation and produced indications of the synthesized material for photocatalytic applications.

© 2022 The Authors. Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding authors at: UNESCO-UNISA Africa Chair in Nanosciences/Nanotechnology Laboratories, College of Graduate Studies, University of South Africa (UNISA), Muckleneuk Ridge, PO Box 392, Pretoria, South Africa (K. Kaviyarasu).

E-mail addresses: philippe_belle@yahoo.fr (P. Belle Ebanda Kedi), uthraloyola@yahoo.com (R. Uthrakumar), kavi@tlabs.ac.za (K. Kaviyarasu).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<https://doi.org/10.1016/j.jksus.2022.102455>

1018-3647/© 2022 The Authors. Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Medicinal plants are special elements known for their therapeutic scope and render long-time remedies for several human diseases. It is mainly because of the healing value of their components used as a raw material in modern medicine. The *couroupita guianensis* (CG) is a tree and its parts are familiar components used for medicinal purposes. Particularly, its chemical ingredients are responsible for various pharmacological and therapeutic properties. The study used to explore the characteristic



Sankaranarayanan-Ramasamy (SR) technique L- threonine lithium chloride single crystal: Unidirectional growth, optical and mechanical studies

K. Parasuraman^a, C. Inmozhi^b, T. Thilagavathi^c, M. Kumar^{d, e, *}, M.E. Raja Saravanan^d,
R. Uthrakumar^d

^a Department of Physics, Sathyabama Institute of Science and Technology, Chennai 600 119, India

^b Department of Physics, Government Arts College for Women, Salem 636008, Tamil Nadu, India

^c Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

^d Department of Physics, Government Arts College (Autonomous), Salem 636007, India

^e Department of Physics, Arignar Anna Government Arts College, Attur 636121, Tamil Nadu, India

ARTICLE INFO

Keywords:

Organic compound
Crystal growth
Nonlinear optical material
Microhardness Optical properties

ABSTRACT

L-Threonine Lithium Chloride (LTLC), an effective semi-organic NLO material, was successfully produced as a bulk single crystal using the Sankaranarayanan Ramasamy (SR) process. According to a study of a single crystal's X-ray diffraction, LTLC crystallises in an orthorhombic system with space group P212121. From the powder XRD method, unidirectional development along the plane (1 1 0) was established. The optical absorption spectra reveals that LTLC has a broad bandgap of 3.75 eV and is very transparent over the whole visible region. On the magnified crystal, photoconductivity measurements show positive photoconductivity. According to the photoluminescence investigation, LTLC emits green light with a wavelength of 535 nm. Vickers microhardness testing was done to determine the crystal's mechanical strength. At different temperatures and frequencies, the dielectric constant and dielectric loss were calculated. In addition to the density functional theory (DFT) calculations such as frontier molecular orbital's (FMOs), analysis was carried out. Utilizing Kurtz's Nd:YAG Laser, powder testing is another method used to assess the produced crystal's SHG activity.

© 20XX

1. Introduction

Higher values of the original frequency can be produced by nonlinear optical materials, which have uses in optoelectronics, fibre optic communication, and optical modulation [1,2]. Numerous researchers have looked for various NLO crystal types for laser applications recently [3]. Due to their high degree of chemical inertness, inorganic crystals were also commonly employed in optoelectronics applications due to their high melting point and outstanding mechanical stability.

Inorganic crystals' optical nonlinearity is typically less than the need for optical devices [4]. Vander Wall and hydrogen bonds that are weak and have a significant degree of delocalization frequently produce organic molecules. Significant drawbacks of organic NLO crystals include their higher optical quality, increased mechanical stability, and difficulty in making crystals at large sizes. [5]. Semi possesses NLO features because of its advantages in both organic and inorganic crystals. [6–7]. The potential for employing new semi-organic materials in a lot of interest has been shown recently in a variety of devices. Amino acids are currently the focus of significant research regarding the hunt for newer nonlinear optical materials since they may have a wide range of possible applications. Many amino acids exhibit outstanding optical

second harmonic generation and are optically active. SHG, which makes them the best option for uses such coherence of blue-green laser generation and frequency doubling [8–10]. Amino acid complexes with inorganic salts frequently combine the benefits of an organic amino acid and an inorganic salt, making them suitable building blocks for SHG properties. As a result, different inorganic acids were utilised to examine the production of salt utilising L-arginine, L-histidine, L-threonium, L-alanine, and L-valine. The result is superb semi-organic compounds have been investigated as prospective NLO application materials, including L-valine picrate [11], L-histidine hydrochloride [12], L-valine hydrochloride [13], L-analine zinc chloride [14], L-threonine cadmium chloride, and L-threonine lithium chloride. However, there is little information available regarding bulk LTLC growth for devices of the second harmonic generation. The SR method is unique for directional solution growth and also having many advantages such as defect-free, high transparency, good healthy bulk crystals in different orientations. Solute to crystal conversion efficiency 100% was achieved in SR method. The unidirectional crystal with a cylindrical shape is more important to avoid the post-growth machining process. The external morphology facets may induce strain through the lattices, it could be avoided in the single interface growth [15,16]. Therefore, higher qual-

* Corresponding author.

E-mail address: kumarsmkgac@gmail.com (M. Kumar).

<https://doi.org/10.1016/j.matpr.2023.05.288>

2214-7853/© 20XX

Antiuro lithiatic Activity of *Phyllanthus Niruri* Leaves Extract

Dr. C. Jayanthi¹, M. Nirmala², M. Jaya³, L. Nandhini⁴

Department of Chemistry, Government College for Women (Autonomous), Kumbakonam,
Tamil Nadu, India

ABSTRACT

Qualitative analysis of the phytochemicals has been carried out in *phyllanthus niruri leaves extract*. It was inferred that the phytochemicals tannin, saponin, flavonoids, steroids, terpenoids, anthroquinone, polyphenol, glycoside and anthocyanin were present. Antiuro lithiatic Activity was carried out by Nucleation assay Aggregation assay. The extract showed prominent inhibition of all Phases of CaOx stone formation. This study rationalizes *phyllanthus niruri leaves* has antiuro lithiatic activity.

INTRODUCTION

Kidney stones are small, hard deposits of mineral and acid salts developed from crystals that separate from the urine within the urinary tract. The most common type of stone contains calcium in combination with either oxalate or phosphate (Tyagi *et al.*, 2012). The problem of calculi formation is observed and reported in all parts of the urinary tract, the kidney, the ureter and the urinary bladder which may considerably vary in size (Havagiray *et al.*, 2010). The rate of occurrence is three times higher in men than women, because of enhancing capacity of testosterone and inhibiting capacity of oestrogen in stone formation (Kalpana *et al.*, 1993). Some of the Indian medicinal plants and their derivatives have been an invaluable source of therapeutic agents to treat many disorders including urolithiasis. An indigenous drug possessing fewer side effects is the major thrust area of the present day research, aiming for a better and safer approach for management of lithiasis (Koehn and Carter, 2005).

The prevalence and incidence of kidney stones is rising worldwide (Romero *et al.*, 2010). Kidney stones are hard deposits of minerals (calcium, oxalate and phosphate) which are formed from dissolved minerals in the urine and are usually excreted in the urethra. Kidney stones are the third most common urinary tract problem after urinary tract infections and prostate disorders. Kidney stones are classified into calcium oxalate, calcium phosphate, uric acid, cysteine, struvite, and mixed stones types, depending on the material of the stones. Calcium stones account for almost 70–80% of all kidney stones (Nikpay *et al.*, 2016).

MATERIALS AND METHODS

Collection of Plant Materials

The leaves powder of *Phyllanthus niruri* were purchased in March 2023 from Siddha Medicinal shop, Thanjavur, Thanjavur district, Tamil Nadu, India.

Preparation for Extract

Take one gram of *Phyllanthus niruri* leaves powder in each extract prepared in 50 ml of ethanol and aqueous solvent, the extract shake it well for 30 minutes by free hand and wait for 24 hours. After extracts were filtered using whatman filter paper No.1 and filtrate used for further analysis.

Phytochemical Screening

Chemical tests were carried out on the extract using standard procedures to identify the constituents as described by Sofowara (1993), Trease and Evans (1989) and Harborne (1973).

Test for Tannins: About 1ml of sample is boiled in 20 ml of water in a test tube and then filtered. A few drops of 0.1% ferric chloride is added and observed for brownish green or a blue-black colouration.

Test for Saponin: About 2 ml of sample is boiled in 20 ml of distilled water in a water bath and filtered. 10ml of the filtrate is mixed with 5 ml of distilled water and shaken vigorously for a stable persistent froth. The frothing is mixed with 3 drops of olive oil and shaken vigorously, then observed for the formation of emulsion.

eguCD8ADFK5 AK n C@CD
M 9 :A@FALE: 8 CD8ADFK
egu g a C@CD
a 12C@CD9t
c c a E9@l 9



e e

A5a s C5 g s D
Ad c e 5n s u t 5n d 5i 1 W 9 2
Cd i t 5n s u t 5n d 5i 1 W 9 2
Pd g 5g c w laZk 5t 8t 5t n 5i
1 W 9 2
u
5 9f 5
5 5 5 9t 5
8 8 9t o m 5i 9t 8
5 5 9s 5 8 5
5 A@8 5 9m 5
5 5 9t 5 5
5 9t 5 8 9f 5
9t 5 8 9



GOVERNMENT COLLEGE FOR WOMEN (AUTONOMOUS)

KUMBAKONAM – 612 001

Affiliated to Bharathidasan University

DST - CURIE Sponsored Institution

IV Cycle of Accreditation

☎ 0435 – 2401391

✉ principal@gwk.ac.in



CRITERION III – RESEARCH, INNOVATIONS AND EXTENSION

FRONT PAGE OF PUBLISHED PAPERS

2021-2022



INFLUENCE OF LEAF EXTRACT OF *CARICA PAPAYA* LEAF AGAINST MERCURIC CHLORIDE INDUCED HEPATIC ENZYMOLOGICAL CHANGES IN *LABEO ROHITA* (HAM.)

A.V. Kavitha¹, S.Vinodhini² and S.Kiruthika³

¹Assistant Professor, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

^{2,3}PG Students, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

ABSTRACT

In the fish *Labeo rohita*, the effect of a sub-lethal concentration of mercuric chloride (0.005 ppm) for 7 days on the activities of ACP, ALP, GOT and GPT, as well as their recovery (Ham.). When liver tissue is exposed to mercuric chloride, it shows a considerable decrease in ACP, ALP as well as an increase in GOT and GPT activities. During the recovery, all of the above metrics returned to near-normal levels thanks to *Carica papaya* treatment of mercury-treated fish. The findings suggested that exposing the fish *Labeo rohita* to *Carica papaya* competitively lowered mercuric chloride toxicity (Ham).

Key words: *Labeo rohita*, *Carica papaya*, ACP, ALP, GOT and GPT

INTRODUCTION

Many freshwater environments are threatened by pollution produced by the rise of industries, technology, and informal settlements. Pollution not only reduces the quality of water, but it also has an impact on all living species in the system. As a result, it's critical to not only identify and manage these pollution sources, but also to keep an eye on their consequences on the aquatic environment's health. Human activities are mostly responsible for increasing environmental pollution by introducing undesirable harmful substances (Bryan,1976).

Heavy metals are abundant in the natural environment, with the majority of them posing serious health risks to organisms (Bamennan and Schiesty, 1996). Pollution has toxicological effects due to its long-term persistence and build up in organisms (Goyer, 1996). Heavy metals play an important part in metabolic pathways, but when their concentrations above a certain threshold, they function as physiological, biochemical, and behavioural inhibitors in organisms. Metals are elements that naturally occur in aquatic habitats as a result of weathering and erosion (Viljoen, 1999). In trace levels, some of these elements are required by living organisms (for example copper and zinc). For growth and reproduction, essential trace elements have a narrow



Hematological studies of the freshwater Zooplankton *Macrocyclus edax*

P.Pavithra¹, K.Subhashini², R.Semmalar³, S.S.Padmapriya⁴, and S.Venkatalakshmi⁵

^{1,2}PG Students, Centre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

³Research Scholar, Centre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

⁴Guest Lecturer, Centre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

⁵Associate Professor and Head, Centre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India.

ABSTRACT

According to the literature review, this is one of the pioneer works in the observation of hemocytes in the cylopoid copepods. The homogenization and centrifugation methods were used to isolate the hemolymph of the zooplanktons. Using eosin stain, hemocytes were stained and identified under a trinocular light microscope. The presence and absence of cytoplasmic granules in the hemocyte cells were used to classify the hemocytes identified. Granulocytes, semi-granulocytes, and hyalinocytes were identified as three forms hemocytes. In a Neubauer counting apparatus, total and differential counts of hemocytes were performed. The results were recorded as 80,000 cells per μl of hemolymph and 45% hyalinocytes, 39% Granulocytes and 16% semi granulocytes.

Key words: Zooplankton, Copepods, hemocytes, Granulocytes, semi-granulocytes, hyalinocytes, *Macrocyclus edax*

INTRODUCTION

Aquatic animals are surrounded by microorganisms in the aquatic environment that are capable of initiating infection in these animals. Unlike fish that possess both innate and adaptive immunity, the lower aquatic animals especially invertebrates lack true adaptive immune systems and they depend upon their innate



Acute Toxicity and Histopathological Effect of Lambda Cyhalothrin and Pretilachlor on Intestine of Indian Earthworm *Lampito mauritii* (Kinberg)

Indhumathi.G¹, Kavitha.V^{2*} and Anandhan.R²

¹Research Scholar, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, India.

²Assistant Professor, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, India.

Received: 18 Feb 2022

Revised: 02 Mar 2022

Accepted: 25 Mar 2022

*Address for Correspondence

V. Kavitha

Assistant Professor,

Department of Zoology, Annamalai University,

Annamalai Nagar, Tamil Nadu, India

Email: kavisamikshaa@gmail.com



This is an Open Access Journal / article distributed under the terms of the **Creative Commons Attribution License (CC BY-NC-ND 3.0)** which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. All rights reserved.

ABSTRACT

Soil pollution has increased over time due to the widespread use of pesticides and chemical fertilizers in agriculture. Earthworms are important members of the soil ecosystem. In most ecological systems, earthworms play a crucial role in soil fertility and the formation of biogenic structures. Furthermore, earthworm gut mucus boosts the biological activities of beneficial soil microorganisms. The use of earthworms to enrich soil with essential nutrients is a cost-effective and environmentally friendly process. To control agricultural pests as well as weeds in India, the organophosphorus insecticide - lambda cyhalothrin and herbicide -pretilachlor are widely used. The present study the toxic effect of lambda cyhalothrin and pretilachlor of acute toxicity and histopathological effect reveals intestine of earthworm *L. mauritii* in a laboratory experiment. For acute study, mortality of *L. mauritii* was observed after 24, 48, 72, 96 and 120 h of the pesticides exposure. A sub-lethal concentration of lambda cyhalothrin and pretilachlor (1/5th of 96 h LC₅₀ value 1.8 ppm kg⁻¹ and 2.7 ppm kg⁻¹) was applied for 30 days. Histology of intestine was observed at 5th, 15th and 30th day of the pesticides exposure. On the 5th, 15th day of experiment, the skin showed significant pathological changes in epidermis layer, circular muscle, and longitudinal muscle. The changes such as vacuolization, degenerated nuclei, damaged epithelial lining of villi and congestion of blood sinuses was observed in 30th day, slight damages was observed. These results suggest that lambda cyhalothrin and pretilachlor could severely affect the intestine up to 15th day thereafter recovered of exposure when compared to 30th day. Histopathological study in *L. mauritii*'s intestine is a suitable parameter for detection of soil contamination by application of pesticides in agricultural field.





Facile synthesis and characterization of ZnO nanoparticles using *Abutilon indicum* leaf extract: An eco-friendly nano-drug on human microbial pathogens

Saraswathi Umavathi^a, Murugesan Subash^b, Kasi Gopinath^{c,*}, Saud Alarifi^d,
Marcello Nicoletti^e, Marimuthu Govindarajan^{b,g,**}

^a Department of Botany, Adhiyaman Arts and Science College for Women, Uthangarai, Tamil Nadu, 635207, India

^b Department of Botany, Arignar Anna Government Arts College, Attur, Tamil Nadu, 636121, India

^c Department of Packaging, Yonsei University, 1 Yonseidae-gil, Wonju-si, Gangwon-do, 26493, South Korea

^d Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^e Department of Environmental Biology, Sapienza University of Rome, Rome, 00185, Italy

^f Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^g Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Plant extract
Phytocompounds
Hexagonal wurtzite structure
ROS generation
Nano-drug carriers

ABSTRACT

The green synthesis of zinc oxide nanoparticles (ZnO NPs) using the leaf extract of *Abutilon indicum* and its characterization have been investigated. UV-vis, FT-IR, Micro-Raman, XRD, and SEM investigations were used to study the optical, structural, and morphological properties of produced ZnO NPs. The antimicrobial affectivity of ZnO NPs tested by well diffusion method counter to bacteria; *S. aureus*, *E. coli* and one fungal strain; *C. albicans*. The absorption peak of UV-vis spectrum was observed at 377 nm. FT-IR function vibration groups had proved the existence of ZnO NPs. The Micro-Raman analysis showed low and high-intensity bands of 330 and 436 cm^{-1} concerning ZnO NPs. The XRD configuration indicated the pure crystalline nature ZnO NPs formation with an approximate crystalline size at 27.89 nm. SEM images showed the structure of NPs to be a spherical shape with 10–30 nm. The ZnO NPs interact with bacterial cells by electrostatic interaction. It has enhanced the surface tension and generates reactive oxygen species (ROS), leading to bacterial lysis. Conveniently, synthesized ZnO NPs can be used against Gram^{-ve} bacteria causing infectious disease in the medical field.

1. Introduction

The rapid emergence of bacterial and fungal infections has become a significant cause of mortalities all over the globe. The availability of multiple resistance drugs for different bacteria and fungi has further made the condition more problematic. Worldwide annually, 50% of hospitalized patients are infected by multiple drug-resistant bacteria [1]. The development of new alternatives for combat drug-resistant bacteria became necessary. Various therapeutic values have been practiced since ancient times and contribute to essential tasks in traditional medicine. Plants will always be an alternative for the enhancement of novel drugs.

Abutilon indicum is a domineering herb with potent biological

activities that belong to the Malvaceae family. In the ancient Ayurvedic system, *A. indicum* was reported with inflammatory and antimicrobial activity and used to treat various diseases, including cough, gonorrhea, piles etc., [2,3]. Besides, essential biologically active constituents such as carbohydrates, alkaloids, starch, glycosides, flavonoids, and phenol are also accessible [4]. It has been commonly used as a stimulant purgative, moisturizing cream, pain-relieving, anti-diabetic, anti-inflammatory, and anti-dot medication as well as to cure leprosy, urinary infection, jaundice, masses, thirst relief, injury and ulcer cleansing, menstrual infections, rheumatism, mumps, respiratory illness, bronchitis, allergies, bloody dysentery, and ear disorders in conventional medicine [5,6]. The antioxidative, antimicrobial, anti-inflammatory, hepatoprotective, anticancer, immunomodulatory

* Corresponding author.

** Corresponding author. Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India.

E-mail addresses: gloriousnanogopi@gmail.com (K. Gopinath), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.jddst.2021.102917>

Received 30 June 2021; Received in revised form 9 October 2021; Accepted 9 October 2021

Available online 20 October 2021

1773-2247/© 2021 Elsevier B.V. All rights reserved.



Factors determine the population characteristics of migratory shorebirds and their prey species in the coastal saltpans

Jeganathan Pandiyan^{a,*}, Shahid Mahboob^b, Khalid A. Al-Ghanim^b, Fahad Al-Misned^b, Zubair Ahmed^b, Ganesan Karthikeyan^a, Kasi Gopinath^c, Marimuthu Govindarajan^{d,e,**}

^a Department of Zoology and Wildlife Biology, AVC College, Mannampandal, Mayiladuthurai, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Department of Packaging, Yonsei University, 1 Yonsei-dae-gil, Wonju-si, Gangwon-do, 26493, South Korea

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Coastal habitats
Man-made saltpans
Shorebirds
Prey
Conservation

ABSTRACT

Naturally-occurring seasonal saltpans serve as a major foraging ground for migratory shorebirds. Nevertheless, little is known about the role of various environmental factors in determining the density, diversity and species richness of shorebirds and their prey in India. This study was designed to collect quantitative data for the first time on the role of season, salinity, water depth, and prey availability in influencing the density, diversity, and species richness of migratory shorebirds in the eastern coastal saltpans of Tamilnadu, India. Selected sites were classified as seasonal saltpans and hypersaline saltpans based on the concentration of salinity in water. Shorebirds were counted systematically from (August) 2012 to (May) 2015 using the total count method. The measurement of salinity level and water depths of each sighting was also collected. The density, diversity and species richness of shorebirds showed highest in the seasonal saltpans than the hypersaline saltpans. The monsoon season attracted more shorebird density, diversity species richness when compared to other seasons. *Calidris alpina* was only the migratory shorebird species recorded in both types of saltpans and across seasons. The micro and macro-invertebrate prey species showed the highest in the seasonal saltpans. The Chironomid larva showed the highest turnover than other species recorded in the studied saltpans. The multiple regression models showed that the year, season, salinity, and water depth influence the shorebird density, diversity, species richness, and prey species' density with a more significant percentage. The study results found that saltpans are also an ideal habitat for migratory shorebirds and their diverse prey species seasonally.

1. Introduction

Shorebirds are fascinating creatures (Richards, 1998). They undertake phenomenal migration in a north-south direction each year propelled by natural selection to circumvent extremer cold weather. About two million shorebirds, most weighing <50 g undertake a monumental roundtrip from Alaska to Australia, covering 25,000 km. They surprisingly live long; the Eurasian oystercatchers, for example, are known to live beyond 40 years (Colwell, 2010). Therefore, a single shorebird has the potential to migrate over 200,000 km with a lifespan of just 10 years. The shorebirds embark on one of the longest-distance migrations

compared to all other species in the animal kingdom. A female shorebird, for instance, was recorded to have flown 11,500 km from Alaska to New Zealand and even without taking a break to feed and rest, which puzzled the scientific community (Hansford, 2007). Similarly, shorebirds caught and ringed by ornithologists at the Arctic Russian tundra, Kazakhstan and Uzbekistan were recaptured at the eastern coastal areas of Tamil Nadu, India, by our colleagues at the Bombay Natural History Society (Balachandran, 2012). Throughout their migratory routes, the shorebirds are naturally wedded to few stopover sites that face enormous threats due to various manmade disturbances.

India's coastal wetlands support thousands of migratory shorebirds

* Corresponding author.

** Corresponding author. Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India.

E-mail addresses: dunlinpandiyan@gmail.com (J. Pandiyan), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.ecss.2021.107490>

Received 5 July 2020; Received in revised form 28 June 2021; Accepted 1 July 2021

Available online 8 July 2021

0272-7714/© 2021 Elsevier Ltd. All rights reserved.



Original article

Phytochemical analysis and fabrication of silver nanoparticles using *Acacia catechu*: An efficacious and ecofriendly control tool against selected polyphagous insect pests



Mathalaimuthu Baranitharan^a, Saud Alarifi^b, Saad Alkahtani^b, Daoud Ali^b, Kuppusamy Elumalai^a, Jeganathan Pandiyan^c, Kaliyamoorthy Krishnappa^c, Mohan Rajeswary^{d,*}, Marimuthu Govindarajan^{d,e}

^aDepartment of Advanced Zoology & Biotechnology, Government Arts College for Men (Autonomous), Chennai 600035, Tamil Nadu, India

^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^cDepartment of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai 609305, Tamil Nadu, India

^dUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^eUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 20 July 2020

Revised 5 September 2020

Accepted 9 September 2020

Available online 17 September 2020

Keywords:

Nanotechnology

Green synthesis

Pest

Spodoptera litura

Helicoverpa armigera

Scanning electron microscope

ABSTRACT

Globally, the farmers are struggling with polyphagous insect pest, and it is the number one enemy of agricultural products, which made plenty of economic deterioration. *Spodoptera litura* and *Helicoverpa armigera* are the agronomically important polyphagous pests. Most of the farmers are predominately dependent on synthetic chemical insecticides (SCIs) for battle against polyphagous pests. As a result, the broad spectrum usage of SCIs led a lot of detrimental outcomes only inconsequently the researchers search the former-friendly phyto-pesticidal approach. In the present investigation, leaf ethanol extract (LEE) and silver nanoparticles (AgNPs) of *A. catechu* (*Ac*) were subjected to various spectral (TLC, CC, UV, FTIR, XRD and SEM) analyses. Larval and pupal toxicity of *A. catechu* *Ac*-LEE and *Ac*-AgNPs were tested against selected polyphagous insect pests. The significant larval and pupal toxicity were experimentally proven, and the highest toxicity noticed in AgNPs than *Ac*-LEE. The larval and pupal toxicity of *Ac*-AgNPs tested against *S. litura* and *H. armigera* LC₅₀/LC₉₀ values were 71.04/ 74.78, 85.33/ 88.91 µg/mL and 92.57/ 96.21 and 124.43/ 129.95 µg/mL respectively. *Ac*-AgNPs could be potential phyto-pesticidal effectiveness against selected polyphagous insect pests. In globally, it is significantly sufficient ratification giving towards the prevention of many unauthorized SCIPs.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Most of the quality and quantity of the agricultural product is directly/ indirectly deteriorated by insect pests in many countries (Elumalai et al., 2010; Kamaraj et al., 2008; Krishnappa et al., 2010; Krishnappa and Elumalai, 2012; Misra, 2014). *Spodoptera litura* is a predominant polyphagous pest occupied a wide range of hosting around 200 floral species globally, in which 74 floral host species noticed from India (Elumalai et al., 2014; Paulraj

et al., 2017). *S. litura* larvae consumed the different parts of host flora, including rhizome and causing severe damage, which gives above 60% of revenue loss in India (Elumalai et al., 2014; Krishnappa et al., 2010a, 2010b). India is a tropical country, polyphagous pest (*S. litura*) surviving and high abundance in that particular climate; therefore, recently, agriculture is facing severe economic losses (Paulraj et al., 2017). *Helicoverpa armigera* is a multivoltine, agronomically predominant polyphagous pest and it consumed a wide range of hosts estimated above 300 floras communities globally (Backiyaraj et al., 2014; Namin et al., 2014). The initial larval stage feeds only soft floral structures then turned to later stages feed on every part of flora (Gokulakrishnan et al., 2012).

Globally, most of the farmers are predominantly depending on synthetic chemical insecticides (SCIs) for battle against polyphagous insect fauna (Elumalai et al., 2010; Krishnappa and Elumalai, 2012). As the results of broad-spectrum usage of SCIs

* Corresponding author.

E-mail address: drrajvectorcontrol@gmail.com (M. Rajeswary).

Peer review under responsibility of King Saud University.





Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com

Original article

Cellulase immobilized magnetic nanoparticles for green energy production from *Allamanda schottii* L: Sustainability research in waste recycling

Shankar Vijayalakshmi^a, Marimuthu Govindarajan^{b,c}, Norah Al-Mulahim^d, Zubair Ahmed^d, Shahid Mahboob^{d,*}^aCO₂ Research and Green Technologies centre, VIT University, Vellore 632014, India^bDepartment of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India^cUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India^dDepartment of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

ARTICLE INFO

Article history:

Received 19 October 2020

Revised 5 November 2020

Accepted 8 November 2020

Available online 13 November 2020

Keywords:

Saccharomyces cerevisiae

Fermentation

Bioethanol

Immobilization

Biofuel

Allamanda schottii

ABSTRACT

This study presents ethanol's fabrication by fermenting the golden trumpet flower (*Allamanda schottii* L) with the yeast strain *Saccharomyces cerevisiae*. The changes in different parameters during fermentation were studied and optimized while producing the ethanol and the end product was subjected to emission test study by blending petrol and ethanol. The *Allamanda* floral substrate contains 65% polysaccharides. The strain *S. cerevisiae* was obtained in the form of baker's yeast from a domestic shop. For 100 ml of slurry, the highest bioethanol yield recorded was about 18.75 ml via optimization of different culture conditions, including a 1:8 ratio for slurry preparation, maintained under 35 °C, 5.5 pH, 72 h. old inoculum with a quantity of 3.75 g 100 ml⁻¹, fermented for 120 h. The highest yield of bioethanol was acquired under the addition of urea. This technique & design is capable of industrial-scale fabrication of bioethanol by using *A. schottii* floral substrates. This research was conducted to fabricate ethanol by fermentation (*A. schottii* L) floral substrate with *S. cerevisiae*. The optimum physiochemical parameters required to obtain the highest yield of bioethanol from *A. schottii* flower by fermentation was studied. The immobilization strategy with a cheap agricultural substrate and magnetic nanoparticles were also studied. The engine performance and emission studies were done with different blends of petrol and bio-ethanol.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Biomass energy reduces GHG emissions on a large scale. The CO₂ released on burning biomass is equal to fossil fuels (Kongkiattikajorn and Sornvoraweat, 2011). However, during photosynthesis, the CO₂ is captured for the growth of biomass and hence a balance exists. The biomass can be grown on underutilized farmland (Rankovic et al., 2009). Biofuels are the only renewable liquid transportation fuels that can reduce dependence on foreign

oil. Huge biomass potential is available in our country to produce biodiesel and bioethanol, so investing in this sector proves to be economical (Raita et al., 2016). When burnt, the biomass can pollute the air, but at low levels than fossil fuels' burning. The Sulphur content, which causes acid rain, are not produced while burning the biomass. The burnt biomass releases CO₂, contributing to GHG emission but compensated by the photosynthesis process during biomass growth (Liu et al., 2018).

The first-generation biofuels are obtained from sugary, starchy and fatty food crops. Molasses, the byproduct of the sugar industry, are used to produce ethanol (Talebna et al., 2010). The 2nd generation biofuels are mainly from lignocellulosic materials. The raw materials like wood, straw, agricultural, horticultural residues and forest waste are available in large quantities (Mittal, 1992). Various methods are available to convert these residues into biofuels. This type of biomass focuses mainly on avoiding food crop resources, which pose a threat to food security (Abdel Ghany et al., 2014). The second-generation processes mainly aim to

* Corresponding author at: Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia.

E-mail address: mushahid@ksu.edu.sa (S. Mahboob).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com

Original article

Optimization strategies for improved biogas production by recycling of waste through response surface methodology and artificial neural network: Sustainable energy perspective research



Lakshmi C. Gopal ^a, Marimuthu Govindarajan ^{b,c,*}, M.R. Kavipriya ^d, Shahid Mahboob ^{e,*}, Khalid A. Al-Ghanim ^e, P. Virik ^e, Zubair Ahmed ^e, Norah Al-Mulhm ^e, Venkatesh Senthilkumaran ^f, Vijayalakshmi Shankar ^{f,*}

^a SJB Institute of Technology, Bengaluru 560060, Karnataka, India

^b Department of Zoology, Annamalai University, Annamalaiagar 608002, Tamil Nadu, India

^c Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

^d Department of Botany, Alagappa University, Karaikudi 630004, Tamil Nadu, India

^e Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^f CO₂ Research and Green Technologies Center, VIT University, Vellore 632014, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 13 October 2020

Revised 3 November 2020

Accepted 16 November 2020

Available online 23 November 2020

Keywords:

Flower waste

Biogas production

Response surface methodology

Artificial neural network

Pretreatments

Sustainable energy

ABSTRACT

Objective: The primary aim of the study is to augment the biogas production from flower waste through optimization and pretreatment techniques.

Methods: Enhancement of biogas production by using response surface methodology (RSM) and artificial neural network (ANN) was done. The time for agitation, the concentration of the substrate, temperature and pH were considered as model variables to develop the predictive models. Pretreatment of withered flowers was studied by using physical, chemical, hydrothermal and biological methods.

Results: The linear model terms of concentration of substrate, temperature, pH, and time for agitation had effects of interaction ($p < 0.05$) significantly. From the ANN model, the optimal parameters for the biogas production process increased when equaled to the model of RSM. It indicates that the artificial neural network model is predicting the yield of biogas efficiently and accurately than the RSM model. Chemical pre-treatments were found to enhance the biogas production from flower waste with higher biomethane kinetics and cumulative yield.

Conclusion: Biogas production was significantly improved with statistical optimization and pretreatment techniques.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding authors at: Department of Zoology, Annamalai University, Annamalaiagar 608002, Tamil Nadu, India (M. Govindarajan); CO₂ Research and Green Technologies Center, VIT University, Vellore 632014, Tamil Nadu, India (S. Vijayalakshmi).

E-mail addresses: drgovind1979@gmail.com (M. Govindarajan), mushahid@ksu.edu.sa (S. Mahboob), vijimicro21@gmail.com (V. Shankar).

Peer review under responsibility of King Saud University.



1. Introduction

In recent times, about 85% of the world's universal energy demands will be obtained from fossil fuels (Edenhofer et al., 2010), these are contributing to a lot of global warming and natural damages (Nigam and Singh, 2011; Rahman et al., 2011). Present approaches deliver the utilization of alternative energy, which is renewable, such as biomass, geothermal, solar and wind, to decrease the GHG. There is an immense possibility for the generation of biofuels from biomass (Naik et al., 2010).

Liquid biofuels or gaseous fuels that are chiefly made by using biomass will be classified as three generations (1st, 2nd and 3rd generations) (Dragone et al., 2010). From the edible portion of plants, such as seeds, grains and sugars, are considered to be

Research Article

Shanmugam Rajeshkumar*, Munusamy Tharani, Vijayarangan Devi Rajeswari, Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, Kasi Gopinath, Natesan Vijayakumar, and **Marimuthu Govindarajan***

Synthesis of greener silver nanoparticle-based chitosan nanocomposites and their potential antimicrobial activity against oral pathogens

<https://doi.org/10.1515/gps-2021-0060>

received July 14, 2021; accepted September 21, 2021

Abstract: In the present investigation, silver nanoparticles (AgNPs) and silver nanoparticle-based chitosan nanocomposite were synthesized using *Cissus arnottiana* leaf extract. The biosynthesized nanoparticles and nanocomposites were characterized using SEM, TEM, and AFM to uncover the morphological characteristics such as size and shape. The SEM image depicts the size of the nanocomposite to be 30–40 nm and shape as spherical. The TEM results reveal the shape of the nanocomposite to be spherical and size around 10–60 nm. The XRD results show the crystalline nature of the AgNPs-based chitosan nanocomposite. The SAED analysis pattern seems to be concordant with the XRD results. The AFM image ensured the precise surface morphology of the synthesized silver nanocomposite

in the 3-dimensional pattern. The antimicrobial efficacy of the biosynthesized AgNPs, AgNP nanocomposite, and chitosan nanoparticles was tested against oral pathogens. The results revealed a potential antimicrobial effect, which states that it must be converted into nanomedicine to meet future biomedical needs.

Keywords: green synthesis, silver nanoparticles, chitosan nanocomposite, *Cissus arnottiana*

1 Introduction

Recently, the field of nanotechnology arised to a greater extent due to its ablazing applications in biomedical field [1]. This attracts huge number of researchers to create different nanomaterials with specific functions to cure a disease or to enhance and perform in equipments or products such as health care products, cosmetics, household products, etc. The nanolevel requirement of these nanomaterials in living cells plays a significant role in outraging the disease-causing pathogens and organisms [2].

Nowadays, the disease-causing pathogens become more resistant to upcoming antibiotic drugs. To combat this issue, metallic nanoparticles like silver nanoparticles (AgNPs) have been reported in several studies by researchers as an effective antimicrobial agent [3–5]. The AgNPs are majorly used in treating burn and open injuries to avoid contamination from wound pathogens and other nosocomial pathogens [6]. AgNPs assume a significant part in science and medication due to their desirable physicochemical properties. AgNPs are known to have antifungal, anti-inflammatory, antiviral, antibacterial, antiangiogenesis, and antiplatelet properties [7–9].

In several studies, chitosan, a natural biopolymer, has been reported to enhance the antibacterial efficacy of the metallic nanoparticles [10–12]. In this present study, chitosan was added to *Cissus arnottiana* leaf

* **Corresponding author: Shanmugam Rajeshkumar**, Department of Pharmacology, Saveetha Dental College and Hospital, SIMATS, Chennai 600077, Tamil Nadu, India, e-mail: ssrajeshkumar@hotmail.com

* **Corresponding author: Marimuthu Govindarajan**, Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalinagar 608002, Tamil Nadu, India; Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

Munusamy Tharani: Department of Pharmacology, Saveetha Dental College and Hospital, SIMATS, Chennai 600077, Tamil Nadu, India

Vijayarangan Devi Rajeswari: Department of Bio-Medical Sciences, School of BioSciences and Technology, Vellore Institute of Technology (VIT), Vellore 632014, Tamil Nadu, India

Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled: Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

Kasi Gopinath: Department of Packaging, Yonsei University, 1 Yonseidae-gil, Wonju-si, Gangwon-do 26493, South Korea

Natesan Vijayakumar: Department of Biochemistry and Biotechnology, Faculty of Science, Annamalai University, Annamalinagar 608002, Tamilnadu, India

Research Article

Ramachandran Ishwarya, Kavitha Saravanan, Daniel Selvaraj, **Marimuthu Govindarajan***, Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, Jeyachandran Sivakamavalli, Thangaswamy Selvakumar, and Baskaralingam Vaseeharan*

Antibacterial greener silver nanoparticles synthesized using *Marsilea quadrifolia* extract and their eco-friendly evaluation against Zika virus vector, *Aedes aegypti*

<https://doi.org/10.1515/gps-2021-0070>

received July 19, 2021; accepted October 24, 2021

Abstract: Fabrication and use of nanoparticles have progressively enlarged within the last decade. Herein the silver nanoparticles (AgNps) were synthesized via the extract from *Marsilea quadrifolia* (*Mq*) as a decreasing and steadying mediator. The *Mq*-AgNPs demonstrated superior toxicity on Zika virus vector, *Aedes aegypti* with the LC₅₀ value of 10.69 µg·mL⁻¹. The *Mq*-AgNPs were established securely to non-target organisms *Artemia nauplii*

and *Ceriodaphnia cornuta*, and no structural or anatomical alterations confirm its security to the surroundings. The antibacterial activity exposed that *Mq*-AgNPs showed superior growth inhibition effect against the tested bacteria. Furthermore, the *Mq*-AgNPs showed strong antioxidant activities when compared to *Mq* leaf extract. Overall, our results highlight that *Mq*-AgNPs are hopeful and biodegradable devices against mosquito vectors of therapeutic significance, with moderate toxicity against non-target aquatic animals.

Keywords: *Marsilea quadrifolia*, Zika virus, *Artemia nauplii*, *Ceriodaphnia cornuta*, antioxidant assay

* **Corresponding author: Marimuthu Govindarajan**, Department of Zoology, Unit of Vector Control, Phytochemistry and Nanotechnology, Annamalai University, Annamalaiagar-608 002, Tamil Nadu, India; Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

* **Corresponding author: Baskaralingam Vaseeharan**, Department of Animal Health and Management, Biomaterials and Biotechnology in Animal Health Lab, Alagappa University, Karaikudi-630004, Tamil Nadu, India, e-mail: vaseeharanb@gmail.com

Ramachandran Ishwarya, Kavitha Saravanan: Department of Animal Health and Management, Biomaterials and Biotechnology in Animal Health Lab, Alagappa University, Karaikudi-630004, Tamil Nadu, India

Daniel Selvaraj: Department of Zoology, Scoot Christian College, Nagercoil-629 003, Tamil Nadu, India

Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled: Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

Jeyachandran Sivakamavalli: Department of Fisheries, Chonnam National University, Yeosu, South Korea

Thangaswamy Selvakumar: Centre for Research in Animal Science and Extension Services, Foundation for Innovative Research in Science and Technology, Kelavannan vilai, NGO Colony Road, Nagercoil-629002, Tamil Nadu, India

1 Introduction

Mosquito vectors are exclusively liable for spreading pathogens and parasites of different infections like dengue, malaria, chikungunya, encephalitis, Zika virus infection, filariasis, and yellow fever [1–3]. World Health Organization (WHO) approximates that dengue is passed on from 50 to 400 million citizens yearly on the earth. The appearance of dengue-resistant strains is a significant challenge to hold; it has extended and impacted human health [4]. Since commercial vaccines for most diseases spread by *Aedes aegypti* do not exist, mosquito control remains a critical component of all prevention and control campaigns, which typically rely on pesticide spraying, biological control agents, and environmental management, among other things. As a result, developing novel, environmentally acceptable, and effective mosquito control techniques are critical to ensuring our future ability to prevent and manage diseases spread by these insects. Mosquito management is being improved in numerous regions; however, essential confronts and the rising mosquito resistances to insecticides require other



Original article

An assessment of level of heavy metals pollution in the water, sediment and aquatic organisms: A perspective of tackling environmental threats for food security

Jeganathan Pandiyan^a, Shahid Mahboob^{b,*}, Marimuthu Govindarajan^{c,d}, Khalid A. Al-Ghanim^b, Zubair Ahmed^b, Norah Al-Mulhm^b, Rajendran Jagadheesan^a, Kaliyamoorthy Krishnappa^a^aDepartment of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai – 609305, Tamil Nadu, India^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar-608 002, Tamil Nadu, India^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 4 November 2020

Revised 13 November 2020

Accepted 22 November 2020

Available online 2 December 2020

Keywords:

Pollution
Heavy metals
Habitats
Benthic
Conservation

ABSTRACT

Pollution is severely threatening the wetland habitats. Heavy metals are one among of the major pollutants in wetland habitats. The cadmium (Cd), copper (Cu), chromium (Cr), cobalt (Co), lead (Pb), mercury (Hg), nickel (Ni) and zinc (Zn), were assessed in the water, sediment, benthic species (polychaetes, mollusc and crustaceans), prawn and fishes. The assessment of heavy metals was done by using double-beam Atomic Absorption Spectrophotometer (AAS). The Hg, Cr and Co were greater in sediment, Ni and Cd were higher in polychaetes and molluscs respectively. However, the Cu and Pb greater in crabs and the Zinc was greater in fishes. The concentration of metals showed significant differences among the various sources examined ($P < 0.05$) except Cr ($P > 0.05$). The inter-correlational analysis among the metals assessed from the various sources showed that the Cr and Pb not correlated among the eight metals examined. However, the Cu and Co were correlated with Hg ($r = 0.307$) and ($r = 0.788$) respectively. The nickel was correlated with Hg ($r = 0.367$), Cu ($r = 0.362$) and Co ($r = 0.432$). The Zinc was correlated with the Cd ($r = 0.331$) and Hg ($r = 0.737$). However, correlation of metals among the different sources shown that the metals of polychaetes correlated with sediment $r = 0.637$, the metals of crabs correlated with the sediment and polychaetes $r = 0.630$ and $r = 0.842$ respectively, the metals of molluscs was also correlated with sediment ($r = 0.636$), polychaetes ($r = 0.889$) and crabs ($r = 0.894$). In addition to that the metals of prawns was correlated with the polychaetes ($r = 0.839$), crabs ($r = 0.628$) and molluscs ($r = 0.634$). The metals of fishes correlated with polychaetes ($r = 0.529$), crabs ($r = 0.710$), molluscs ($r = 0.493$) and prawns ($r = 0.593$). Indeed the multiple regression model explained that the metals of sediments influence the accumulation of metals in biotic species such as polychaetes, molluscs, crustaceans, prawns and fishes with 84% ($F = 21.079$; $p < 0.001$). The order of the heavy metals in the water, sediment and biotic species was $Hg > Pb > Ni > Cr > Zn > Co > Cu > Cd$. The study found that the level of heavy metals at various sources in the sanctuary is showing considerable warning and the sanctuary is required intensive assessment on various aspects of pollution since the Point Calimere Wildlife Sanctuary is supporting several species of migratory and endangered shorebirds seasonally.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The wetlands habitats are most productive ecosystem in the world, providing suitable habitats for various species of aquatic organisms which are depends on it. Nevertheless the coastal wetland habitats including intertidal mudflats are functioning as a proper shelter, feeding and breeding grounds for various species (Balachandran, 2012; Sivaperuman and Venkatraman, 2014). But recently several wetlands are under severe threats due to various

* Corresponding author.

E-mail address: mushahid@ksu.edu.sa (S. Mahboob).

Peer review under responsibility of King Saud University.





Facile synthesis of gold and platinum doped titanium oxide nanoparticles for antibacterial and photocatalytic activity: A photodynamic approach

Shahid Mahboob^{a,*}, Ravi Nivetha^b, Kasi Gopinath^{c,*}, Chinnasamy Balalakshmi^d, Khalid A. Al-Ghanim^a, Fahad Al-Misned^a, Zubair Ahmed^a, Marimuthu Govindarajan^{e,f}

^a Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^b Center for Nanotechnology Research, Vellore Institute of Technology, Vellore, 632014, Tamilnadu, India

^c School of Materials and Energy, Southwest University, Chongqing, 400715, China

^d Department of Nanoscience and Technology, Alagappa University, Karaikudi, 630 003, Tamilnadu, India

^e Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, 608 002, Tamil Nadu, India

^f Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Phytosynthesis
Enterolobium saman
Metal-doping
Photoanode
Photocatalytic activity
Antibacterial activity

ABSTRACT

A simple method has been needed to synthesize nanoparticles (NPs) to avoid environmental pollution, an alternative chemical and physical method. This current study deals with phytosynthesis of gold (Au) and platinum (Pt) metal doped with titanium oxide (TiO₂) NPs using *Enterolobium saman* bark extract. This extract plays a vital role in reducing and stabilizing Au and Pt doped into the TiO₂ NPs lattices. Phytosynthesized samples were characterized by XRD, SEM, ED-XRF, TEM, FTIR, Raman, and UV-vis-DRS analyses. The metal doping effect has decreased bandgap energy and particle size, whereas increased conductivity for TiO₂/M-Au and TiO₂/M-Pt NPs compared to pristine TiO₂ NPs. Phytosynthesized NPs were fabricated for dye-sensitized solar cell (DSSC) and photocatalytic behaviour against methylene blue (MB) dye was studied. An obtained result demonstrates that TiO₂/M-Au NPs have excellent feasibility for applying DSSC and photocatalytic application due to particle size, crystallite size, absorption ability, and bandgap energy. Besides, synthesized samples were measured with cyclic voltammetry and impedance spectroscopy found that the metal doping is drifted the dielectric and increases that the metal doping is drifted the dielectric increases electro-catalytic of the TiO₂. Different concentrations of all NPs were tested against *Escherichia coli* MTCC 40 and *S. aureus* ATCC 6633 bacteria by a well-diffusion method. The 10 mg concentration of all NPs showed better antibacterial activity. However, we believe that the proposed simple phytosynthesized method provides an efficient way to overcome the chemical and physical methods.

1. Introduction

Apparently, due to the lightweight, easy accessibility, high performance, and non-toxicity, biologically derived titanium dioxide (TiO₂) based nanomaterials have recently been very desirable in the field of energy harvesting and environmental remediation [1]. The Food and Drug Administration (FDA) and Environmental Protection Agency (EPA) regulations have reported that 50 µg/kg body weight/day of TiO₂ nanoparticles (NPs) is a safe dose for humans. In contrast, the toxicity of TiO₂ NPs in cellular function depending on the size, shape, surface defect, coating, adsorbed groups, solubility, and surface charge [2]. TiO₂-based nanomaterial has considerable potential for dye-sensitized solar cells (DSSCs) applications. DSSC promises third-generation solar

cells for research and development in the area of renewable energy harvesting. High power conversion efficiency (PCE) and stable solar cells because of their cost-effectiveness. Due to the optical band gap and effective charge isolation separation, the TiO₂ nanostructure has also been used in photocatalytic dye degradation applications [3]. A traditional DSSC consists of four components: dye as a primary sun-absorbing part to generate photoelectron, nanocrystalline semiconducting material to anchor dyes, and develop the electron-hole pair, an effective electro-catalytic counter-electrode to transmit the produced electron-holes pair to complete the circuit [4]. Compared to other metal oxides such as SnO₂ and ZnO [5], regular nanocrystalline TiO₂ layers are used as photoanodes. Dyes based on ruthenium are more effective and are used conventionally in different dyes tested for better absorption [6].

* Corresponding authors.

E-mail addresses: mushahid@ksu.edu.sa (S. Mahboob), gloriousnanogopi@gmail.com (K. Gopinath).

<https://doi.org/10.1016/j.pdpdt.2020.102148>

Received 26 October 2020; Received in revised form 30 November 2020; Accepted 11 December 2020

Available online 17 December 2020

1572-1000/© 2020 Elsevier B.V. All rights reserved.



Contents lists available at ScienceDirect

Journal of King Saud University – Science

journal homepage: www.sciencedirect.com



Review

A review on biological carbon sequestration: A sustainable solution for a cleaner air environment, less pollution and lower health risks

R. Gayathri ^a, Shahid Mahboob ^{b,*}, Marimuthu Govindarajan ^{c,d,*}, Khalid A. Al-Ghanim ^b, Zubair Ahmed ^b, Norah Al-Mulhm ^b, Masa Vodovnik ^e, Shankar Vijayalakshmi ^{f,*}^a School of Biosciences and Technology, VIT University, Vellore 14, India^b Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^d Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India^e University of Ljubljana, Groblje 3, 1230 Domžale, Slovenia^f CO₂ Research and Green Technologies Centre, VIT University, Vellore 14, India

ARTICLE INFO

Article history:

Received 4 November 2020

Revised 13 November 2020

Accepted 7 December 2020

Available online 17 December 2020

Keywords:

Carbon dioxide

Health effects

Utilization

Storage

Sequestration

ABSTRACT

Carbon dioxide gas is the key element of the carbon cycle and a major source for photosynthesis, but for the past 150 years, the atmospheric CO₂ has been increased drastically from 250 to 418 ppm due to the extreme utilization of fossil fuels. This accelerated release of CO₂ acts as a major source for climatic change due to the greenhouse gas effect resulting in global warming and melting of polar ice caps, alteration in biogeochemical cycles, altered rainfall, ocean acidification, eutrophication of lakes, imbalance in the ecological communities and extinction of some species, effects on soil fertility, changes in the metabolism and at the molecular level. Reduce, reuse and recycle strategy can be applied to control elevated CO₂ levels by preventing deforestation, using renewable energy as an alternative for fossil fuels and reusing the atmospheric CO₂. Carbon capture and storage (CCS), Carbon capture and utilization (CCU) are the two technologies adapted to capture the atmospheric CO₂, utilize it, and focus on permanent storage in the geological sites. Captured CO₂ is used to produce many value added products such as polymers, bio-fuels, reactants etc. Plants and microorganisms act as a natural CO₂ filter. Several biomolecules such as carbohydrates, proteins, and lipids are produced due to the biological carbon fixation process using photosynthesis. Six different photosynthetic pathways and some non-photosynthetic pathways to fix atmospheric CO₂ have been reported in diverse species of plants and microbes such as bacteria, fungi, yeast, algae etc. Algae are the most potent microbe in CO₂ utilization and biological carbon fixation compared to other microbes and used widely on a large industrial scale for biofuel production. Algal biofuel production using captured CO₂ is the best productive method to recycle and reduce atmospheric CO₂.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	2
2. Causes for CO ₂ emission	2
2.1. Natural emission of CO ₂	2

* Corresponding authors at: Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India. CO₂ Research and Green Technologies Centre, VIT University, Vellore 14, India.

E-mail addresses: mushahid@ksu.edu.sa (S. Mahboob), drgovind1979@gmail.com (M. Govindarajan), vijimicro21@gmail.com (S. Vijayalakshmi).

Peer review under responsibility of King Saud University.



<https://doi.org/10.1016/j.jksus.2020.101282>

1018-3647/© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



Review

A review on microbial degradation of drinks and infectious diseases: A perspective of human well-being and capabilities

Vijayalakshmi Shankar^a, Shahid Mahboob^{b,*}, Khalid A. Al-Ghanim^b, Zubair Ahmed^b, Norah Al-Mulhm^b, Marimuthu Govindarajan^{c,d}^aCO₂ Research and Green Technologies Centre, VIT University, Vellore 14, India^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 5 November 2020

Revised 20 November 2020

Accepted 10 December 2020

Available online 29 December 2020

Keywords:

Beverages

Microbial spoilage

Infectious diseases

Prevention

Human well-being

ABSTRACT

Various microorganisms present in beverages as contaminants, yet few can develop in the presence of acidic and low oxygen conditions. Notably, yeast is the dominant microbe group present in the fruit juices and drinks in the fruit juices and drinks. Due to the microbe development and secondary metabolite production, such as polluting compounds, carbon dioxide, and spoilage, the beverages were identified. Yeast and molds are considered to be a vital microbe that causes deterioration. The primary reason for the deterioration in fruits and fruit juices is because of the contamination caused by fungi and yeast, and sometimes damage may cause by insects. Sugars and sugar concentrates are generally polluted with osmophilic yeasts, for instance, *Z. rouxii*. This review mainly focuses on the types of spoilages in soft drinks and beverages and their control measures.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Contents

1. Introduction	2
2. Soft drinks	2
2.1. Spoilage of soft drinks by yeast	2
2.2. Bacterial spoilage	3
2.3. Fungal spoilage	4
3. Alcoholic beverages:	4
3.1. Yeast spoilage	4
3.2. Bacterial spoilage	4
3.3. Microbiological health risks linked with beverages	6
4. Presence of pathogenic bacteria	6
4.1. Presence of viruses and parasites	6
4.2. Mycotoxins in beverages	6
5. Other microbial metabolites in beverages	7
6. Factors microbial growth in beverages	8
6.1. pH and acidity	8

* Corresponding author.

E-mail address: mushahid@ksu.edu.sa (S. Mahboob).

Peer review under responsibility of King Saud University.





Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com



Original article

Green synthesis of ZnO nanoparticles for antimicrobial and vegetative growth applications: A novel approach for advancing efficient high quality health care to human wellbeing

Saraswathi Umavathi ^a, Shahid Mahboob ^{b,*}, Marimuthu Govindarajan ^{c,d}, Khalid A. Al-Ghanim ^b, Zubair Ahmed ^b, P. Virik ^b, Norah Al-Mulhm ^b, Muruges Subash ^e, Kasi Gopinath ^f, C. Kavitha ^a^a Adhyan Arts and Science College for Women, Uthangarai, Tamil Nadu 635207, India^b Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India^d Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India^e Arignar Anna Govt. Arts College, Attur, Tamil Nadu 636121, India^f School of Materials and Energy, Southwest University, Chongqing 400715, PR China

ARTICLE INFO

Article history:

Received 24 November 2020

Revised 9 December 2020

Accepted 13 December 2020

Available online 19 December 2020

Keywords:

Leaf extract

ZnO nanoparticles

Reactive oxygen species

Sesamum indicum

Health care

Human wellbeing

ABSTRACT

The present work aims to synthesize zinc oxide (ZnO) nanoparticles via green approaches using leaf extract of *Parthenium hysterophorus*. UV-vis and FT-IR tests confirmed the existence of biomolecules, active materials, and metal oxides. The X-ray diffraction structural study exposes the ZnO nanoparticles formation with hexagonal phase structures. SEM and TEM analysis reveal surface morphologies of ZnO nanoparticles and most of them are spherical with a size range of 10 nm. ZnO nanoparticles were revealed strong antimicrobial activity against both bacterial and fungal strains. The germination of seeds and vegetative growth of *Sesamum indicum* has been greatly improved.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Nanotechnology is a multidisciplinary scientific domain and has been used in various science fields, including chemical, physical, biological, pharmaceutical and material science (Porter and Youtie, 2009; Govindarajan et al., 2016a,b; Govindarajan, M. and Benelli, 2016, 2017; Balalakshmi et al., 2017; Divya et al., 2018; Fahimmunisha et al., 2020). The promising application of nanotechnology unlocked up a new scope and perspective in agriculture. The relatively small size, high surface to volume ratio and characteristics optical properties of nanomaterials find the

application from plant protection to nutrition and management practices in the farm (Shang et al., 2019). The perceptive of nanotechnology provides a new precision to agriculture with particular reference to fertilizer. The effects and efficiency of nanoparticle uptake on growth and metabolic activities may vary between the plants (Rastogi et al., 2017). The uptake concentration of nanoparticle influences the germination process and plant growth. Deficiency of zinc (Zn) is one of the major micronutrient problems affecting crop production, mostly calcium carbonate-rich alkaline soils (Takkar and Walker, 1993). The calcium carbonate abundant soils and alkaline pH may reduce both the obtainability and solubility of Zinc to the crops (Alloway, 2009; Rashid and Ryan, 2004). The Zn fertilizers such as zinc oxide (ZnO) and zinc sulphate were used to compensate the Zn deficiency in soils (Mortvedt, 1992) but were limited to their applications due to Zn non-availability to the plants. Apart, the application of chemical fertilizer leads to adverse effects on livestock, beneficial soil microorganisms and finally reduces soil fertility. In order to combat this problem, more effective and non-persistent fertilizer such as controlled release formulation is therefore required. ZnO nanoparticles

* Corresponding author.

E-mail address: mushahid@ksu.edu.sa (S. Mahboob).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Review

Microbial degradation of plastics: Sustainable approach to tackling environmental threats facing big cities of the future

S. Venkatesh^a, Shahid Mahboob^{b,*}, Marimuthu Govindarajan^{c,d}, Khalid A. Al-Ghanim^b, Zubair Ahmed^b, Norah Al-Mulhm^b, R. Gayathri^a, S. Vijayalakshmi^a^aCO₂ Research and Green Technologies Centre, VIT University, Vellore 14, India^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 21 December 2020

Revised 14 January 2021

Accepted 28 January 2021

Available online 10 February 2021

Keywords:

Microorganisms

Decomposition

Bioplastic

Solid wastes

ABSTRACT

Most microorganisms are used as a foundation of bioplastic production and also used for the decomposition of plastics. Although bioplastics production is considered expensive than artificial plastic, it has many advantages over them. Some bio-polymers have also gained public acceptance and are now being produced. The useful breakdown of plastic bags takes more than a thousand years. For the decomposition of plastics, microorganisms should be calculated extensively so that solid wastes can be decomposed. Thus, microbes have been played an important role in decompose as well as the production of plastics.

© 2021 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	1
2. Available methods for study of plastic biodegradation	2
3. Plastic degradation by bacteria and fungi	2
4. Microbial mechanism of PE degradation	5
5. Investigating the techniques involved in the bioremediation of plastic	6
6. Toxicity of biodegradable plastic	7
7. Conclusion	7
Declaration of Competing Interest	7
Acknowledgement	7
References	7

* Corresponding author.

E-mail address: mushahid@ksu.edu.sa (S. Mahboob).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

1. Introduction

Plastic usage has transformed our life in various ways. The production and utilization of plastics are always increasing due to the rising demand. They are inexpensive, strong, lightweight, corrosion-resistant, duration and electrical insulation properties and have high thermal (Aruna and Shanthi, 2015). Five hundred



Ulvan loaded graphene oxide nanoparticle fabricated with chitosan and D-mannose for targeted anticancer drug delivery

Sonia Kesavan^{a,**}, K.S Meena^b, S. Aruna Sharmili^c, Marimuthu Govindarajan^{d,e},
Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^g, Ahmed S. Alobaidi^f,
Khalid F. Alanzi^f, Baskaralingam Vaseeharan^{g,*}

^a Department of Chemistry, Queen Mary's College (Autonomous), Chennai, Tamil Nadu, 600004, India

^b Department of Chemistry, Alagappa Government Arts College, Karaikudi, Tamil Nadu, 630003, India

^c Department of Biotechnology, Stella Maris College (Autonomous), Chennai, Tamil Nadu, 600086, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^g Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

ARTICLE INFO

Keywords:

Graphene oxide
Chitosan
D-mannose
Ulva lactuca
Biocompatibility
Glioblastoma

ABSTRACT

Cancer growth is a developing significant public health threat, and notwithstanding the advances in biomedical research and innovation, a pressing need is felt for the progress of new anticancer drugs. Sulfated polysaccharides, such as ulvan from green macroalgae, exhibit a diverse range of biological applications. In this study, a novel D-mannose-mediated targeted drug delivery system (GO-CH-Ma) for targeting glioblastoma cancer was developed by loading *Ulva lactuca* as the anticancer model drug onto functionalized graphene oxide. Ultraviolet spectroscopy, Fourier-transform infrared spectroscopy (FT-IR), X-ray diffraction (XRD) and Zeta potential were used to describe the main physicochemical properties of the chitosan-functionalized graphene oxide (GO-CH). Similarly, scanning electron microscopy (SEM) and transmission electron microscopy (TEM) enabled the study of surface morphology. The entrapment of ulvan on GO-CH-Ma has been observed to be 88%. The biocompatibility of the nanocarrier and drug-loaded nanocarrier was studied via hemolysis and anti-inflammatory assay. The *in vitro* drug release profile of ulvan revealed a pH-dependent-controlled release system observed by UV-Visible analysis. Moreover, a human glioblastoma cell line (U87) was used to examine the preliminary *in vitro* cytotoxicity. Finally, a mannose-decorated GO-CH carrier loaded with ulvan demonstrated a promising targeted drug delivery system to treat *in vitro* glioblastoma.

1. Introduction

Nanotechnology has made a meaningful contribution to nanomedicine's advancement, thereby demonstrating extraordinary guarantees to improve restorative methodologies against various diseases. Nanomedicine speaks to a chance of focusing on procedures and multifunctionality. Despite the significant advancement in medicine, the majority of tumors are still incurable, and malignant gliomas being the most concerning. Additionally, glioblastoma (GBM) is the most malignant and exceptionally vigorous, having a place to evaluate IV gliomas as per the WHO [1,2].

Marine polysaccharides are a vital source of compounds with therapeutic applications because they are stable, inexpensive, biocompatible, non-toxic, safe, and biodegradable. Many researchers have studied polysaccharide-based nanomaterials for biomedical applications such as drug delivery, gene delivery, cancer therapy, wound dressing, and antimicrobial activity in recent years [3–5]. Green algae (Chlorophyceae), red algae (Rhodophyceae), and brown algae (Phaeophyceae) are the three major types of macroalgae. In particular, macroalgae are an excellent candidate for sulfated polysaccharides and have piqued the interest of cosmetics, nutraceuticals and pharmaceuticals [6].

Ulvan (UL) is a naturally occurring anionic sulfated polysaccharide

* Corresponding author.

** Corresponding author.

E-mail addresses: soniakcsavanmc2@gmail.com (S. Kesavan), vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jddst.2021.102760>

Received 25 March 2021; Received in revised form 1 July 2021; Accepted 24 July 2021

Available online 30 July 2021

1773-2247/© 2021 Elsevier B.V. All rights reserved.



Acute Toxicity and Histopathological Effect of Lambda Cyhalothrin and Pretilachlor on Intestine of Indian Earthworm *Lampito mauritii* (Kinberg)

Indhumathi.G¹, Kavitha.V^{2*} and Anandhan.R²

¹Research Scholar, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, India.

²Assistant Professor, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, India.

Received: 18 Feb 2022

Revised: 02 Mar 2022

Accepted: 25 Mar 2022

*Address for Correspondence

V. Kavitha.

Assistant Professor,

Department of Zoology, Annamalai University,

Annamalai Nagar, Tamil Nadu, India

Email: kavisamikshaa@gmail.com



This is an Open Access Journal / article distributed under the terms of the **Creative Commons Attribution License** (CC BY-NC-ND 3.0) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. All rights reserved.

ABSTRACT

Soil pollution has increased over time due to the widespread use of pesticides and chemical fertilizers in agriculture. Earthworms are important members of the soil ecosystem. In most ecological systems, earthworms play a crucial role in soil fertility and the formation of biogenic structures. Furthermore, earthworm gut mucus boosts the biological activities of beneficial soil microorganisms. The use of earthworms to enrich soil with essential nutrients is a cost-effective and environmentally friendly process. To control agricultural pests as well as weeds in India, the organophosphorus insecticide - lambda cyhalothrin and herbicide -pretilachlor are widely used. The present study the toxic effect of lambda cyhalothrin and pretilachlor of acute toxicity and histopathological effect reveals intestine of earthworm *L. mauritii* in a laboratory experiment. For acute study, mortality of *L. mauritii* was observed after 24, 48, 72, 96 and 120 h of the pesticides exposure. A sub-lethal concentration of lambda cyhalothrin and pretilachlor (1/5th of 96 h LC₅₀ value 1.8 ppm kg⁻¹ and 2.7 ppm kg⁻¹) was applied for 30 days. Histology of intestine was observed at 5th, 15th and 30th day of the pesticides exposure. On the 5th, 15th day of experiment, the skin showed significant pathological changes in epidermis layer, circular muscle, and longitudinal muscle. The changes such as vacuolization, degenerated nuclei, damaged epithelial lining of villi and congestion of blood sinuses was observed in 30th day, slight damages was observed. These results suggest that lambda cyhalothrin and pretilachlor could severely affect the intestine up to 15thday thereafter recovered of exposure when compared to 30th day. Histopathological study in *L. mauritii*'s intestine is a suitable parameter for detection of soil contamination by application of pesticides in agricultural field.



Article

Interplay between Binary and Three-Body Interactions and Enhancement of Stability in Trapless Dipolar Bose–Einstein Condensates

Sabari Subramaniyan¹, Kishor Kumar Ramavarmaraja², Radha Ramaswamy^{1,*} and Boris A. Malomed^{3,4,*}

¹ Centre for Nonlinear Science, Department of Physics, Government College for Women(A), Kumbakonam 612001, India; ssabari01@gmail.com

² Centre for Quantum Science, Dodd-Walls Centre for Photonic and Quantum Technologies, Department of Physics, University of Otago, Dunedin 9054, New Zealand; kishor.bec@gmail.com

³ Center for Light-Matter Interaction, Department of Physical Electronics, School of Electrical Engineering, Faculty of Engineering, Tel Aviv University, Ramat Aviv, Tel Aviv P.O. Box 39040, Israel

⁴ Instituto de Alta Investigación, Universidad de Tarapacá, Casilla 7D, Arica 1000000, Chile

* Correspondence: vittal.cnls@gmail.com (R.R.); malomed@tauex.tau.ac.il (B.A.M.)

Abstract: We investigate the nonlocal Gross–Pitaevskii (GP) equation with long-range dipole-dipole and contact interactions (including binary and three-body collisions). We address the impact of the three-body interaction on stabilizing trapless dipolar Bose–Einstein condensates (BECs). It is found that the dipolar BECs exhibit stability not only for the usual combination of attractive binary and repulsive three-body interactions, but also for the case when these terms have opposite signs. The trapless stability of the dipolar BECs may be further enhanced by time-periodic modulation of the three-body interaction imposed by means of Feshbach resonance. The results are produced analytically using the variational approach and confirmed by numerical simulations.

Keywords: Bose–Einstein condensates; Gross–Pitaevskii equation; dipole-dipole interaction; variational method; Runge–Kutta method; Crank–Nicolson method



Citation: Subramaniyan, S.; Ramavarmaraja, K.K.; Ramaswamy, R.; Malomed, B.A. Interplay between Binary and Three-Body Interactions and Enhancement of Stability in Trapless Dipolar Bose–Einstein Condensates. *Appl. Sci.* **2022**, *12*, 1135. <https://doi.org/10.3390/app12031135>

Academic Editor: Chiara D’Errico

Received: 13 December 2021

Accepted: 17 January 2022

Published: 21 January 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

The advent of Bose–Einstein condensates (BECs) in ⁵²Cr [1,2], ¹⁶⁴Dy [3,4] and ¹⁶⁸Er [5] accompanied by long-range dipole-dipole (DD) interactions superimposed on the contact inter-atomic collisions has impacted the investigation of ultracold quantum gases [6]. The anisotropic character and long-range nature of DD interactions endows the dipolar BECs (DBECs) with several distinct features such as the subordination of stability on the trap geometry [1,2], roton-maxon character of the excitation spectrum [7,8], new dispersion relations for elementary excitations [9,10], novel quantum phases [11,12], explicit [13,14] and hidden vortices [15], specific vortex-antivortex pairs [16], anisotropic multidimensional solitons [17,18], quantum droplets stabilized by beyond-mean-field effects [19–21], etc. The above phenomena arise due to the interplay between the contact *s*-wave interactions and the dipolar attraction or repulsion [22]. Tuning contact interactions by means of the Feshbach resonance is an important tool in analyzing the properties of DBECs [7,8]. In particular, DBECs with pure long-range interactions can be made by tuning the contact interactions to zero.

The theoretical description of a dilute weakly interacting DBEC is based on the Gross–Pitaevskii (GP) equation with the nonlocal DD-interaction term [1,2,23–25]. In particular, the combination of local and nonlocal terms in the GP equation can support various species of bright and dark matter-wave solitons. In the alkali BECs, bright solitons exist when the negative (attractive) contact interaction exactly balances the dispersion (kinetic-energy) term [26,27]. In DBEC, the nonlocal DD interaction term may reinforce local ones originating from the *s*-wave contact interaction. DD interactions are strongest in the atomic condensates

GENERAL LAND USE PATTERN IN THANJAVUR DISTRICT, TAMIL NADU: A GEOGRAPHICAL ANALYSIS

Kalyani. V, Assistant Professor, Department of Geography, Government College for Women (Autonomous), Kumbakonam – 612001.
Neelakantan. R, Professor, Department of Industries and Earth Science, Tamil University, Thanjavur - 613010.

E-Mail: kalyanisatheesh@gmail.com

Abstract

Thanjavur District is geographically located in the Cauvery delta area, and it is known as the "rice bowl" of Tamil Nadu State. Fruitful terrain was equally fertile ground for the creation of art, architecture, and culture. The study of the general land use pattern in Thanjavur District, Tamil Nadu, is an essential part of agricultural geography since it offers a solid foundation for the planning and development of the farming sector in the district. The agricultural industry is the most important source of revenue for the people of the Thanjavur District. Finally, this research discovers that the Thanjavur districts land use classifications have been determined taluk-by-taluk utilising the Geographic Information System (GIS). As a result, the current paper covers the overall land use pattern in Thanjavur District, Tamil Nadu: A Geographical Analysis.

Keywords: Land use/land cover, Geographic Information System (GIS), Thanjavur district.

Introduction

Agriculture is the science and art of producing plant life from the soil to benefit humans and other living things. It includes all the productive efforts he put into raising different products by increasing with other natural recourses to satisfy his needs and wants. Agriculture is the backbone of the Indian economy. In India, nearly 70 % of the total population and 90 % of the rural population is engaged in agriculture. Land use can be considered an important aspect of the land-man relationship, and it has been studied particularly relevant to agricultural geography (Yuvaraj et al., 2020; Gumma, 2015; Sujatha et al., 2011).

Numerous foreign and Indian geographers have interpreted the idea of land use differently. For, the land must be exploited to meet the legitimate wants and aspirations of the people as feasible (NRSA, 2006, 2007). Essentially self-explanatory, the phrase land use refers to the real and precise usage of an earth's surface in terms of natural main land use,

such as the land beneath the forest, grassland, farming, or other similar land uses, as opposed to other land uses. The term "land usage" refers to all industrialised and developing areas on their surface, for a specific point at a given time and space (Arulbalaji, 2019; Gumma et al., 2015; Freeman, 1968). Land use is any permanent or cyclic human intervention on the environment to satisfy human needs (Ganeshmoorthi, 2018; Prabu and Dar, 2018; Sangeetha and Baskaran, 2011). Land use appropriateness is a tract's potential to accommodate various land use patterns under specific cultural and socio-economic circumstances (Lenin Sundar et al., 2019; Vink, 1975). Symons (1978) states that land-use maps are crucial instruments for regional development and planning. (Krishnan, 2014).

Thanjavur district has been known for its agricultural activity from ancient times. The Granary of Tamil Nadu is in the deltaic area of the famed river Cauvery and is criss-crossed by a vast network of irrigation systems and irrigation systems. Tropical vegetation abounds in this coastal area. An important part of agricultural geography is the study of general land-use patterns in the Thanjavur District.

The Study Area

Located in southern India, Thanjavur is a significant centre of art and architecture. The city of Thanjavur is home to the majority of the magnificent living Chola temples, which are all designated as UNESCO World Heritage Sites. On the eastern coast of the Indian state of Tamil Nadu, Thanjavur district is situated. Its latitude ranges from 90° 50' to 11° 25' north, while its longitude ranges from 78° 45' to 70° 25' east. It has a total area of 3,411 square kilometres (Fig.1). The Coloroon River separates the district from the Perampalure and Thiruchirappalli districts to the north. It is bordered on the east by the Thiruvarur and Nagapattinam districts, on the south by the Palk Strait and Pudhukottai district, and on the west Pudukkottai and Thiruchirappalli districts. It is also bordered on the east by the Thiruvarur and Nagapattinam districts. Thanjavur district is divided into three revenue divisions with eight taluks with fourteen blocks and three municipalities.



SURVEY OF MARINE TOURISM AND AROUND NAGAPATTINAM DISTRICT – A STATISTICAL ANALYSIS

Abstract

The Marine tourism industry has different characteristics and it has a hospitality business. This study examines the various components of socio-economic and coastal tourism in Nagapattinam, Tamil Nadu and the perception of tourism people about the ecological status of tourism. The researcher concluded that the primary data from the 150 respondents in the study area were obtained through interview methods, as well as using statistical tools such as SPSS techniques and frequency and communication. The findings of the study suggest that it is necessary to make the contribution and growth of residents more sustainable.

Keywords- Coastal Tourism, Socio-Economic, Environment, Development.

1. Introduction

Tourism is one of the largest industries in the world. The coastline of Tamil Nadu is about 1076 kms, which is 15% of the total coastline of India, stretching along the Bay of Bengal, the Indian Ocean and the Arabian Sea. It is one of the longest coastlines of 187.9 kms and the second longest coastline. One of the most common types of tourism is coastal tourism. It is strongly dependent on natural and cultural aspects. Coastal tourism has become the main face of modern life. Coastal tourism is a process that attracts tourists, especially coastal ecosystems. Most coastal tourism takes place on the coast and in the immediate vicinity of the coast. The growth of tourism in general and in coastal areas in particular is related to three main factors: increased personal incomes, leisure time and improvements in transportation system and greater public awareness of world destinations due to improved communications (EEA, 2001). We can examine the interdependent relationships between tourism and the environment in terms of the interaction of the visitor, the place and the host community (UNWTO, 2004). Sustainability principles refer to the environmental, economic and socio-economic aspects of tourism development and a suitable balance must be established between these three dimensions to guarantee its long-term sustainability (UNEP & UNWTO, 2005). This growth, which has reached its peak in recent decades exert pressures on the environmental and cultural resources of coastal areas patterns of tourist destinations (Marina & Alessio, 2009).

2. Study area

The present study area is located in the eastern part of Tamil Nadu and Nagapattinam is one of the coastal districts of the state. It is a peninsular delta district, bordered on the east by the Bay of Bengal, on the south by the Balk Strait and coastal plain land on the west and north. The district is geographically located between 10° 10' and 11° 50' North latitude and between 79° 30' and 79° 50' East longitude. The total

CUSTOMER ACHIEVEMENT ALONG WITH LIFE INSURANCE CORPORATION

G. ABIRAMY, M.Com M.Phil.
GUEST LECTURER, PG & RESEARCH DEPARTMENT OF COMMERCE
GOVT. COLLEGE FOR WOMEN (A), KUMBAKONAM
(AFFILIATED TO BHARATHIDASAN UNIVERSITY)

Dr. P. HEMALATHA., M.Com. M.Phil., M.Ed. Ph.D.
Research Supervisor & Principal,
Raja Doraisingam Government Arts College,
Sivagangai.

ABSTRACT:

This study investigates achievement level of clients towards Life Insurance Corporation in Chennai city. Because of expanding mindfulness among individuals about their vulnerability of life and expanding rivalry in Insurance area, it is critical for Insurance Companies to comprehend the essential of their clients. The primary point of this investigation is to know the purchaser achievement towards LIC. This investigation depends on essential information which is gathered through survey among 160 policyholders in Chennai and information were broke down with factual device like rate is utilized to recognize the components in charge of buyer achievement towards LIC.

Keywords: LIC; Consumer Achievement; Consumer Awareness.

1. INTRODUCTION:

Protection business wad quickly developed in eighteenth century. Individuals are shaky towards dangers and vulnerabilities. Time of death of an individual isn't sure. Each judicious man is progressively wary about keeping them from hazard or limiting or giving impact against it. Protection segment have made conceivable to cover the hazard which is inconceivable. Insurance agencies began to move the hazard and hazardous to the interpretation of who are happy to acknowledge the weight for monetary thought. The principle goal of protection organization is to dispense with the hazard on the substitution conviction for dubious and in this manner it makes collaboration among the policyholders for spreading their hazard. Because of expanding rivalry in Insurance division and expanding desire for policyholder's prompts measure the mindfulness and achievement level of the clients. LIC began to give more administrations so as to fulfill the necessity of the client and for their prosperity and survival in the Insurance Sectors. Because of progress in taste and inclination of the policyholder it is getting to be hard for the insurance agency to comprehend and quantify the administration viably.

Financial Performance Of Oriental Insurance Company Ltd

G. Abiramy,

m.com m.phil.

Guest lecturer, pg & research department of commerce
Govt. College for women (a), kumbakonam
(affiliated to bharathidasan university)

Dr. P. Hemalatha.,

m.com. M.phil., m.ed. Ph.d.

Research supervisor & principal,

Raja doraisingam government arts college,

Sivagangai.

Abstract

Insurance is been an vital part of the Indian Financial system, after 17 years of Liberalization of Insurance Industry in India, it has taken a new trajectory as many life insurers have managed to list themselves in the Indian bourses. The Indian Insurance is growing faster since 2000, as it has been liberalized after more than 50 years, where private life insurers have entered the insurance sector with innovative practices leading to more business as it clearly needs a lot of finance that can now be mobilized through the cleaner equity capital from a longer perspective. The Oriental Insurance Company actions adds to transparency, governance, accountability after liberalization of sector and also to improve the focus towards profitability of the Indian Insurance Industry. This study attempts to measure the financial performance of oriental insurance company ltd.

Keywords: Life Insurers, Financial system, Profit after tax, Net premium, Net Claim, Investment Income, financial performance

Introduction

Insurance is the backbone in managing the risk of the country. The insurance providers offer diversity of products to business, providing protection from risk thereby ensuring financial security. It helps individual and organization to minimize the consequences of risk which impart significant cause on the growth and development of insurance industry. Indian insurance industry is facing major challenges in reaching out willing customers, providing them services, acquiring and retaining players, product and distribution innovation etc. Apart from addressing the challenges of customers, improving the performance to achieve profitable growth is another big challenge faced by Indian life insurers. To sustain the profitable growth, private companies are struggling in spreading awareness about need of insurance, developing brand strength, meeting regulatory demands, establishing wide network of distribution channels and setting infrastructure. Oriental insurance sector anticipate different segments of customers with different needs thereby raising the importance of new and competitive dynamics.

Evaluation of Budget Analysis of Kumbakonam Municipality

Mrs.M.Banumathi,

Guest Lecturer,

PG& Research Dept. of Commerce

Govt. college for Women (A),

Kumbakonam. -612001

E-mail. id : mbanu2615@gmail.com

Dr. Mrs. P. Hemalatha, M.Com,

M. phil., M.Ed., Ph. D,

Research Supervisor & Principal,

Raja Doraisingam Government Arts College,

Sivagangai-630561.

(Affiliated to Bharathidasan University)

Abstract:

The financial structure of Urban Local Bodies in India is mainly comprises of tax and non-tax revenues as major source of revenues and these local bodies also receive grants-in aid and other concessions from the Government. The expenditure of the municipalities are classified as establishment expenditures, administration expenditures, operating, repairs & maintenance expenditures finance expenses and depreciation. Therefore this study is focussed on the evaluation of budget analysis of Kumbakonam Municipality using trend analysis using actual and estimated values. For further study Chi-square test is used.

Key words: Urban Local Bodies, Source of revenues, Grants-in aid.

Introduction

Financial Management is used in each and every organisation, whether it is trading or non-trading organisation. Finance is needed to promote or establish business, acquire fixed assets, make investigations such as market survey, develop product, keep men and machine at work It is necessary as it guides in

- Financial planning and successful promotion of an enterprise,
- In acquisition of funds at a minimum cost,
- In efficient usage and allocation of funds,
- In taking sound financial decisions and
- In promoting and mobilising individual and corporate savings.

Urban Local Government(ULG):



AN ENHANCED ANT COLONY OPTIMIZATION BASED TASK SCHEDULING ALGORITHM FOR CLOUD COMPUTING

E.Suganthi¹ **Dr.F.KurusMalaiSelvi**²

¹Research Scholar, PG&Research Department of Computer Science ,
Govt College for Women(A), Kumbakonam, Bharathidasan University, India
E-mail: e.suganthi@gcwk.ac.in

²Associate Professor & Head , PG&Research Department of Computer Science ,
Govt College for Women(A), Kumbakonam, Bharathidasan University, India
E-mail: f.kurusmalaiselvi@gcwk.ac.in

Corresponding author: E.Suganthi(e.suganthi@gcwk.ac.in)

8627

ABSTRACT

Cloud computing is a dispersed collection of resources that is made available to users. It provides users with nearly unlimited pay-per-use computing services without the responsibility for the underlies. The aim of cloud service providers is to allow effective use of resources and optimise benefit. These results in task planning, in cloud computing, to be a central and challenging problem. Cloud task forecast is an NP-hard optimisation problem, which has been suggested for solving by several meta-heuristic algorithms. A successful task manager should adapt his planning approach to evolving conditions and task styles. This research paper proposes a advance Ant colony optimization algorithm by adding up ant onlookers for the adaptation and diversification of strengthening strategies with the added ants.

KEYWORDS: Cloud Computing, Task Scheduling, Optimization technique, Ant Colony Optimization, Multi-Objective

DOI Number: 10.14704/nq.2022.20.6.NQ22854

NeuroQuantology 2022; 20(6): 8627--8634

1. INTRODUCTION

The end user working in cloud environment is allowed to access the entire stack of computer resources, from hardware to software. Cloud storage systems are on pay-as-you-go basis. Depending on applications demands, the service user may lessen or increase the usage of resources available. This is solitary benefit of cloud computing [1],[2] but the expense for this benefit should be paid by service users. The customer of the cloud service can charge and release resources at any time. Any service that is based on application specifications is free for the cloud service customer. When the user does not select a correct service it will create a great problem. Therefore, the planning of tasks and allocation of resources are compulsory components of cloud science. The resource use efficiency depends not on the

random allocation of resources, but on the planning and load balance methodologies. Computers in the cloud are used extensively to solve complex tasks (user requests). The use of the scheduling algorithm is recommended for solving complex task problems. The tools are leveraged in such planning algorithms.

Task scheduling is one of the intricate issue in cloud computing. An effective use of resources may lead to an adequate scheduling of tasks. The greatest value of cloud computing is the optimum usage of resources [3]. Distribution of resources and task scheduling are thus two sides of one coin. Each one has an impact on the other. Without considering hosting infrastructure, users on the internet, currently have access to content from anywhere, anyplace and anytime. Many computers form part of the hosting



Analysis of Socio, Economic Impact on Tourism Development in Nagapattinam District using in SPSS

¹Nithya S* AND ²Neelakantan R

¹Research Scholar

Department of Industries and Earth Sciences Tamil University, Thanjavur, Tamil Nadu, India

and

²Professor

Dean, Faculty of Science

Department of Industries and Earth Sciences Tamil University, Thanjavur, Tamil Nadu, India

ABSTRACT

The present paper analyze about the tourism of Nagapattinam district with various dimensions. "Coastal and marine tourism includes those recreational activities which involve travel away from one's place of residence which have as their host or focus the marine environment and/or the coastal zone." ICMTS. The present study analyzed using the Simple statistical methods. Chi square Test, ANOVA Test and SPSS Factor analysis. Nagapattinam is a developing district and it also a major pilgrimage centre for Hindu, Christians and Muslim. So the district has the religious values. It also having the coastal tourist spots of Kodyakkarai, Tharangambadi, Poombuhar and Velankanni. From the findings there is significant development in economy of Nagapattinam district due to tourism. In general, Nagapattinam district has enormous the potential to attract tourists which will generate employment opportunities economic status of the people.

Key words: Tourism, Development, Determining factors, social impact.

INTRODUCTION:

Tourism is a Major industry and it plays a essential role in the economic progress of a country. It gives immense pleasure to the people. So people desire to seen various places and have interest to see different places. The number of Foreign Tourist Arrivals (FTAs) in India during 2018 increased to 10.56 million as compared to 10.04 million in 2017. The growth rate in FTAs during 2018 over 2017 was 5.2% as compared to 14.0% during 2017 over 2016 (India Tourism Statistics 2019. According to India Tourism Statistics 2019, "Tourism continues to play an important role as a foreign exchange earner for the country. In 2018, foreign exchange earnings (FEE) from tourism were US\$ 28.59 billion as compared to US\$ 27.31 billion in 2017, registering a growth of 4.7%." According to United Nations World Tourism Organization (2020, "Travel as the activity of moving between different locations offense for any purpose but more so for

“member’s satisfaction on salary and wage benefits with trade unions - at tnstc kumbakonam division “

Turkish Online Journal of Qualitative Inquiry (TOJQI)
Volume 12, Issue 10, October 2021: 2032-2036

“MEMBER’S SATISFACTION ON SALARY AND WAGE BENEFITS WITH TRADE UNIONS - AT TNSTC KUMBAKONAM DIVISION “

1R. Kalaimathi, 2Dr. P. Hemalatha

1Research Scholar in Commerce, Govt. College for Women (Autonomous), Kumbakonam, Affiliated to Bharathidasan University, Tiruchirappalli, Tamil Nadu

2Principal & Research Supervisor, **Raja Doraisingam Govt.Arts College, Sivagangai**, Affiliated to Bharathidasan University, Tiruchirappalli, Tamil Nad

Abstract

The existence of a strong and recognized trade union is a prerequisite to industrial peace. Decisions taken through the process of collective bargaining and negotiations between employer and unions are more influential. Trade unions play an important role and are helpful in effective communication between the workers and the management. A union acts as an intermediary between employees and Corporation and provides a necessary complement to legislated benefit and protection. Unions bargain with management for the successful increase workers salary and wages benefits through collective bargaining. The impact of unions on worker's wage and salary benefits therefore, the aim of the study is to analyze member’s satisfaction salary and wage benefits with the activities of trade unions in the TNSTC kumbakonam division Tamil Nadu. The result of the study reveals that among the skilled, semi-skilled and unskilled workers, the majority of the skilled workers perceive moderate satisfaction, whereas the majority of the semi-skilled and unskilled workers perceive a high level of satisfaction with the activities of the trade unions in the study unit.

INTRODUCTION

A trade union is an organized association of workers formed to protect their rights and interest. In the rapid programme of industrial development of India and with the implementation of new economic policy, trade unions have come to occupy a critical position in the success of industrial relations in the country. The Trade Unionism in India has been undergoing rapid changes due to socioeconomic transformation also. It is an important instrument to promote class collaboration and harmony. They are found as an instrument of solving social economic, political and psychological problems of large scale industry, machine, technology and mass production. Trade unions can reshape the relationship between the employers and employees in order to maintain an atmosphere of industrial peace and harmony. Trade unions play a significant role in securing the nature of the job and enforcing the rights on the job. Unions act as an intermediary between employees and Corporation and provide a necessary complement to legislated benefit and protection. Unions bargain with management for the successful increase of worker wages, other benefits through collective bargaining. The impact of unions on worker's wage and salary benefits is presented in Table 4.11

“an impact and level of collective bargaining system of tnstc employees and its trade union – kmbakonam division”

Turkish Online Journal of Qualitative Inquiry (TOJQI)
Volume 12, Issue 10, October 2021: 2023-2031

“AN IMPACT AND LEVEL OF COLLECTIVE BARGAINING SYSTEM OF TNSTC EMPLOYEES AND ITS TRADE UNION – KMBAKONAM DIVISION”

¹R. Kalaimathi, ²Dr. P. Hemalatha

¹Research Scholar in Commerce, Govt. College for Women (Autonomous), Kumbakonam, Affiliated to Bharathidasan University, Tiruchirappalli, Tamil Nadu

²Principal & Research Supervisor, Raja Doraisingam Govt.Arts College, Sivagangai, Affiliated to Bharathidasan University, Tiruchirappalli, Tamil Nadu

Abstract

This Paper addresses the collective bargaining of TNSTC trade union members, the future of collective bargaining, the conditions necessary for successful collective bargaining, some of the advantages of collective bargaining, issues of concern for employers and guidelines for employers on the process of bargaining itself. Collective bargaining is specifically an industrial relations mechanism or tool, applicable to the employment relationship as a process. In collective bargaining the union always has a collective interest since the negotiations are for the benefit of several employees. Where collective bargaining is not for one employer but for several, collective interests become a feature for both parties to the bargaining process. In collective bargaining certain essential conditions need to be satisfied, such as the existence of the freedom of association and a labour law system. collective bargaining took a new turn as trade unions were forced to examine seriously their approach to the preparation of proposals and the presentation of their case to bargain collectively is the performance of the mutual obligation of the employer and the representative of the employees to meet at reasonable times and confer in good faith with respect to wages, hours, and terms and conditions of employment,

Keywords: TNSTC employees, Collective bargaining, Transport employees.

CONCEPT OF COLLECTIVE BARGAINING

TNSTC is the second largest government bus transport corporation in India and is well known for its route coverage almost too every remote area within Tamil Nadu. In Tamil Nadu, TNSTC (Kumbakonam) Ltd accounts major stake. Transportation service here is more than 40 years old and is the back bone of the economy of this division. In this study area, majority of the TNSTC workers are getting membership in various trade unions. So that the researcher concentrate the trade union’s collective bargaining system. The phrase “*collective bargaining was first coined by Sidney and Beatrice Webb*. This was widely accepted, particularly in the developed countries. Generally by collective bargaining we mean, an essential element of economic democracy, is a ‘two party’ procedure for arriving at a commonly agreed solution. The term is thus used to describe the procedure, whereby employers must attempt to reach agreement about wage-rates and basic conditions of labour with trade unions, instead of with individual workers. In other words, it is the process of discussion and negotiation between an employer and a union culminating in a written



In vitro evaluation of cytotoxicity, anti-inflammatory and wound healing properties of otolith stones from pomadasys maculates

C. Kanimozhi¹, S. Ganesan², G. Gokila³, and R. Manimekalai^{1,*}

¹PG and Research Department of Physics, A.V.V.M. Sri Pushpam College (Affiliated to Bharathidasan University, Tiruchirappalli), Poondi, Thanjavur, Tamil Nadu 613 503, India

²PG and Research Department of Zoology, A.V.V.M. Sri Pushpam College (Affiliated to Bharathidasan University, Tiruchirappalli), Poondi, Thanjavur, Tamil Nadu 613 503, India

³Department of Physics, PRIST University, Vallam, Thanjavur, Tamil Nadu 613 403, India

Received: 16 November 2021

Accepted: 27 May 2022

Published online:

27 June 2022

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2022

ABSTRACT

Pomadasys maculatus (Blotched-grunt, saddle grunter) generally, often known as the “Dhoter,” is a commercially significant fish of the coast of India-Pakistan. Fishes are high in nutrients such as polyunsaturated fatty acids, amino acids, vitamins, and minerals. It is important in the prevention and treatment of coronary heart disease, asthma, eye problems and vitamin shortages. The natural crystalline material collected from the head of the Pomadasys maculatus fishes are used for various characterization techniques such as single crystal XRD, powder XRD, EDAX, SEM, FT-IR and FT-Raman, and the other biological properties of the Otolith stones are analyzed. The unit cell characteristics of the naturally collected otolith stones were determined using single crystal X-ray diffraction. The FT-IR and FT-Raman spectra investigated the various functional groups found in the otolith stones. The current study also sought to investigate the cytotoxicity, anti-inflammatory and wound healing properties of otolith stones found in Pomadasys maculatus fishes. One of the goals of this study is to determine whether otolith stones have cytotoxic effects on fibroblast cells. The inhibitory effects of otolith stones in LPS-stimulated RAW 264.7 macrophages were used to assess their anti-inflammatory properties. The wound scratch assay assured the otolith stone’s wound healing efficacy. The EDAX analysis exhibits the various constituents present in otolith stones.

Handling Editor: Annela M. Seddon.

Address correspondence to E-mail: maniabi64@gmail.com

<https://doi.org/10.1007/s10853-022-07385-4>



Robustness of Measurement–Induced Correlations Under Decoherence Effect

S. Bhuvaneswari¹ · R. Muthuganesan² · R. Radha¹

Received: 29 December 2020 / Accepted: 7 May 2021 / Published online: 26 May 2021
© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

In this article, we study the dynamics of quantum correlation measures such as entanglement and measurement–induced nonlocality (MIN). Starting from an arbitrary Bell-diagonal mixed state under Markovian local noise such as bit-flip, depolarizing and generalized amplitude damping channel, we provide the decays of the entanglement measured by concurrence and quantum correlation captured by different forms of MIN (trace distance, Hilbert–Schmidt norm and relative entropy) as a function of the decoherence parameters. The effect of local noises on the dynamical behaviors of quantum correlation is also observed. We show the existence of specific and important features of MIN such as revival, noise robustness and sudden change with respect to decoherence parameter. It is observed that all the noises cause sudden death of entanglement for partially entangled states. Further, we show the existence of separable quantum states with non-zero quantum correlations in terms of MIN.

Keywords Entanglement · Quantum correlation · Dynamics · Decoherence

1 Introduction

Nonlocality, a peculiar property of quantum system plays an important role in understanding the characteristics of composite quantum systems and makes a fundamental departure

✉ R. Radha
vittal.cnls@gmail.com

S. Bhuvaneswari
bhuvanajkm85@gmail.com

R. Muthuganesan
rajendramuthu@gmail.com

¹ Centre for Nonlinear Science (CeNSc), Government College for Women (A), Kumbakonam, 612 001, Tamil Nadu, India

² Centre for Nonlinear Science & Engineering, School of Electrical, Electronics Engineering, SASTRA Deemed University, Thanjavur, Tamil Nadu 613 401, India



Contents lists available at ScienceDirect

Physica A

journal homepage: www.elsevier.com/locate/physa

Spotlighting quantum phase transition in spin -1/2 Ising–Heisenberg diamond chain employing Measurement-Induced Nonlocality



S. Bhuvaneswari^a, R. Muthuganesan^b, R. Radha^{a,*}

^a Centre for Nonlinear Science (CeNSc), PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

^b Centre for Nonlinear Science & Engineering, School of Electrical and Electronics Engineering, SASTRA Deemed University, Thanjavur, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 1 October 2020

Received in revised form 12 January 2021

Available online 19 March 2021

Keywords:

Quantum correlation
Phase transition
Diamond chain
Nonlocality

ABSTRACT

We examine thermal quantum correlations characterized by Measurement-Induced Nonlocality (MIN) in an infinite spin-1/2 Ising–Heisenberg spin chain with Dzyaloshinskii–Moriya (DM) interaction. We evaluate MIN analytically in the thermodynamic limit using the transfer matrix approach and show that the MIN and its first-order derivative may spotlight the quantum criticality and quantum phase transition (QPT). We observe that the DM interaction reduces the role of anisotropy parameter in initiating QPT. Further, the DM interaction also induces the nonlocality in the system if the spins are unentangled and greatly enhances the quantum correlations if the spins are correlated. The impact of the magnetic field and temperature on quantum correlations is also brought out at a critical point.

© 2021 Elsevier B.V. All rights reserved.

1. Introduction

Nonlocality, a unique and fundamental characteristic feature of composite quantum systems [1–3], has been proven to be a useful resource for real-life applications, notably in the domain of secure communication, cryptography, etc. In the realm of Bell nonlocality [4], the violation of any Bell inequality can manifest itself as nonlocality and entanglement can be recognized as one of the most important signatures of nonlocal aspects of quantum states. Any pure entangled two-qubit state violates the Clauser, Horne, Shimony, and Holt (CHSH) inequality, while it is not true for mixed states [5]. Werner showed that the entanglement is not a complete picture of nonlocality [6]. Further, the deterministic quantum computation with one qubit (DQC1) [7] protocol is demonstrated through the separable state (zero entanglement), implying that there exists a special kind of property beyond entanglement, which is also efficient in solving some classically intractable problems. In recent times, the detailed investigation on these domains reveal that the entanglement does not measure the quantum correlations present in the quantum system completely [7]. To address this issue, various measures have been identified to measure the quantum correlations which cannot be grasped by the entanglement [8–12].

In particular, Luo and Fu introduced a new measure called Measurement-Induced Nonlocality (MIN) [11], which is based on the fact that local disturbance due to von Neumann projective measurements on marginal state can influence globally. Under these local measurements, the marginal states are invariant, and due to the local invariance, MIN is

* Corresponding author.

E-mail address: vittal.cns@gmail.com (R. Radha).

LOCALIZED EXCITATIONS AND THEIR COLLISIONAL DYNAMICS IN (2+1)-DIMENSIONAL BROER-KAUP-KUPERSHMIDT EQUATION

R. RADHA^{1,*}, C. SENTHIL KUMAR²

¹Centre for Nonlinear Sciences, Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam – 612001, India

*Corresponding author Email: vittal.cnls@gmail.com

²Department of Physics, Vinayaka Mission's Kirupananda Variyar Engineering College, Vinayaka Mission's Research Foundation (Deemed to be University), Salem – 636308, India

Compiled September 28, 2021

In this paper, we investigate the (2+1)-dimensional Broer-Kaup-Kupershmidt equation employing the truncated Painlevé approach and we generate localized solutions like dromions, lumps, and rogue waves. The collisional dynamics of the localized solutions is then studied. While the dromions are found to undergo inelastic collision exchanging energy among them, the lumps are found to be noninteracting. We also bring out the unstable nature of rogue waves.

Key words: Truncated Painlevé approach, Dromions, Rogue waves, Lumps.

1. INTRODUCTION


The revival of interest in the investigation of (2+1)-dimensional nonlinear partial differential equations (PDEs) can be attributed to the identification of rogue waves [1–10]. In general, rogue waves occur in deep ocean. The most important characteristic feature of rogue waves is that they appear from nowhere and disappear without a trace. In recent times, rogue waves have been identified in various fields such as hydrodynamics [11], nonlinear optics [12], Bose-Einstein condensates [13], and plasma physics [14]. Dromions [15–20], which are exponentially localized solutions [21–26], are another interesting classes of solutions in (2+1) dimensional integrable systems. They essentially arise by virtue of coupling of the field variable to a potential. In addition to rogue waves and dromions, there exist lumps that are algebraically decaying solutions in (2+1)-dimensional nonlinear PDEs [27–31]. While dromions have been known to undergo both elastic and inelastic collisions, lumps do not interact at all. On the other hand, rogue waves are found to be highly unstable. In addition to the above, (2+1)-dimensional nonlinear PDEs also admit traveling wave solutions [32], breathers [33, 34], etc. The contrasting nature of these localized solutions makes the investigation of (2+1) nonlinear PDEs an interesting journey.

In this paper, we investigate the (2+1)-dimensional Broer-Kaup-Kupershmidt (BKK) system [35] employing the truncated Painlevé approach and generate lo-



OPEN

Bayesian Optimization of Bose-Einstein Condensates

Tamil Arasan Bakthavatchalam^{1,4}, Suriyadeepan Ramamoorthy^{2,4},
Malaikannan Sankarasubbu^{2,4}, Radha Ramaswamy^{3,4} & Vijayalakshmi Sethuraman^{1,4}

Machine Learning methods are emerging as faster and efficient alternatives to numerical simulation techniques. The field of Scientific Computing has started adopting these data-driven approaches to faithfully model physical phenomena using scattered, noisy observations from coarse-grained grid-based simulations. In this paper, we investigate data-driven modelling of Bose-Einstein Condensates (BECs). In particular, we use Gaussian Processes (GPs) to model the ground state wave function of BECs as a function of scattering parameters from the dimensionless Gross Pitavetskii Equation (GPE). Experimental results illustrate the ability of GPs to accurately reproduce ground state wave functions using a limited number of data points from simulations. Consistent performance across different configurations of BECs, namely Scalar and Vectorial BECs generated under different potentials, including harmonic, double well and optical lattice potentials pronounces the versatility of our method. Comparison with existing data-driven models indicates that our model achieves similar accuracy with only a small fraction ($\frac{1}{50}$ th) of data points used by existing methods, in addition to modelling uncertainty from data. When used as a simulator post-training, our model generates ground state wave functions 36 x faster than Trotter Suzuki, a numerical approximation technique that uses Imaginary time evolution. Our method is quite general; with minor changes it can be applied to similar quantum many-body problems.

At ultra-cold temperatures, when a gas is dilute enough such that the interactions are weak, the atoms crash into the ground state, leading to an exotic state of matter called Bose-Einstein Condensates (BECs)¹. BECs are in general governed by the mathematical model described by Gross-Pitaevskii equation (GPE), a variant of Nonlinear Schrödinger equation (NLSE). In GPE, there are two variants, namely nonlinear interaction parameter (coupling strength) and the trapping potential and both could be spatio-temporal in general. GPE in its most general form cannot be solved analytically. It can be solved analytically² only for specific choices of coupling strengths and trapping potentials. In other cases, one has to resort to numerical techniques³ like Crank-Nicholson Scheme, Trotter-Suzukii Approximation, Finite Element Analysis, etc.

Numerical techniques rely on numerical differentiation which is the finite difference approximation of derivatives using values of the original function evaluated at some sample points. Numerical approximations of derivatives are inherently ill-conditioned and unstable due to introduction of truncation and round-off errors⁴. On the other hand, Automatic Differentiation (AD) also called algorithmic differentiation is a family of techniques for efficiently and accurately evaluating derivatives of numeric functions expressed as computer programs⁵. AD exploits the fact that every computer program executes a sequence of elementary arithmetic operations and elementary functions. By applying the chain rule repeatedly to these operations, derivatives of arbitrary order can be computed automatically and accurately to working precision. AD is both efficient and numerically stable compared to Numerical Differentiation. AD has become the beating heart of Modern Machine Learning, leading to a new paradigm of programming called Differentiable Programming.

In recent years, Scientific Computing community has started adopting data-driven approaches from Machine Learning, as faster and efficient alternatives to numerical simulation techniques. This new regime called Data-driven Scientific Computing has shown great promise already by faithfully modelling physical phenomena using scattered noisy observations from coarse-grained grid-based numerical simulations. The learned models are capable of predicting the dynamics of the system under study in a fixed number of CPU cycles irrespective

¹Department of Physics, Presidency College (Autonomous), University of Madras, Chennai 600005, India. ²Saama AI Research Lab, Chennai 600032, India. ³Post-Graduate and Research Department of Physics, Centre for Nonlinear Science (CeNSc), Government College for Women (Autonomous), Kumbakonam 612001, India. ⁴These authors contributed equally: Tamil Arasan Bakthavatchalam, Suriyadeepan Ramamoorthy, Malaikannan Sankarasubbu, Radha Ramaswamy, and Vijayalakshmi Sethuraman. [✉]email: tamilarasanbakthavatchalam@gmail.com



Modulational instability of spin-orbit coupled Bose-Einstein condensates in discrete media

S. Sabari^a, R. TamilThiruvalluvar^b, R. Radha^{a,*}

^a Centre for Nonlinear Science, Department of Physics, Government College for Women (A), Kumbakonam 612001, India

^b Department of Physics, Pondicherry University, Puducherry - 605014, India

ARTICLE INFO

Article history:

Received 23 May 2021

Received in revised form 12 September 2021

Accepted 13 September 2021

Available online 23 September 2021

Communicated by B. Malomed

Keywords:

Bose-Einstein condensates

Spin-orbit coupling

Modulational instability

Discrete media

Linear stability analysis

ABSTRACT

We address the impact of intra-site spin-orbit (SO) coupling and associated inter-component Rabi coupling on the modulational instability (MI) of plane-wave states in two-component discrete Bose-Einstein condensates (BECs). Conditions for the onset of the MI and the respective instability are found analytically. SO coupling allows us to produce the MI even for a small initial wavenumber ($q < \pi/2$) for miscible states. In particular, SO coupling introduces MI even in the absence of hopping coefficient, a concept which may have wider ramifications in heavy atomic BECs. We have also shown how our results of the linear stability analysis can be corroborated numerically. The fact that we have brought out the stability criteria in different domains of system parameters means that our model is tailor made experiments.

2021 Elsevier B.V. All rights reserved.

1. Introduction

Even though the ubiquitous nature of the spin-orbit (SO) coupling, which involves the interaction between quantum particle's spin and its momentum is quite well known, it came to the fore only after the experimental realization of Bose-Einstein condensates (BECs), a new state of matter comprising of ultracold atoms getting piled up in the ground state like a giant matter wave. It must be emphasized that BECs provide an ideal platform (launching pad) to engineer SO coupling in a neutral atomic BEC by dressing two atomic spin states with parts of laser. In fact, in the presence of laser coupling [1], the interaction between the two dressed atomic spin states are modified driving a quantum phase transition from a spatially spin mixed state (laser off) to a phase separated state (stripped phase). The stripped phase which occurs above a critical laser density initiates a phase transition by engineering SO coupling thereby leading to the rich possibility of generating synthetic electric and magnetic fields. Recent investigations have explored the possibility of identifying tunable spin-orbit coupled (SOC) BECs with various trapping potentials and stable domains of condensates have been observed.

The SOC-BECs involving the interplay between the nonlinear two-body interaction with the linear interaction of SOC have been investigated in the last few years. For instance, it has been reported that the new ground-state phase can be created in two-component nonlinear systems such as stripes, phase separation, etc. [2], tricritical points [3], different type of solitons [4–7], 2D solitons with embedded vorticity and vortex lattice [8–11].

The combinations of optical lattice (OL) potential with the interaction of SO coupling was shown to possess interesting phenomena like flattening of Bloch potential [12], atomic Zitterbewegung [13] and new topological phases [14]. Generally, BECs become fragmented on each lattice when it is trapped by deep OL potential. Such a system has been effectively described in the tight-binding approximation by discrete version of the corresponding Gross-Pitaevskii (GP) equation [15–17]. Recent investigations on composite solitons and localized modes are carried out on Rashba type SOC-BEC trapped in deep OL [11,18].

The modulational instability (MI) is a generic phenomenon which occurs in a dynamical system like fluid dynamics, nonlinear optics, plasma physics due to the interplay between nonlinearity and dispersion (or diffraction in spatial domain) resulting in the fragmentation of carrier waves into a train of localized waves [19]. Recently, the onset of MI in scalar [20–22] and vector BECs [23–25] within the framework of continuous NLS type (GP) equations has been investigated. The above scenario could drastically change in discrete multicomponent settings [26–30].

* Corresponding author.

E-mail addresses: ssabari01@gmail.com, sabari.cnls@gcw.ac.in (S. Sabari), taphy1905@gmail.com (R. TamilThiruvalluvar), vittal.cnls@gmail.com (R. Radha).

PRIMER ON SOLVING DIFFERENTIAL EQUATIONS USING MACHINE LEARNING TECHNIQUES

TAMIL ARASAN BAKTHAVATCHALAM^{1,2,a}, SELVAKUMAR MURUGAN^{2,b}, SURIYADEEPAN RAMAMOORTHY^{2,c}, MALAIKANNAN SANKARASUBBU^{2,d}, RADHA RAMASWAMY^{3,e}, VIJAYALAKSHMI SETHURAMAN^{1,f}, BORIS A. MALOMED^{4,g}

¹Department of Physics, Presidency College (Autonomous), University of Madras, Chennai - 600005, India

Email:^a tamilarasanbakthavatchalam@gmail.com

Email:^f vijiphysics.nld@gmail.com

²Saama AI Research Lab, Chennai -600032, India

Email:^b selvakumar.murugan@saama.com

Email:^c suriyadeepan.ramamoorthy@saama.com

Email:^d malaikannan.sankarasubbu@saama.com

³Post-Graduate and Research Department of Physics,

Centre for Nonlinear Science (CeNSc), Government College for Women (Autonomous),

Kumbakonam - 612001, India

Email:^e vittal.cnls@gmail.com

⁴Department of Physical Electronics, School of Electrical Engineering, Faculty of Engineering and Center for Light-Matter Interaction, Tel Aviv University, P.O.B. 39040, Ramat Aviv, Tel Aviv, Israel

Email:^g malomed@tauex.tau.ac.il

Received February 1, 2022

Abstract. Machine Learning (ML) has shown a substantial impact on computational sciences in recent years. The adaptation of ML techniques to deal with various systems in physical sciences has gained ground in addition to the existing numerical methods. In this work, we introduce the readers to machine learning with special reference to Artificial Neural Networks (ANNs) that can solve ordinary differential equations (ODEs) and partial differential equations (PDEs) including those which are subject to specific symmetries. This paper will be helpful for graduate and undergraduate students as an introductory material to early career researchers interested in applying ML techniques to solve problems in computational sciences. In particular, we choose elementary differential equations that describe systems from various fields of science to illustrate the proficiency of ANNs to capture the regularities that underlie such systems in the hope of adding ML techniques to the physicists' toolbelt.

Key words: Machine Learning; Neural Networks; Differential Equations; Deep Learning; Nonlinear Dynamics; Mathematical Modelling; Numerical Methods; Differential Programming.

1. INTRODUCTION

The amount of data accumulated in science over the years is abundant, and the process and algorithms were nothing new. Despite all the above, the rise of comput-

Role of higher-order interactions on the modulational instability of Bose-Einstein condensate trapped in a periodic optical lattice

S Sabari,^{1,*} OT Lekeufack,^{2,3,†} SB Yangoue,^{4,‡} R Tamilthiruvalluvar,^{5,§} and R Radha^{5,¶}

¹*Institute of Theoretical Physics, UNESP – Universidade Estadual Paulista, 01156-970 São Paulo, SP, Brazil.*

²*Pure physics Laboratory: Group of nonlinear physics and complex systems, Department of Physics, Faculty of Science, University of Douala, P.O. Box 24157, Douala, Cameroon*

³*Centre d'Excellence Africain des Technologies de l'Information et de la Communication (CETIC), Université de Yaoundé I, Yaoundé, Cameroun*

⁴*Department of Physics, University of Bamenda, P.O Box 39 Bamenda, Cameroon*

⁵*Centre for Nonlinear Science(CeNSc), Department of Physics, Government College for Women(Autonomous), Kumbakonam – 612001, India*

(Dated: September 23, 2022)

In this paper, we investigate the impact of higher-order interactions on the modulational instability (MI) of Bose-Einstein Condensates (BECs) immersed in an optical lattice potential. We derive the new variational equations for the time evolution of amplitude, phase of modulational perturbation, and effective potential for the system. Through effective potential techniques, we find that high density attractive and repulsive BECs exhibit new character with direct impact over the MI phenomenon. Results of intensive numerical investigations are presented and their convergence with the above semi analytical approach is brought out.

PACS numbers: 03.75.Kk, 03.75.Lm, 05.30.Jp, 67.85.De

I. INTRODUCTION

Solitons in a Bose-Einstein condensate (BEC) trapped in an optical lattice (OL) potential have attracted great attention recently. Generally, BEC in OL is an ideal test bed for condensed matter theory with non-linearity, which gives variety of phenomena such as gap solitons, localization, breathers, diffusion, vortices in lattice, etc... Such a rich dynamics that arises in BECs in OL can be attributed to the phase transition that takes place from superfluid to mott-insulator. In particular, the dispersion dynamics in OL can give rise to stable localized matter wave states in the form of gap solitons. This is represented by stationary solutions of the associated Gross-Pitaevskii (GP) equation with the eigenvalue located at optically induced finite bandgap, in the repulsive condensate also [1].

In the ultracold regime, most of the results of experiments in BECs are reproduced and described by the theoretical model based on the nonlinear mean-field GP equation with two-body interaction [2, 3] and the number density dependent three-body interaction [4, 5]. However, it should be pointed out that the impact of higher order interactions have not yet been considered [2, 6, 7]. The fact that the BEC density increases in various experiments with strong compression of the trap, [8, 9] makes the introduction of higher-order interaction more realistic and inevitable in the description of the dynamics. In very

recent findings, higher-order interactions challenged the three-body interactions over the MI of BECs [10]. Earlier studies bring about stabilization in BECs even with repulsive two body interaction with/without nonlinear management in which the role of higher order interactions has been made more crucial [11, 12].

The inclusion of the shape dependent higher-order interaction at higher density through residual nonlinearity is justified as new phenomena and may bring about changes into the dynamics of BECs especially when immersed in traps like OL potentials. It can be recalled that the recent first real time solitary matter wave pulse was observed experimentally in Rb atoms by the process of modulational instability, where the condensate was kept in periodically varying optical lattice structure (optical waveguide). Thus, in the present work, we examine the effects of higher-order interactions and how they impact the MI [13] of plane wave for a collective interactions including residual nonlinearity of the system consisting of BECs trapped in an OL potential driven by harmonic trap. To produce localized patterns such as solitons, breathers and fundamental vortices which form the basis for energy transport mechanism in nonlinear physical systems, the investigation of the onset of MI is inevitable. Some of the recent investigations towards the exploration of MI parametric domain by means of linear stability analysis from the perspective of higher order interactions in recent years include BECs with zero-nonlinearity in single component [13], residual nonlinearity in two-component condensate [14], Spin-Orbit coupled (SOC) BECs mixture [15], inter-component asymmetry interaction in quantum droplets and SOC-BECs in optical lattice [16, 17].

In the present investigation, through the time-dependent variational approach (TDVA) [10, 20, 21] and

* ssabari01@gmail.com

† lekeufackolivier@gmail.com

‡ sergebruno@yahoo.fr

§ taphy1906@gmail.com

¶ vittal.cnl@gmail.com



Contents lists available at ScienceDirect

Physica A

journal homepage: www.elsevier.com/locate/physa

Quantum correlations and coherence in a moving Unruh–deWitt detector

S. Bhuvaneshwari^a, R. Muthuganesan^b, R. Radha^{a,*}^a Centre for Nonlinear Science (CeNSc), PG & Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India^b Department of Physics, Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Břehová 7, 115 19 Praha 1-Staré Město, Czech Republic

ARTICLE INFO

Article history:

Received 4 May 2022

Received in revised form 24 June 2022

Available online 14 July 2022

Keywords:

Quantum correlation

Coherence

Unruh–deWitt detector

Uncertainty

ABSTRACT

In this paper, we investigate the quantum correlations and coherence of two accelerating Unruh–deWitt detectors coupled to a scalar field in $3 + 1$ Minkowski space–time. We show that the entanglement is completely destroyed in the limit of infinite acceleration while the local quantum uncertainty and l_1 -norm of coherence remain nonzero. In addition, we also highlight the role of Unruh temperature and energy spacing of detectors on quantum correlations for different choices of initial states.

© 2022 Elsevier B.V. All rights reserved.

1. Introduction

Quantum resource theory is an essential tool to understand the nature of physical systems from the perspective of nonlocality and provides an operational interpretation of various quantum effects [1]. Quantum coherence, superposition, entanglement and quantum correlations beyond entanglement are few notable resources which can prove to be advantageous over the classical algorithms. Understanding these resources in physical systems continue to be a formidable and interesting task even today. The quantum information processing is usually studied within the framework of entanglement-versus-separability. The entanglement is considered to be the only early version of nonlocal aspects of quantum systems [2–4] and is demonstrated through the violation of Bell inequality [5]. The seminal work of Werner and Almedia et al. reveals that the entanglement is an incomplete manifestation of nonlocality implying that quantum correlations cannot only be limited to entanglement [6,7] and the separable states can also come in handy in the implementation of some specific quantum tasks. To study this quantum correlation beyond entanglement, Ollivier and W. H. Zurek introduced a measure called quantum discord which captures the nonlocality of separable (unentangled) states [8].

Other than entanglement, many other types of quantum correlations have been discovered in recent years [8–10]. As a result, a number of quantum correlation concepts have emerged with each being motivated by specific applications in quantum information science competing for recognition as the most accurate quantumness metric. For example, the skew-information and quantum Fisher information play an important role in parameter estimation and bring out the limitations on the variance of the observable. As stated, each quantum correlation captures different quantumness in quantum systems due to their distinct type of measurement techniques. Among them, skew information based quantum correlation measures such as local quantum uncertainty (LQU) and uncertainty-induced nonlocality (UIN) are widely used

* Corresponding author.

E-mail addresses: rajendramuthu@gmail.com (R. Muthuganesan), vittal.cnl@gmail.com (R. Radha).



PAPER

Elusive exotic structures and their collisional dynamics in (2+1)-dimensional Boiti-Leon-Pempinelli equation

RECEIVED
1 July 2022REVISED
6 October 2022ACCEPTED FOR PUBLICATION
11 November 2022PUBLISHED
23 November 2022R Radha^{1,*}, Sudhir Singh², C Senthil Kumar³ and Senyue Lou⁴¹ Centre for Nonlinear Science (CeNSc), Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam-612001, India² Department of Mathematics, National Institute of Technology, Tiruchirappalli 620015, India³ Department of Physics, Vinayaka Mission's Kirupananda Variyar Engineering College, A Constituent College of Vinayaka Mission's Research Foundation (Deemed to be University) NH-47, Sankari Main Road, Periyaseeragapadi, Salem-636308, India⁴ School of Physical Science and Technology, Ningbo University, Ningbo, 315211, People's Republic of China

* Author to whom any correspondence should be addressed.

E-mail: vittal.cnl@gmail.com**Keywords:** localized solutions, truncated Painlevé expansion approach, singular manifold, dromions, lumps, rogue waves, breathers**Abstract**

In this paper, we investigate the (2+1) dimensional Boiti-Leon-Pempinelli (BLP) equation employing truncated Painlevé expansion approach and extract a plethora of localized nonlinear waves, including multi-dromions, multi-lumps, multi-rogue waves, generalized-breathers etc. The dromions are characterized as bright, dark and mixed (bright-dark) based on their intensity. The collisional dynamics of dromions shows that they change their shape or form upon interaction in addition to undergoing a phase change. The lump solutions of orders one and two are also extracted through appropriate test functions and observed to be non-interacting in nature. Also, the first-order and second-order rogue waves are also obtained through rational polynomials and shown to be unstable. The generalized breathers are obtained by utilizing the three-wave test function. The highlights of our investigation is that one encounters a strange coherent structure called 'dromion filter' which contains a dynamic and a stationary dromion. In addition, we are also able to unearth a 'coexistent dromion-line soliton'.

1. Introduction

The advent of dromions [1–5] had a huge impact (bearing) in the quest for subsequent identification of other localized solutions like rogue waves [6–9], interaction waves [10–13] and lumps [14–19]. While dromions which originate at the cross point of the intersection of two nonparallel ghost solitons decay exponentially in all directions and are being driven by boundaries, Lumps are essentially algebraically decaying solutions which do not interact with each other. The above class of localized solutions find applications in a variety of contexts. While rogue waves have been identified in hydrodynamics [20], Bose–Einstein Condensates [21], Plasma Physics [22], etc., dromions appear in the domain of meta materials as plasmon dromions [23], in the study of flexural-gravity waves [24], electron acoustic waves in space plasmas [25], in ultrafast lasers [26], etc.

The richness in the structure of localized solution in the (2+1) dimensional nonlinear partial differential equations (pdes) can be attributed to the existence of lower dimensional arbitrary functions of space and time and it is believed that manoeuvring these lower dimensional arbitrary functions of space and time can give rise to newer and unknown localized structures. It is this peculiar perspective of the (2+1) dimension nonlinear pdes which makes their investigation interesting even today. In this direction, we wish to analyse the localized solutions admitted by Boiti-Leon-Pempinelli (BLP) equation [27].

$$v_t - v_{xx} - 2uv_x = 0, \quad (1)$$

$$u_{yt} - (u^2 - u_x)_{xy} - 2v_{xxx} = 0. \quad (2)$$

Letter

Signatures of intrinsic decoherence and weak measurement on quantum correlations

S Bhuvaneshwari^{1,*}, R Muthuganesan² and R Radha^{1,*}¹ Centre for Nonlinear Science (CeNSc), PG and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India² Centre for Nonlinear Science and Engineering, School of Electrical and Electronics Engineering, SASTRA Deemed University, Thanjavur, Tamil Nadu, IndiaE-mail: bhuvanajkm85@gmail.com and vittal.cnl@gmail.com

Received 19 June 2021

Accepted for publication 16 November 2021

Published 2 December 2021

**Abstract**

In this article, we consider a pair of spin-1/2 particles with squeezing coupling serving as the physical carrier of quantum information. We then examine the dynamics of quantum correlation quantified by the entanglement and measurement-induced nonlocality (MIN) under the intrinsic decoherence. The impact of intrinsic decoherence on the dynamical behaviors of quantum correlations is investigated. We show that the MIN quantities are more robust, while intrinsic decoherence cause sudden death in entanglement. Besides, we highlight the role of spin squeezing coupling and external magnetic field on quantum correlation measures. Finally, we investigate the impact of weak measurement on MIN.

Keywords: quantum correlation, intrinsic decoherence, spin squeezing model, nonlocality

(Some figures may appear in colour only in the online journal)

1. Introduction

Quantum information science has turned out to be a rapidly growing multidisciplinary research area in recent times exploiting the properties of collective quantum phenomena such as quantum correlation and entanglement. These collective phenomena make a fundamental difference between classical and quantum systems and are believed to be a vital resource for various quantum technology applications [1–8]. Entanglement refers to a ‘spooky action at a distance’ and considered as a certified manifestation of nonlocal attributes of a quantum system [9–11]. The presence of entanglement in the quantum system is witnessed through the violation of Bell inequality [12]. For pure states, it is believed that

the entanglement is the complete manifestation of a quantum system’s nonlocality. On the other hand, the nonlocal behaviors of mixed states are still not understood. Seminal work of Bell [13] and the measures of quantum correlations beyond entanglement [14–17] reveal that the entanglement does not capture the entire spectrum of nonlocality of a quantum state due to the mixedness [18]. In the light of the above observation, it is concluded that the entanglement is an incomplete manifestation of nonlocality and to capture complete picture of nonlocality of quantum systems, we are in dire need of better indicators of quantum correlations.

A decade ago, a new variant and more reliable quantum correlation measure was introduced by Luo and Fu [17]. This captures bipartite correlation or nonlocal effects due to locally invariant eigenprojective measurements on any one of the subsystems. The local measurements on a subsystem can influence globally and this notion of nonlocality is more

* Authors to whom any correspondence should be addressed.

CHANGES OBSERVED ON STRESS AFTER TWELVE WEEKS OF PRANAYAMA PRACTICES AMONG COLLEGE WOMEN

Ms. P. SANMUGAPRIYA,

Research Scholar in Yoga, Centre for Yoga Studies, Annamalai University.

Dr. R. VIDHYASREE,

Assistant Professor, Department of Physical Education, Annamalai University.

ABSTRACT

The purpose of the study was designed to examine the effect of pranayama practices on stress of college women students. For the purpose of the study, thirty women students from various Departments at Annamalai University were selected as subjects. They were divided randomly into two equal groups. Each group consisted of the fifteen subjects. Group I underwent pranayama practices for five days per week for twelve weeks. Group II acted as control who did not undergo any special training programme apart from their regular programme. The following variable namely stress was selected as criterion variable. All the subjects of two groups were tested on selected dependent variable at prior to and immediately after the training programme. The analysis of covariance [ANCOVA] was used to analyze the significant difference, if any among the groups. The .05 level of confidence was fixed as the level of significance to test the 'F' ratio obtained by the analysis of covariance, which was considered as an appropriate. The results of the study showed that there was a significant difference among pranayama practices group and control group on stress. And also the results of the study showed that there was a significant reduction on stress due to twelve weeks of pranayama practices.

KEYWORDS:

PRANAYAMA PRACTICES, STRESS, ANCOVA, COLLEGE WOMEN



EFFECT OF SIMPLIFIED KUNDALINI YOGA WITH AND WITHOUT SUPER BRAIN YOGA ON SELF CONFIDENCE AMONG ADOLESCENT GIRLS WITH LEARNING DISABILITY FACTORS

B.Yamuna, Ph.D Scholar- Yoga, Centre for Yoga Studies, Annamalai University
Dr. R.Vidhyasree, Assistant Professor, Department of Physical Education, Government College for women, Kumbakonam

ABSTRACT

The purpose of the study was to effect of the effect of simplified kundalini yoga with and without super brain yoga on self confidence among adolescent girls with learning disability factors. The study was conducted on 90 adolescent girls with learning disability factors. Totally three group's namely experimental group I, II and control group consisting of 30 adolescent girls with learning disability factors in each group who underwent sixteen weeks of yoga practices whereas the control group did not undergo any type of training. The self confidence was measured before and after the experimentation using the standardized questionnaire to measure the behaviour parameter and analyzed by Analysis of Covariance (ANCOVA) and it was concluded that the simplified kundalini yoga with super brain yoga had significant ($P < 0.05$) effect on the self confidence.

Key words: simplified kundalini yoga, super brain yoga, self confidence, behaviour parameter, adolescent girls, and learning disability.

INTRODUCTION:

The term Yoga suggests that 'Yuj' in Indo-Aryan which implies union. Yoga may be a union of body, mind and soul. in keeping with Vethathiri Maharishi (1992) 'Yoga may be a systematic psychic apply to boost awareness, to develop will-power and to comprehend the self'. Yoga is generally classified into bhakthi yoga, karma yoga, yoga and jnana yoga. These branches of yoga is the bottom for the origin of various yoga forms like kriya yoga, laya yoga, kundalini yoga, sahaja yoga, raja yoga, mantra yoga, etc. Simplified kundalini yoga (SKY) may be a combination of raja yoga and kundalini yoga initiated by Yogiraj Vethathiri Maharishi. Simplified kundalini yoga is AN integrated system consisting of four major parts like kayakalpa exercises, simplified physical exercises, meditation and self-examination. Kayakalpa exercises and simplified physical exercises facilitate to boost the standard of life-force and promote sensible physical health. Meditation apply helps in elevating the mind of the professional person and eventually, self-examination courses facilitate a private to research his/her emotions, that ultimately results in religious development through temperament transformation (Krishnan's, 2006).

In day to day life, solely bound elements of the body ar functioning in keeping with the character of labor. All the elements of the body ought to perform properly to keep up the body match. it's necessary that a simplified exercise may be a should to create the elements of the body perform to a minimum level. There ar several types of physical exercises within the world. Vethathiri Maharishi has chosen the simplest ones from round the world, analyzed them together with his medical expertise, practiced them for an extended amount, and enjoyed their edges. Then he gave it to United States as a simplified physical exertion. This exercise doesn't strain the body, consumes less time, and edges ar higher.

STATEMENT OF THE PROBLEM

The present study was designed to effect of simplified kundalini yoga with and without super brain yoga on selected self confidence among adolescent girls with learning disability factors.



Anti-viral effectuality of plant polyphenols against mutated dengue protein NS2B47-NS3: A computational exploration

J. Maria Jancy Rani^a, K. Kalaimathi^b, S. Vijayakumar^c, G. Varatharaju^d, K. Karthikeyan^e,
G. Thiagarajan^e, K. Bhavani^f, P. Manogar^g, S. Prabhu^{h,*}

^a Department of Chemistry, Annai Vallankanni Arts and Science College, Bishop Sundaram Campus (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu-613 007, India

^b Department of Chemistry, Government College of Engineering, Sengipatti, Thanjavur, Tamil Nadu-613 402, India

^c PG and Research Department of Botany, AVVM Sri Pushpam College (Autonomous) Poondi, (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu- 613 503, India

^d Department of Botany, Sri Kaliswari College (Autonomous), Sivakasi, Tamil Nadu-626 123, India

^e Department of Botany, Government Arts College (Autonomous), Kumbakonam, (Affiliated to Bharathidasan University), Tamil Nadu- 612 002, India

^f Department of Zoology Government College for Women (Autonomous), Kumbakonam (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu- 612 001, India

^g Department of Botany, Nehru Memorial College (Autonomous), Puthanampatti (Affiliated to Bharathidasan University), Tamil Nadu-621 007, India

^h Department of Botany, Annai Vallankanni Arts and Science College, Bishop Sundaram Campus (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu-613 007, India

ARTICLE INFO

Keywords:

Anti-dengue
Phytopolyphenols
Inhibitor validation
Interaction analysis
Pharmacokinetics parameters

ABSTRACT

Dengue fever is a tropical endemic disease that is transmitted by the female *Aedes aegypti* mosquitoes around the world. In dengue patients, the severe vascular leak, bleeding and organ failure underlines the severity of this disease. It affects 100 million people each year and half a million cases of dengue hemorrhagic fever are recorded worldwide. In addition, the combined infections of COVID-19 and dengue fever cause serious health problems for those infected. Because polyphenolic compounds have the ability to regulate and restore the immune system, the intervention of biological tools is required to find an accurate polyphenolic compound from plants to combat these viral diseases. In this computational study, we used a range of software to explore the anti-dengue potential of the selected polyphenols, while the database was also used to explore diseased enzymes and drug targets in the dengue pathways of *Homo sapiens*. All examined polyphenols showed excellent docking values after molecular docking and also showed remarkable hydrogen bond interactions between the ligand and the dengue protein. Among the polyphenols tested, (R)-(+)-rosmarinic acid has a docking score of -8451 with an energy value of -59,860. After that, curcumin has the second best docking score of -8.221, followed by D-(+)-catechin (-7.042) and epicatechin (-7.069). In addition, molecular dynamic simulations demonstrate that the complex of (R)-(+)-rosmarinic acid and dengue protein has a solid understanding of the precise interactions. The current study suggests that this is the first report of such compounds against the dengue virus, despite extensive research on these substances in previous research. Moreover, we expect these polyphenols to show promising effects when used in antiviral research such as *in-vitro* and *in-vivo* studies. In addition, toxicological studies such as adsorption, distribution, metabolism, and excretion are required before further exploration of the polyphenols, as this will help to understand the biological processes of these potential polyphenols.

1. Introduction

Dengue fever seems to be extremely prevalent around the world in the past few decades. It is mainly transmitted to humans by the infected female *Aedes aegypti* mosquitoes. This virus can be transmitted up to two

days before symptoms of dengue appear (Duong et al., 2015) and it can be active for up to two days after the disease has resolved (Nguyen et al., 2013). The large number of dengue cases goes undiagnosed because of the mild symptoms and are treated by infected people themselves. As a result, it is found that the actual number of dengue cases is not

* Corresponding author.

E-mail address: prabhusbotany@gmail.com (S. Prabhu).

<https://doi.org/10.1016/j.genrep.2022.101546>

Received 21 October 2021; Received in revised form 2 January 2022; Accepted 3 February 2022

Available online 8 February 2022

2452-0144/© 2022 Elsevier Inc. All rights reserved.



Anti-dengue Potential of Mangiferin: Intricate Network of Dengue to Human Genes

K. Kalaimathi¹ · J. Maria Jancy Rani² · S. Vijayakumar³ · N. Prakash² · K. Karthikeyan⁴ · G. Thiyagarajan⁴ · K. Bhavani⁵ · S. Prabhu⁶ · G. Varatharaju⁷

Received: 28 December 2021 / Accepted: 14 April 2022 / Published online: 7 May 2022
© The Author(s) under exclusive licence to Sociedade Brasileira de Farmacognosia 2022

Abstract

Dengue fever has become one of the deadliest infectious diseases and requires the development of effective antiviral therapies. It is caused by members of the Flaviviridae family, which also cause various infections in humans, including dengue fever, tick-borne encephalitis, West Nile fever, and yellow fever. In addition, since 2019, dengue-endemic regions have been grappling with the public health and socio-economic impact of the ongoing coronavirus disease 19. Co-infections of coronavirus and dengue fever cause serious health complications for people who also have difficulty managing them. To identify the potentials of mangiferin, a molecular docking with various dengue virus proteins was performed. In addition, to understand the gene interactions between human and dengue genes, Cytoscape was used in this research. The Kyoto Encyclopedia of Genes and Genomes software was used to find the paths of Flaviviridae. The Kyoto Encyclopedia of Genes and Genomes and the Reactome Pathway Library were used to understand the biochemical processes involved. The present results show that mangiferin shows efficient docking scores and that it has good binding affinities with all docked proteins. The exact biological functions of type I interferon, such as interferon- α and interferon- β , were also shown in detail through the enrichment analysis of the signaling pathway. According to the docking results, it was concluded that mangiferin could be an effective drug against the complications of dengue virus 1, dengue virus 3, and non-structural protein 5. In addition, computational biological studies lead to the discovery of a new antiviral bioactive molecule and also to a deeper understanding of viral replication in the human body. Ultimately, the current research will be an important resource for those looking to use mangiferin as an anti-dengue drug.

Keywords Dengue proteins · Protein-protein interaction · Pharmacokinetics · Drug probabilities

Introduction

Dengue fever is a serious viral disease that is ubiquitous in tropical and subtropical regions (Guzman and Harris 2015). According to the World Health Organization, 100 million cases of dengue fever occur every year. In 500,000 of these cases,

hospitalization is required and in 25,000 cases, conditions worsen and lead to death. According to a recent report, there are 390 million dengue infections worldwide each year, more than three times the estimates of the World Health Organization (Qamar et al. 2019). This mosquito-borne virus disease is endemic in over 100 countries and triggered by the dengue virus transmitted

✉ S. Prabhu
prabhu@avasctnj.edu.in

¹ Department of Chemistry, Government College of Engineering, Sengipatti, Thanjavur, Tamil Nadu 613402, India

² Department of Chemistry, Annai Vailankanni Arts and Science College, Bishop Sundaram Campus (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu 613007, India

³ PG and Research Department of Botany, AVVM Sri Pushpam College (Autonomous), Poondi (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu 613503, India

⁴ Department of Botany, Government Arts College (Affiliated to Bharathidasan University), Kumbakonam, Tamil Nadu 612002, India

⁵ Department of Zoology, Government Arts College for Women (Autonomous), Affiliated to Bharathidasan University, Kumbakonam, Thanjavur, Tamil Nadu 612001, India

⁶ Department of Botany, Annai Vailankanni Arts and Science College, Bishop Sundaram Campus (Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu 613007, India

⁷ Department of Botany, Sri Kaliswari College (Autonomous), Sivakasi, Tamil Nadu 626123, India



The impact of public spending in education and health on human development in India

Martin Ruzima^a and P. Veerachamy^b

^aResearch Unit, Institute of Policy Analysis and Research – Rwanda (IPAR-Rwanda), Kigali, Rwanda;

^bDepartment of economics, Government college for Women(A), Kumbakonam, India

ABSTRACT

This study examined the effect of public spending on education and health sectors on human development in India. Data for the period 1990–2018, obtained from UNDP and World Bank databases, were used. The ARDL model technique was used for data analysis. In the long run, the findings showed that public expenditure on health and education has positive and negative significant effects, respectively, on human development in India. In the short run, results indicated that the public expenditure on health and education has positive and negative respectively insignificant impacts on human development in India. In addition, GDP per capita has been found to have positive but insignificant impact on human development for both long run and short run. The findings of this article suggest that government of India should rise funding for health and education with proper targeting.

KEYWORDS

Public spending; human development; education; health; ARDL; GDP

1. Introduction

It is widely agreed among economists that the human development contributes to sustainable economic development. Human development emerged *via* the works of Sen and Mahbub in 1990. They argued that people are the wealth of the nation (Malik 2014). In line with this, the United Nation Development Programme (UNDP) developed Human Development Index (HDI) to track the progress and well-being of people, and every year UNDP releases a report with different topics relating to human welfare. The HDI is calculated based on literacy, life expectancy and income levels in the country (UNDP 1990). Human development is improved by investing in human capital (Schultz 1961; Becker 1993). Investment in human capital is done through spending on health and education at any level.

Public spending or investment in health and education is known as ‘social spending’ or simply investment in human capital. Thus, increased expenditure in these two sectors leads to a rise in human development and consequently economic growth (Gupta et al. 1998). An increase in government spending on health care reduces the mortality rate, which results in higher life expectancy (Kim and Lane 2013; Jaba,

Stability window of trapless polariton Bose-Einstein condensates

S. Sabari, R. Kishor Kumar, R. Radha, and P. Muruganandam

Phys. Rev. B **105**, 224315 – Published 27 June 2022

Article

ABSTRACT

We theoretically explore the possibility of stabilizing the trapless polariton Bose-Einstein condensates (pBECs). Exploiting the variational method, we solve the associated nonlinear, complex Gross-Pitaevskii equation and derive the equation of motion for the amplitude and width of the condensate. These variational results described by ordinary differential equations are rewritten to perform a linear stability analysis to generate a stability window in the repulsive domain. A set of coupled nonlinear ordinary differential equations obtained through the variational approach are then solved by numerical simulations through the fourth-order Runge-Kutta method, which are further supported by the split-step Crank-Nicholson method, thereby setting the platform for stable pBECs. In particular, we generate a window containing system parameters in the $g_1-\gamma_{\text{eff}}$ space within which the system can admit stable condensates. The highlight of the results is that one observes beating effects in the real time evolution of the condensates with attractive interactions much similar to multicomponent BECs, and their periodicity can be varied by manipulating linear and nonlinear loss/gain terms. For repulsive condensates, one notices the stretching of the density.

- Received 6 June 2021
- Accepted 9 June 2022

DOI:<https://doi.org/10.1103/PhysRevB.105.224315>

©2022 American Physical Society

Physics Subject Headings (PhySH)

1. [Research Areas](#)
[Exciton polariton](#)
1. [Physical Systems](#)
[Bose-Einstein condensates](#)[Polariton condensate](#)

Nonlinear Dynamics

AUTHORS & AFFILIATIONS

[S. Sabari](#)^{1,2}, [R. Kishor Kumar](#)³, [R. Radha](#)², and [P. Muruganandam](#)⁴

- [Institute of Theoretical Physics, UNESP – Universidade Estadual Paulista, 01156-970 São Paulo, São Paulo, Brazil](#)

- ²Centre for Nonlinear Science (CeNSc), Department of Physics, Government College for Women (A), Kumbakonam 612001, India
- ³Department of Physics, Centre for Quantum Science, and Dodd-Walls Centre for Photonic and Quantum Technologies, University of Otago, Dunedin 9054, New Zealand
- ⁴Department of Physics, Bharathidasan University, Tiruchirappalli 620024, India

GOVERNMENT COLLEGE FOR WOMEN (AUTONOMOUS)

KUMBAKONAM – 612 001

Affiliated to Bharathidasan University

DST - CURIE Sponsored Institution

IV Cycle of Accreditation



☎ 0435 – 2401391


✉ principal@gwk.ac.in



CRITERION III – RESEARCH, INNOVATIONS AND EXTENSION

FRONT PAGE OF PUBLISHED PAPERS

2020-2021



The aromatic ginger *Kaempferia galanga* L. (Zingiberaceae) essential oil and its main compounds are effective larvicidal agents against *Aedes vittatus* and *Anopheles maculatus* without toxicity on the non-target aquatic fauna

Mohamad S. AlSalhi^{a,b,**}, Kuppusamy Elumalai^c, Sandhanasamy Devanesan^{a,b},
Marimuthu Govindarajan^{d,e,*}, Kaliyamoorthy Krishnappa^f, Filippo Maggi^g

^a Research Chair in Laser Diagnosis of Cancers, Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh 11451, Saudi Arabia

^b Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh 11451, Saudi Arabia

^c Department of Advanced Zoology & Biotechnology, Government Arts College for Men (Autonomous), Chennai, 600035, Tamil Nadu, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

^f Department of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai, 609305, India

^g School of Pharmacy, University of Camerino, Via Sant'Agostino 1, 62032 Camerino, Italy

ARTICLE INFO

Keywords:

Aedes vittatus
Anopheles maculatus
ecofriendly approach
essential oil
Kaempferia galanga
novel insecticide

ABSTRACT

Kaempferia galanga L. (Zingiberaceae), also known as aromatic ginger, is used in the food and cosmetics, pharmaceuticals and maquillage materials, as well as in the traditional Indian medicine. The essential oil (EO) obtained from its rhizome by hydrodistillation is of economic interest and contains ethyl *p*-methoxycinnamate as the main component, which is used as an anticancer, antimicrobial, skin protector and whitening and nematocidal agent. To our knowledge, the mosquitocidal properties of the aromatic ginger EO have hitherto been poorly investigated. Therefore, in the present work, we tested this EO and its major chemical constituents (MCCs), namely ethyl *p*-methoxycinnamate, *trans*-ethyl cinnamate and *trans*-cinnamaldehyde for their larval toxicity on *Aedes vittatus* and *Anopheles maculatus*. The EO chemical composition, studied by GC-MS analysis, was made up of 39 compounds, with ethyl *p*-methoxycinnamate (30.6%), *trans*-ethyl cinnamate (26.8%) and *trans*-cinnamaldehyde (11.5%) as the major constituents. The EO and its MCCs showed LC₅₀ values of 39.22, 10.36, 28.26 and 36.35 µg/ml on *Ae. vittatus*, and of 41.36, 12.56, 30.23 and 38.47 µg/ml on *An. maculatus* third instar larvae, respectively. The ecofriendliness of EO and MCCs on the aquatic fauna was demonstrated as they were safe to *Anisops bouvieri*, *Acilius sulcatus* and *Gambusia affinis*. Our findings showed that the *K. galanga* rhizome EO and its MCCs are promising larvicidal agents without affecting non-target organisms (NTOs). They also give new insights into the potential economic exploitation of this EO and its MCCs in the fabrication of effective and eco-friendly mosquitocidal formulations.

1. Introduction

The human communities are extremely suffered by the nuisance of growing Dipteran insect populations in both agricultural and medical fields (Pavela et al., 2020a, b,c). Mosquitoes (Diptera: Culicidae) are known to transmit several dreadful human diseases, creating negative

impacts on human health and, therefore, they have been declared as a 'public enemy' (Pavela et al., 2019a; Benelli et al., 2020a, b). Bites of vector mosquitoes not only transmit many diseases but also cause irritation and allergic response to the skin (Cheng et al., 2003; Awosolu et al., 2018; Thongwat et al., 2018). Worldwide, mosquitoes are responsible for causing millions of deaths every year (WHO, 2011),

* Corresponding author at: Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India.

** Corresponding author at: Research Chair in Laser Diagnosis of Cancers, Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh 11451, Saudi Arabia.


E-mail addresses: malsalhi@ksu.edu.sa (M.S. AlSalhi), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.indcrop.2020.113012>

Received 12 July 2020; Received in revised form 19 August 2020; Accepted 11 October 2020

Available online 20 October 2020

0926-6690/© 2020 Elsevier B.V. All rights reserved.



Synthesis and characterization of ZnO and Ca-ZnO nanoparticles for potential antibacterial activity and plant micronutrients

Saraswathi Umavathi¹, Mohamad S. AlSalhi², Sandhanasamy Devanesan², Sivasamy Kadhiraivan³, Kasi Gopinath⁴, Marimuthu Govindarajan^{5,6,*}

¹ Department of Botany, Adhiyaman Arts and Science College for Women, Uthangarai, Krishnagiri 635 207, Tamil Nadu, India

² Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh, 11451, Kingdom of Saudi Arabia

³ Department of Chemistry, Swami Dayananda College of Arts and Science, Manjakkudi, Tiruvarur-612 610, Tamil Nadu, India.

⁴ School of Materials and Energy, Southwest University, Chongqing 400715, PR China

⁵ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India

⁶ Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Nanotechnology
Precipitation
Antibacterial activity
Electron microscopy
In vitro regeneration

ABSTRACT

Zinc oxide nanoparticles (ZnO NPs) and calcium-doped zinc oxide (Ca-ZnO) NPs were produced via coprecipitation assay. Physicochemical properties of synthesized NPs have been studied by X-ray powder diffraction (XRD), Ultraviolet-visible (UV-vis), Photoluminescence (PL), scanning electron microscopy (SEM), Energy Dispersive X-ray (EDAX) with fluorescent elemental mapping, High-Resolution Transmission Electron Microscopy (HR-TEM) analyses. UV-vis analysis shows the presence of Ca doping in the Zn lattice, which reduces the bandgap. The XRD patterns of the NPs has a hexagonal wurtzite structure. SEM and HR-TEM images exhibited spherical morphology. The PL spectral measurements indicated six bands embedded due to zinc and oxygen vacancies. The crystallization temperature of the as-prepared samples was measured by a thermogravimetric (TG) analysis. Moreover, the antibacterial activity of the NPs was tested against *Staphylococcus aureus* and *Escherichia coli* bacteria using a disk diffusion method. The antibacterial activity of ZnO NPs was advanced compare to Ca-ZnO NPs. By contrast, Ca-ZnO NPs reduced antibacterial efficacy. Additionally, *Sphaeranthus indicus* shootlets were tested in different concentrations (0-5 mg/L) of ZnO NPs and Ca-ZnO NPs in Murashige and Skoog media. An impact of Ca-ZnO NPs significantly induced the development of shootlets, leaves and node initiation on *S. indicus*, compared to pure ZnO NPs. Plant development was increased by increasing the concentrations of NPs. Therefore, these biocompatible Ca-ZnO NPs can be used for target-drug delivery in biomedicine and as a micro-nutrient for *in vitro* plant growth and development.

1. Introduction

In recent years, Zinc oxide nanoparticles (ZnO NPs) have excellent physicochemical properties such as (i) II-VI semiconductors, (ii) direct bandgap value at 3.36 eV, (iii) n-manner semiconductors, (iv) binding energy of 60 meV, (v) UV-light absorbance, (vi) excellent stability, and (vii) hexagonal wurtzite structure [1–3]. ZnO NPs have potential applications in optoelectronics, photonics [4–6], dye-sensitized solar cells, sensors, photocatalysis, and plant tissue culture, have antimicrobial, anticancer, anti-inflammatory, wound healing properties, and have applications in food packing [7–15] and as next-generation inorganic light-emitting diodes [16]. The optoelectronic characteristics of ZnO

NPs fundamentally depend on changes in the material's disorder level, which is affected by electron-phonon interactions [17]. In early literature, the photoluminescence defect level was shown to be reduced in Cu doped ZnO NPs (Cu-ZnO NPs); thus, these NPs could be useful in optical devices [18,19].

Zinc (Zn) is an important micronutrient for plant development, playing a pivotal role in physiological functions such as nitrogen metabolism, energy transfer, protein synthesis, plant growth hormone production, and inter-node elongation [20]. Consequently, zinc deficiency can alter plant growth processes such as starch accumulation, small leaf formation, flowering and fruiting, reduced seed yield, and increased oxidative stress [21–24]. Calcium (Ca) is also an essential

* Corresponding author.

E-mail address: drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.surfin.2020.100796>

Received 10 August 2020; Received in revised form 28 October 2020; Accepted 31 October 2020

Available online 7 November 2020

2468-0230/© 2020 Elsevier B.V. All rights reserved.



Photocatalytic, antiproliferative and antimicrobial properties of copper nanoparticles synthesized using *Manilkara zapota* leaf extract: A photodynamic approach

Rose Mary Kiriyanthan^a, S. Aruna Sharmili^{a,*}, R. Balaji^b, S. Jayashree^a, Shahid Mahboob^c, Khalid A. Al-Ghanim^c, Fahad Al-Misned^c, Zubair Ahmed^c, Marimuthu Govindarajan^{d,e}, Baskaralingam Vaseeharan^{f,*}

^a Department of Biotechnology, Stella Maris College (Autonomous), Chennai, 600086, Tamil Nadu, India

^b Centre for Advance Study in Botany, University of Madras, Guindy Campus, Chennai, 600025, Tamil Nadu, India

^c Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women (A), Kumbakonam, 612 001, Tamil Nadu, India

^f Department of Animal Health and Management, Alagappa University, Science Block, 6th Floor, Burma Colony, Karaikudi, 630 004, Tamil Nadu, India

ARTICLE INFO

Keywords:

Copper nanoparticles
Manilkara zapota
 Characterization
 Antimicrobial assay
 Antiproliferative assay

ABSTRACT

Copper nanoparticles were synthesized using *Manilkara zapota* leaf extract. The synthesis of the nanoparticle was primarily visualized when the colour of the reaction mixture turned into reddish-brown. Biosynthesized nanoparticles were characterized by UV-vis, FT-IR, XRD, SEM and EDX. The UV spectra showed maximum absorption at 584 nm. FT-IR studies showed stretching frequency at 592.76 cm^{-1} , which is the fingerprint region for Cu-O bond. The crystallinity of the synthesized copper nanoparticles (Mz-Cu NPs) was revealed through XRD analysis. The synthesized Mz-Cu NPs were spherical with an average size of 18.9–42.5 nm and it was shown by SEM analysis. EDX analysis displayed that the nano sample contains 58 % of copper. The antimicrobial property of the synthesized nanoparticles was evaluated against fungal plant pathogens *Rhizoctonia solani* (MTCC 12232), *Sclerotium oryzae* (MTCC 12230) and bacterial species, namely *Bacillus subtilis* (ATCC 23857), *Escherichia coli* (ATCC 25922), *Staphylococcus aureus* (ATCC 25923), *Vibrio harveyi* (ATCC 35084), *Vibrio parahaemolyticus* (ATCC 33845). In *in-vitro* haemolytic assay, the particle showed 5.73, 3.34, 0.5 % hemolysis at 100, 50, 25 $\mu\text{g}/\text{mL}$ concentration respectively. In the antiproliferative assay, the IC_{50} values of MCF7 and Vero cells were found to be 53.89 and 883.69 $\mu\text{g}/\mu\text{l}$. The particle degraded Methyl violet, Malachite green and Coomassie brilliant blue by 92.2, 94.9 and 78.8 %, within 50, 40 and 60 min, respectively, through its photocatalytic activity.

1. Introduction

Nanotechnology is manipulating materials measuring less than 100 nm in size to generate various properties seldom found in their corresponding bulk materials [1,2]; a few such properties are catalysis, imaging, energy conversion and conservation [3]. Nanobiotechnology is exploiting these properties in biotechnology, physics, chemistry, and material science [4]. Metal nanoparticles are currently the hub of research due to wide-ranging applications. They can be manufactured using biological, chemical, and physical methods. Amongst, physical and chemical methods are cost-effective and toxic to the environment

due to reagent, toxicity, and tedious processes. Contrary wise, the biological method is an eco-friendly approach, where organic compounds like plant/microbial extract, animal products, and other biomolecules are used as reducing, capping, and stabilizing agent. These compounds are mostly water-soluble, non-toxic, and biodegradable; hence the biological method is usually called green synthesis. Plant extracts are generally preferred over other organic compounds as it is more efficient, economical, easy to prepare and store [5–8].

Copper has been the metal of choice due to its low cost of preparation and also because of its incomparable physicochemical properties which are analogous to that of noble metals such as gold and silver [9]. They

* Corresponding authors.

E-mail addresses: arunasharmili@stellamariscollege.edu.in (S.A. Sharmili), vaseeharan@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.pdpdt.2020.102058>

Received 19 September 2020; Received in revised form 27 September 2020; Accepted 2 October 2020

Available online 13 October 2020

1572-1000/© 2020 Elsevier B.V. All rights reserved.



Biogenic synthesis of aromatic cardamom-wrapped zinc oxide nanoparticles and their potential antibacterial and mosquito larvicidal activity: An effective eco-friendly approach

Viswanathan Vinotha^a, Mariappan Yazhiniprabha^a, Daniel Selva Raj^b, Shahid Mahboob^c, Khalid A. Al-Ghanim^c, Fahad Al-Misned^c, Marimuthu Govindarajan^{a,e}, Baskaralingam Vaseeharan^{a,*}

^a Nanobioscience and Nanopharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Zoology, Scott Christian College, Nagercoil 629 003, Tamil Nadu, India

^c Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Editor: Zhang Xiwang

Keywords:

Elettaria cardamomum
Zinc oxide nanoparticles
Antibiofilm
Aedes aegypti
Culex tritaeniorhynchus
Enzymes reduction

ABSTRACT

In the current study, *Elettaria cardamomum* (*Ec*) seed extract-assisted zinc oxide nanoparticles (ZnO NPs) were synthesized by a secure co-precipitation method and characterized through UV-vis, XRD, FTIR, SEM and TEM. The surface plasmon resonance of *Ec*-ZnO NPs was distinguished at 373 nm. The XRD pattern authenticates the crystalline nature and matches with JCPDS file No. 36-1451. The result acquired from the FTIR spectrum conveyed the potential biomolecules that take place in the *Ec*-ZnO NPs by sharp express peaks at 3472.30, 1416.81, 1573.95, 1557.02, and 848.25 cm^{-1} . SEM and TEM images make known the homogenous spherical form and average size (18.72 nm) of *Ec*-ZnO NPs, respectively. The *Ec*-ZnO NPs have better antioxidant activity at 100 $\mu\text{g/ml}$ concentration than *Ec* seed extract. Additionally, the *Ec*-ZnO NPs were encompassed with enhanced antibiofilm activity at 100 $\mu\text{g/ml}$ concentration against four different bacteria *Enterococcus faecalis*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*, and *Proteus vulgaris*. Among them, Gram-negative bacteria were more susceptible to *Ec*-ZnO NPs. Moreover, when compared to *Ec* seed extract, the *Ec*-ZnO NPs was highly potential agent against *Aedes aegypti* ($\text{LC}_{50} = 13.27 \mu\text{g/ml}$, $\text{LC}_{90} = 25.36 \mu\text{g/ml}$) and *Culex tritaeniorhynchus* ($\text{LC}_{50} = 15.09 \mu\text{g/ml}$, $\text{LC}_{90} = 29.70 \mu\text{g/ml}$). Compare to *Cx. tritaeniorhynchus*, *Ae. aegypti* was more susceptible to *Ec*-ZnO NPs. Thus, the *Ec*-ZnO NPs can probably utilize to reduce the bacterial biofilm in the discipline of biomedical and pharmaceuticals and as well as remarkable larvicidal agent to the constraint of mosquito vectors.

1. Introduction

Mosquitoes are dangerous for millions of lives, including animals worldwide, since their action as a transmitter for harmful microorganisms of diverse dreadful diseases [1,2]. *Aedes aegypti* and *Culex tritaeniorhynchus* are the foremost vectors that cause numerous principal diseases like dengue and Japanese encephalitis, respectively [3–5]. Among them, the dengue fever has been proclaimed to increase rapidly, and most spreading mosquito-borne illnesses in the equatorial and subtropical regions around the world [6,7] and the prevention of these

diseases only depends in finding the way to knock down the vector population [3].

The usage of chemical insecticides like diflubenzuron, organophosphates, organochlorines and carbamates has been employed as an effective method to control insects. The continual usage of these insecticides develops resistance in mosquitoes, and they are haphazard and harmful [3,8–12]. Resistance in mosquitoes, risky impact on lives and the earth, there is an extreme necessity towards finding the target-specific, biodegradable, eco-friendly, and cost-effective drugs against mosquitoes [3,12,13].

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jece.2020.104466>

Received 6 August 2020; Received in revised form 5 September 2020; Accepted 7 September 2020

Available online 13 September 2020

2213-3437/© 2020 Elsevier Ltd. All rights reserved.



Original article

Down-regulation of hepatic G-6-Pase expression in hyperglycemic rats: Intervention with biogenic gold nanoconjugate

Sengani Manimegalai^a, Shahid Mahboob^b, Khalid A. Al-Ghanim^b, Fahad Al-Misned^b,
Marimuthu Govindarajan^{c,d}, Krishnan Anbarasu^e, Vijayarangan Devi Rajeswari^{a,*}^aDepartment of Bio-Medical Sciences, School of BioSciences and Technology, Vellore Institute of Technology (VIT), Vellore 632 014, Tamil Nadu, India^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^dDepartment of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India^eDepartment of Bioinformatics, School of Life Sciences, Vels Institute of Science, Technology & Advanced Studies, Pallavaram, Chennai 600 117, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 22 July 2020

Revised 30 August 2020

Accepted 13 September 2020

Available online 22 September 2020

Keywords:

Diabetes

Glucose 6 phosphatase

Couroupita guianensis

Gold nanoparticles

Quantitative RT-PCR

ABSTRACT

Chronic diabetes extensively complicates the glucose metabolism to onset and progress the complication. Concurrently, several contemporary medicines, especially organo-metallic formulations, are emerging to treat hyperglycemia. The current study aims to emphasize the gold nanoparticles (GNPs) potential for glucose metabolism regulation in Streptozotocin (STZ) induced diabetes. Quantitative real-time polymerase chain reaction (RT-PCR) was carried out to detect the mRNA expression of Glucose transporters 2 (GLUT2), Glucokinase (GK) and Glucose 6 Phosphatase (G-6-Pase). The study shows remarkable results such as the prognostic effect of GNPs in reinforcing the repression of enzyme complex G-6-Pase about 13.3-fold when compared to diabetes control. Also, molecular docking studies showed significant inhibition of G-6-Pase by the terpenoid ligands with alpha and beta amyryn from leaf extract of *Couroupita guianensis*. Thus the study explored the novel mechanism of G-6-Pase downregulated by GNPs intervention that majorly contributes to the regulation of circulatory glucose homeostasis during diabetes.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Diabetes mellitus is one of the deadliest disorder that challenges the healthcare system by requiring lifetime treatment (Clarke et al., 2004). High circulatory blood glucose levels represent it, eventually resulting in microvascular abnormalities observed in the retina, glomerulus, and lower extremities of the brain nervous (American Diabetes Association, 2017). Usually, microvascular abnormalities directly accelerate atherosclerosis and consequently increase the risk of myocardial infection (Peluso and Serafini, 2017). Substantially, there is an essential need for new pharmaceutical interventions to discover a cure for hyperglycemia and associated complications. In general, organic anti-oxidants are capable of

controlling oxidative stress, which constrains the complication progression in all types of diabetes (Sies, 1997). Many studies reported that plants and their secondary metabolites are potential biological anti-oxidants and also possess anti-diabetic activity (Martel et al., 2016). Specifically, anthraquinones and its derivatives are proved as potent inhibitors of the critical hydrolyzing enzyme, Glucose 6 phosphatase (G-6-P ase) (Westergaard and Madsen, 2001). The herbal plant *Couroupita guianensis* is majorly encompassed with triterpenes and anthraquinone isatin, which acts as a specific, reversible and competitive G-6-P translocase inhibitor (Shekhawat and Manokari, 2016). The leaf extract suppresses the G-6-Pase expression and declines the hepatic glucose production during gluconeogenesis.

Similarly, 7-hydroxy-5-methoxy-6,8-dimethyl flavanone from *C. guianensis* protects pancreatic β cells from hyperglycemia-induced glucotoxicity (Hu et al., 2012). Altogether, polyphenols effectively enhance glucose uptake and reduce hepatic glucose output by increasing insulin activity (Kumar et al., 2011; Martínez et al., 2012). At the same time, these organic sources are found to be lacking in parameters like dissolution, bioavailability, and sufficient dosage (Yang et al., 2008).

* Corresponding author.

E-mail address: vdevirajeswari@vit.ac.in (V. Devi Rajeswari).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



OPEN

Chitosan overlaid Fe₃O₄/rGO nanocomposite for targeted drug delivery, imaging, and biomedical applications

Viswanathan Karthika^{1,2}, Mohamad S. AlSalhi^{2,3}✉, Sandhanasamy Devanesan^{2,3}, Kasi Gopinath¹, Ayyakannu Arumugam⁴ & Marimuthu Govindarajan^{5,6}

A hybrid and straightforward nanosystem that can be used simultaneously for cancer-targeted fluorescence imaging and targeted drug delivery *in vitro* was reported in this study. A chitosan (CS) polymer coated with reduced graphene oxide (rGO) and implanted with Fe₃O₄ nanoparticles was fabricated. The fundamental physicochemical properties were confirmed via FT-IR, XRD, FE-SEM, HR-TEM, XPS, and VSM analysis. The *in vivo* toxicity study in zebrafish showed that the nanocomposite was not toxic. The *in vitro* drug loading amount was 0.448 mg/mL⁻¹ for doxorubicin, an anticancer therapeutic, in the rGO/Fe₃O₄/CS nanocomposite. Furthermore, the pH-regulated release was observed using folic acid. Cellular uptake and multimodal imaging revealed the benefit of the folic acid-conjugated nanocomposite as a drug carrier, which remarkably improves the doxorubicin accumulation inside the cancer cells over-express folate receptors. The rGO/Fe₃O₄/CS nanocomposite showed enhanced antibiofilm and antioxidant properties compared to other materials. This study's outcomes support the use of the nanocomposite in targeted chemotherapy and the potential applications in the polymer, cosmetic, biomedical, and food industries.

One current focus of the scientific community is combating cancer using nanotherapeutics, chemotherapy, and gene therapy. Current strategies for the drug delivery of cancer therapeutics are focused on increasing the drug concentration at the target site and reducing the systemic distribution. In this regard, functionalized nanoparticles (NPs) are of interest because they can prevent the systemic metabolism and subsequent elimination of the drug, thus ensuring pharmacological effect with less toxicity¹. Advancements in nanoscience have facilitated the selective transportation of drugs into the target site by a unique mechanism (ligand-mediated targeting and receptor-mediated targeting molecules)—this makes to deliver the specific antitumor molecule to the cancer tissue. The sensitive nanocarriers are size- and dimension-dependent. After exposure to an external stimulus, such as pH, enzymatic systems, magnetic gradient, temperature, and ultrasound, the nanocarriers change their physicochemical properties^{1,2}. Superparamagnetic Fe₃O₄ NPs have gained attention for their multifunctionality, including target-based carriage, localized hyperthermia therapy, stem cell labeling and tracking, and contrast agents for magnetic resonance imaging (MRI)^{3–6}.

Additionally, nanocomposites with these superparamagnetic NPs and a 2-dimensional graphene derivative with a large surface area can be used to rational design the drug delivery system. Thus, nanocomposites with a high drug loading capacity and magnetically controlled carrier can be accomplished simultaneously. The preferred graphene derivative for drug loading and delivery efficiency is graphene oxide (GO), because its viable surface chemistry contains layers of graphene sheets with various organic functional groups, such as carboxylic acid, epoxide, and hydroxide, on its surface⁷. The peripheral carboxylic group (–COO) can stabilize the colloidal system and create a pH-responsive negative charge surface. However, the basal plane having hydroxyl (–OH)

¹Department of Packaging, Yonsei University, 1 Yonseidaegil, Wonju-si, Gangwon-do 26493, South Korea. ²Research Chair in Laser Diagnosis of Cancers, College of Science, Department of Physics and Astronomy, King Saud University, Riyadh 11451, Kingdom of Saudi Arabia. ³Department of Physics and Astronomy, College of Science, King Saud University, Riyadh 11451, Kingdom of Saudi Arabia. ⁴Department of Botany, Alagappa University, Karaikudi, Tamil Nadu 630 003, India. ⁵Department of Zoology, Unit of Vector Control, Phytochemistry and Nanotechnology, Annamalai University, Annamalainagar, Tamil Nadu 608 002, India. ⁶Department of Zoology, Unit of Natural Products and Nanotechnology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu 612 001, India. ✉email: malsalhy@gmail.com



Research paper

Facile synthesis and biophysical characterization of egg albumen-wrapped zinc oxide nanoparticles: A potential drug delivery vehicles for anticancer therapy



Tharumasivam Siva Vijayakumar^a, Shahid Mahboob^b, Giridharan Bupesh^{c,*}, Sakthivel Vasanth^c, Khalid A. Al-Ghanim^b, Fahad Al-Misned^b, Marimuthu Govindarajan^{d,e,**}

^a PG & Research Department of Biotechnology, Srimad Andavan Arts & Science College, Srirangam, Tiruchirappalli, 620005, India

^b Department of Zoology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Research & Development Wing, Sree Balaji Medical College & Hospital (SBMCH), Bharath University, BIHER, Chrompet, Chennai, 600044, Tamil Nadu, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^e Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

ZnO nanoparticles
Egg white (albumin)
Anticancer activity
RT-PCR
Protein expression
Drug development

ABSTRACT

In the present investigation, an innovative method was evolved for the synthesis of egg albumin doped Zinc oxide nanoparticles (Ea-ZnO NPs). The biophysical characterizations of Ea-ZnO NPs were assessed by UV-Vis., FT-IR, XRD, TEM and DLS analysis. UV spectrum exhibited a distinctive absorption peak at 310 nm. The synthesized Ea-ZnO NPs materialized as a spherical and hexagonal platelet-like structure with 20–60 nm in diameter through TEM. The anticancer and biocompatibility potential of Ea-ZnO NPs was evaluated against MCF-7 cancer cell line. Anticancer activity of Ea-ZnO NPs on MCF-7 by MTT assay having significant cytotoxicity and proportionately decreased cellular viability. Gene expression study (RT-PCR) and Western blot analysis indicated that the Ea-ZnO NPs stimulated ROS, which in turn up regulated the transcription of mRNA levels of apoptotic genes such as p53, bcl-2, caspase-3 and caspase-9 while significantly downregulated the expression of anti-apoptotic gene Bcl-2. The outcome deliberates that Ea-ZnO NPs can specifically repress the gene expression of MCF-7 by ROS damage and cytotoxicity intervened cell death. Overall, the Ea-ZnO NPs can be a potential drug for anticancer therapy.

1. Introduction

Presently nanotechnology has brought a fantastic consideration due to more extensive uses of nanoparticle in multiple fields of science. Nanotechnology has risen at the leading edge of science and innovation for many recent decades. For the foremost part, it includes the creation of gadgets or materials in the range of nanometer extend, for instance, 1–100 nm. Metal oxides nanoparticles were widely used as functional materials in various applications. Among metal oxides, FeO, TiO₂, ZnO, MgO, and CaO have drawn incredible consideration as a result of their more extensive applications in pharmaceutical, which includes anticancer, antidiabetic, anti-inflammatory antimicrobial, antifungal and antiviral properties. The key functional material has identified by their compatible properties, piezoelectric and biphasic semiconductor

capacities. Those properties are significantly present in Zinc oxide (ZnO); thus, it has used in biosensors, optoelectronic & photovoltaic gadgets, lasers transducers [1,2] lead (drug) transporters and cosmetics [3]. The biosafety and molecular properties of nanoparticles are still not investigated by successful synthesizers. Many studies will be needed to educate and substantiate the uses and their environmental and health risks of the nanoparticles [4].

In general, ZnO NPs are synthesized by mechanical, physical, chemical, and biological methods. Nevertheless, specific strategies are applied for the synthesis of ZnO NPs such as sol-gel [5,6], chemical co-precipitation [7], hydrothermal [8–10], flame spray-pyrolysis [11–15], microwave mediated [16–19] and thermal decomposition [20]. However, these techniques are quite high energy, prolonged timing and are not eco friendly because of multistep synthesis.

* Corresponding author.

** Corresponding author. Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India.

E-mail addresses: bupeshgiri55@gmail.com (G. Bupesh), drgovind1979@gmail.com (M. Govindarajan).

<https://doi.org/10.1016/j.jddst.2020.102015>

Received 17 July 2020; Received in revised form 8 August 2020; Accepted 11 August 2020

Available online 23 August 2020

1773-2247/© 2020 Elsevier B.V. All rights reserved.



Original article

Entomofaunal survey and larvicidal activity of greener silver nanoparticles: A perspective for novel eco-friendly mosquito control

Kuppusamy Elumalai ^a, Shahid Mahboob ^b, Khalid A. Al-Ghanim ^b, Fahad Al-Misned ^b, Jeganathan Pandiyan ^c, Putta Muralidharan Kousik Baabu ^c, Kaliyamoorthy Krishnappa ^{c,*}, **Marimuthu Govindarajan** ^{d,e,**}^aDepartment of Advanced Zoology & Biotechnology, Government Arts College for Men (Autonomous), Chennai 600035, Tamil Nadu, India^bDepartment of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cDepartment of Zoology and Wildlife Biology, A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai 609305, India^dUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, 608 002 Tamil Nadu, India^eDepartment of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 7 July 2020

Revised 22 July 2020

Accepted 30 August 2020

Available online 4 September 2020

Keywords:

Mosquito density and diversity

Physico-chemical characteristics

Blumea mollis

Green synthesis

Larval toxicity

ABSTRACT

The entomofaunal survey and its toxicity of *Blumea mollis* (Asteraceae) leaf aqueous extract-mediated (*Bm*-LAE) silver nanoparticles (AgNPs) were assessed against selected human vector mosquitoes (HVMs). A total of 1800 individuals of 29 species belongs to 7 genera were identified. Month-wise and Genus-wise abundance of HVMs larval diversity were calculated and one-way ANOVA statistically analyzed the average physico-chemical characteristics. The relationship between physicochemical characteristics and HVMs larvae in KWS was interpreted. The total larval density and container index were 23530.18 and 1961.85 examined against 10 different containers. Various spectroscopic and microscopic investigation characterized *Bm*-AgNPs. The *Bm*- AgNPs tested against HVMs larvae, the predominant LC₅₀/LC₉₀ values of 18.17/39.56, 23.45/42.49 and 21.82/40.43 µg/mL were observed on *An. subpictus* Cx. *vishnui* and *Ae. vittatus*, respectively. The findings of this investigation, improperly maintained drainages, containers and unused things in study sites, are engaged to HVMs development. This will be essential for designing and implementing HVMs control. The larval toxic potentiality of *Bm*- AgNPs had a prompt, inexpensive and compelling synthesis of multi-disperse action against HVMs.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Mosquitoes are well adapting to breed and survive in various extreme habitats (Senthamarai Selvan et al., 2015; Senthamarai Selvan and Jebanesan, 2016; Franklin and Whelan, 2009). World-wide, mosquitoes are great blood-sucking ectoparasites which provide major health issues to the public. Due to the climate change, urbanization, artificial containers, availability of food sources are

significant factors for utilization and rapid multiplication of human vector mosquitoes (HVMs) (Gokulakrishnan et al., 2012; Krishnappa and Elumalai, 2012a; Thomson, 2014; Rafael et al., 2019). They are transmitting many dreadful vector-borne diseases (VBDs) to public as the results causing millions of mortality and morbidity has been occurring every year (Govindarajan et al., 2005, 2013; WHO Zika virus report, 2016a; WHO malaria report, 2016b; WHO malaria report, 2018; WHO filariasis report, 2019; Govindarajan et al., 2016a; Benelli et al., 2017a). The VBDs are mainly transmitted by four genera of *Aedes*, *Anopheles*, *Culex*, and *Mansonia* of mosquitoes (Sivakumar et al., 2011; Benelli, 2015; Benelli and Govindarajan, 2017; Benelli et al., 2018). Recently, VBDs are very challenging problems and gives social-economic crisis in worldwide (Krishnappa et al., 2012a; Govindarajan and Rajeswary, 2015; Govindarajan et al., 2016b; Benelli et al., 2017b; Rekha et al., 2019).

Aedes is a cosmopolitan major HVM they breeds and proliferates in different household water containers even in a tree hole, flower pot, etc., (Govindarajan and Sivakumar, 2012; Govindarajan et al., 2013; Albaba et al., 2015; Sudeep and Shil, 2017). It considerably

* Corresponding author.

** Corresponding author at: Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, 608 002 Tamil Nadu, India.

E-mail addresses: krishnappaprofessor@gmail.com (K. Krishnappa), drgovind1979@gmail.com (M. Govindarajan).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

Research Article

Chinnasamy Balalakshmi*, Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, Khalid F. Alanzi, Kasi Gopinath, Ayyakannu Arumugam, and Marimuthu Govindarajan*

Development of chitosan/agar-silver nanoparticles-coated paper for antibacterial application

<https://doi.org/10.1515/gps-2020-0070>

received September 19, 2020; accepted November 19, 2020

Abstract: The radical proliferation of pathogenic bacteria and their infections causes significant issues for human health and the environment. Today, biopolymers are used to produce different nanoparticles. In the present investigation, the fabricated chitosan/agar-silver nanoparticles (Cht/Ar-AgNPs)-coated papers were tested for antibacterial applications. Agar was used as a reducing agent for the synthesis of AgNPs. Synthesized Ar-AgNPs were examined through optical, phase crystallinity and topological analysis. Cht and Ar-AgNPs solution was mixed with various ratios of 9:1, 8:2, 7:3, 6:4, and 5:5 by weight. In addition to that, the conformity of Cht/Ar-AgNPs-coated papers was characterized by structural, spectral, and morphological analysis. However, Cht/Ar-AgNPs-coated papers were subjected to antibacterial properties. The ratio of (6:4) Cht/Ar-AgNPs-coated paper showed excellent antibacterial agent, and it can be used as extending the food product shelf life.

Keywords: drug development, cellulose paper, bar-coating, antibacterial activity, reactive oxygen species

1 Introduction

Biopolymers are considered an ideal candidate for nanotechnology's recent advent due to the biocompatibility, reducing/capping ability, and different physicochemical properties, especially nontoxicity to an environment. Depending on the applications, biopolymers are varied. Moreover, they are highly edible as well as biodegradable films in food packaging. The high recyclable natural biomolecules-based packaging materials like fatty acids, amino acids, and polysaccharides are convenient environmental agents than petrochemical-based polymers. Biopolymer films and coatings are used for many applications: it serves as the best food packaging material, minimizes food deterioration, extends food life by serving as solute and gas barriers, and acts as various additives (antioxidants, antimicrobials, coloring agents, and nutrients) [1–5].

The biopolymers and papers-associated materials are providing an eco-friendly approach. Paper has unique characters as well as used for multiple applications: less weight, low cost, superior mechanical properties, low environmental impact, paper sensors, filters, indicators, writing/printing prepossess, packaging, household products, resistance from insufficient grease and oil, low barrier properties [6–10], and due to its lipophilic nature, it is prone to react with fatty molecules and leads to damage in printed papers, and enables usage as synthetic polymers, which leads to recycling issues and causes environmental problems [6–11]. Moreover, many researchers pushed towards the advanced and new approaches for developing antimicrobial papers by administering eco-friendly synthesis processes [12–14].

Many researchers are likely attracted by agar (Ar) because it is highly renewable, biodegradable, and of

* Corresponding author: Chinnasamy Balalakshmi, Department of Nanoscience and Technology, Alagappa University, Karaikudi 630 003, Tamil Nadu, India, e-mail: samyasribala@gmail.com

* Corresponding author: Marimuthu Govindarajan, Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalinagar 608 002, Tamil Nadu, India; Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India, e-mail: drgovind1979@gmail.com

Naiyf S. Alharbi, Shine Kadaikunnan, Jamal M. Khaled, Khalid F. Alanzi: Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

Kasi Gopinath: School of Materials and Energy, Southwest University, Chongqing, 400715, People's Republic of China

Ayyakannu Arumugam: Department of Botany, Alagappa University, Karaikudi 630 003, Tamil Nadu, India



Impact of pesticide monocrotophos on microbial populations and histology of intestine in the Indian earthworm *Lampito mauritii* (Kinberg)

Vaithiyanathasamy Kavitha^{a,b,*}, Ramachandran Anandhan^{a,c}, Naiyf S. Alharbi^d, Shine Kadaikunnan^d, Jamal M. Khaled^d, Taghreed N. Almanaa^d, Marimuthu Govindarajan^{a,b,**}

^a Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, 608 002, India

^b Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India

^c Department of Zoology, Government Arts College (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India

^d Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:

Insecticide
Ecotoxicology
Microbes
Earthworm
Intestine
Histology

ABSTRACT

Soil contamination has enlarged over the decades due to intensive use of pesticides and chemical fertilizers in agronomy. Earthworms are significant organisms in the soil community. Earthworms are the major role in soil fertility in most ecological system and the production of biogenic structures. Moreover, earthworm gut mucus enhances the beneficial soil microorganism potential biological activities. They are used as model organisms for assessing the ecological risks of chemicals. Enrichment of essential nutrients in soil through earthworm is a cost-effective and eco-friendly approach. In India, the organophosphorus pesticide monocrotophos is commonly used to control agricultural pests. Hence, it is important to study the effect of monocrotophos on the gut microbiota in *Lampito mauritii*. A 15-day exposure to a low (1/10th of the LC₅₀ after 96 h i.e., 0.093 ppm kg⁻¹) and high sublethal concentration (1/3rd of the LC₅₀ after 96 h i.e., 0.311 ppm kg⁻¹) of monocrotophos led to reduced proliferation of the gut microbiota in *L. mauritii*. However, exposure for 30 days led to a recuperation of the microbial populations to near control values. Among the eight bacterial and five fungal species that inhabit the gut of *L. mauritii*, only six bacterial and three fungal species were able to survive after exposure to monocrotophos. In addition to the study, histopathological changes were observed in the intestine of *L. mauritii* after application of lower sublethal concentration of monocrotophos. Severe pathological changes such as vacuolization, degenerated nuclei, damaged villi and congestion of the blood sinuses were noticed in the intestine on 1st and 5th day of the experiment. But in 30th day the damages were slowly recovered due to degradation of monocrotophos by the presence of some pesticides degrading bacterial and fungal species and regenerative capability of chloragogen cells in the intestine. The results suggested that reduced microbial populations and pathological damages in intestine were observed during the application of monocrotophos. So, the monocrotophos have several harmful impacts on earthworms.

1. Introduction

There is a growing concern for soil contamination due to the widespread and global use of agropesticides. Knowledge about the effect(s) of pesticides in the soil environment is imperative since the resident soil organisms may be adversely affected. Depending on the type of soil organisms affected, this may have damaging consequences for soil fertility. A large proportion (80%) of the biomass of terrestrial invertebrates comprises earthworms that are important for structuring and increasing the nutrient content of soil. Therefore, they are suitable bioindicators for the chemical contamination of soil in terrestrial

ecosystems that can be used to provide an early warning for deteriorating soil quality [1,2]. This is important for protecting the natural environments and is of increasing interest in protecting human health [3] as well as other terrestrial vertebrates that prey on earthworms. Earthworms are suitable bioindicators of soil toxicity since they ingest large quantities of decomposed litter, manure, and other organic matter deposited on soil, thereby aiding the conversion into rich topsoil [4]. The pesticides in soil may accumulate in worms via ingesting contaminated soil or the absorption from soil water. Monocrotophos (dimethyl (E)-1-methyl-2-(methylcarbamoyl) vinyl phosphate) is an organophosphorus pesticide that is used to control a broad spectrum of

* Corresponding author. Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, 608 002, India.

** Corresponding author. Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India.

E-mail addresses: kavisamikshaa@gmail.com (V. Kavitha), drgovind1979@gmail.com (M. Govindarajan).



β -glucan extracted from eukaryotic single-celled microorganism *Saccharomyces cerevisiae*: Dietary supplementation and enhanced ammonia stress tolerance on *Oreochromis mossambicus*

Mani Divya^a, Narayanan Gopi^a, Arokiadhas Iswarya^a, Marimuthu Govindarajan^{b,c}, Naiyf S. Alharbi^d, Shine Kadaikunnan^d, Jamal M. Khaled^d, Taghreed N. Almanaa^d, Baskaralingam Vaseeharan^{a,*}

^a *Biomaterials and Biotechnology in Animal Health Lab, Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India*

^b *Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India*

^c *Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India*

^d *Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia*

ARTICLE INFO

Keywords:

β -glucan
Yeast
Ammonia stress
Antioxidant properties
Immune activities
Fish

ABSTRACT

Ammonia is a widespread pollutant that is toxic to living organisms in aquaculture. This study aimed to evaluate the effects of a diet supplemented with beta-glucan from yeast, *Saccharomyces cerevisiae* (*Sc- β G*), on the stress response of *Oreochromis mossambicus* (Tilapia) to ammonia. Fish were divided into four groups, including a control fed a basal diet and three experimental groups fed diets supplemented with *Sc- β G* at 2, 5 and 10 mg/g respectively. After 8 weeks, experimental groups were exposed to ammonia at 100 mg L⁻¹ for 1 week. Growth was measured after the 8-week feeding trial and serum, mucus, and liver tissue were sampled before and after the ammonia challenge. Compared with the control diet, feed supplemented with *Sc- β G* at 10 mg/g significantly ($p < 0.05$) improved growth performance (7.8–9.9 g increase in weight). The cellular immune responses (myeloperoxidase, reactive oxygen species, and reactive nitrogen species), humoral immune responses (alkaline phosphatase, lysozyme, and peroxidase inhibition), and antioxidant response (catalase, superoxide dismutase, and glutathione) were tested in serum, mucus and liver tissue. Compared with the control, these responses were significantly ($p < 0.05$) enhanced at 10 mg/g supplementation with *Sc- β G*. This study demonstrates that *Sc- β G* may be applied to induce stress tolerance and improve growth performance in aquaculture.

1. Introduction

Aquatic animals and fish are consumed for their nutritional benefits in many Asian countries [1]. The considerable growth in aquaculture production observed in the last 50 years has improved the global nutrition security [2]. With reference to the food and agricultural organization (FAO), utilization of aquatic commodities such as fish by human diet reached up to 47 % in 2006 and is expected to reach 50 % in the next few years [3]. The global per capita fish consumption increased from 19.7 kg in 2013 to above 20 kg in 2015 [4]. The fish dietary production has hastily at 5.0 % per year in the preceding five decades [2]. The dietary supplementation practice has been utilized in the aquatic field since 2000. However, often aquaculture is not economically viable due to varying demand and value across fish species

and consumers [5].

Although aquatic organisms can be cultured at high densities, infectious diseases are often a major constraint to the aquaculture industry. In general, the occurrence of diseases is caused by several factors, a few of which are considered relevant to fish and aquatic pathogens [6]. Further, there is an evidence that high level of pollutants increase mortality in aquatic species [7]. To avoid environmental impacts, intensive culture systems have been utilized to manage ammonia levels and mitigate the effect on fish health. Stressors can exacerbate the presence of diseases in impacted habits and frequent stressors can directly lead to fish mortality [8]. Among environmental stressors, ammonia is a common aquatic pollutant and is toxic to fish. Ammonia is the most important compound in intensive culture systems as it causes stress and damage to fish tissue, even in small amounts, and can be

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.micpath.2019.103917>

Received 11 November 2019; Received in revised form 3 December 2019; Accepted 9 December 2019

Available online 10 December 2019

0882-4010/ © 2019 Elsevier Ltd. All rights reserved.



Green fabrication, characterization and antibacterial potential of zinc oxide nanoparticles using *Aloe socotrina* leaf extract: A novel drug delivery approach



Basheer Ahmed Fahimmunisha^a, Ramachandran Ishwarya^a, Mohamad S. AlSalhi^{b,**}, Sandhanasamy Devanesan^b, Marimuthu Govindarajan^{c,d}, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh, 11451, Saudi Arabia

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

Nanotechnology
Confocal laser scanning microscope
XRD
SEM
TEM
Drug delivery
Antibiofilm

ABSTRACT

Currently, nanotechnology is a prolific area of scientific research. As they can be synthesized in an eco-friendly manner, zinc oxide nanoparticles (ZnO NPs) provide an attractive solution to combat bacterial infection. In this study, synthesis of ZnO NPs using *Aloe socotrina* (As) leaf extract and characterized by UV, FTIR, XRD, SEM, and TEM. UV spectroscopy analysis showed that the maximum absorption was at about 315 nm. TEM images of the ZnO NPs definite the structure of ZnO NPs ranging from 15 to 50 nm in size while XRD spectra revealed their crystallinity. To understand the antibacterial activity of NPs, extensive experiments were performed using several assays such as agar well diffusion, minimum inhibitory concentration, growth kinetics, intracellular uptake, reactive oxygen species (ROS) generation, and antibiofilm activity. The *A. socotrina* capped ZnO NPs (As-ZnO NPs) revealed significant activity against biofilms formed by four bacterial pathogens, which can interfere with the management of drug resistant bacterial diseases caused by these biofilms. In summary, this novel biosynthesis technique of As-ZnO NPs with potent bactericidal activity offers an effective solution for the management of UTI pathogens.

1. Introduction

In recent years, urinary tract infection (UTI) has emerged as one of the most widespread diseases in the world. This condition is characteristically aggressive and has contributed to the rise in nosocomial bacterial infection in ambulatory and hospitalized patients as the UTI can spread from the urethra to the bladder [1]. The annual incidence of UTI is approximately between 150 and 250 million cases globally and accounts for about 8 million doctor visits. Moreover, approximately 10% of all hospitalized cases contract these infections despite the sterile hospital conditions [2]. Prolonged use of urinary catheters increases the risk of infection and directly contributes to the rise in UTI [3]. Development of biofilms on the exterior of the catheter is one of the main problems associated with the management of UTI [4,5] as it can prolong the presence of the microorganisms resulting in persistent infections that tend to be resistant to antimicrobial therapy leading to


chronic disease. Some of the most common microbes that infect the urinary catheter and produce biofilms are *Escherichia coli*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Enterococcus*, *Staphylococcus aureus*, *Proteus mirabilis*, *Klebsiella pneumoniae* and *Morganella morganii* [6]. In recent times, antibiotic resistance is an important global public health issue due to the uncontrolled use of antibiotics resulting in the development of resistance in bacteria including uropathogens [7]. Therefore, the antibiotics that currently used have become ineffective and there is an increasing need to develop more efficient antibiotics against specific microbial populations.

Since the early years, microbial pathogens have posed a huge challenge to the scientific community with microbial contamination being one of the major concerns [8]. Most antimicrobial agents have frequent issues such as instability at high temperature or pressure and are often toxic and cause irritation. Moreover, development of antibiotic resistant pathogens has turned into a severe health problem and

* Corresponding author.

** Corresponding author.

E-mail addresses: malsalhi@ksu.edu.sa (M.S. AlSalhi), vaseeharanb@gmail.com (B. Vaseeharan).



Isolation of β -glucan from *Eleusine coracana* and its antibiofilm, antidiabetic, antioxidant, and biocompatible activities

Mani Divya^a, Sivashanmugam Karthikeyan^b, Cyril Ravi^c, Marimuthu Govindarajan^{b,e},
Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f, Taghreed N. Almanaa^g,
Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b School of Bioscience & Technology, VIT University, Vellore, 632014, Tamilnadu, India

^c Department of Zoology and Microbiology, Thiagarajar College (Autonomous), Madurai, 625 009, Tamil Nadu, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:

Finger millet
 β -Glucan
Polysaccharides
Biomolecules
Antibacterial activity
Hemolysis assay

ABSTRACT

Eleusine coracana (Finger millet) has high nutritional value with numerous health benefits and is of low cost. Isolation of beta-glucan (β G) from *E. coracana* (*Ec-βG*) has gained increasing research attention. UV-vis spectroscopy used to measure the surface plasmon resonance at 361 nm to confirm the presence of polysaccharides (glucan molecules) in *Ec-βG*. X-ray diffraction analysis of *Ec-βG* displayed a crystalline nature and confirmed the presence of the β G molecule. Further, the bioactive compounds of *Ec-βG* were screened using gas chromatography-mass spectrometry. The antibacterial activity of *Ec-βG* against both Gram-positive (*Lysinibacillus fusiformis*, *Enterococcus faecalis*) and Gram-negative (*Proteus vulgaris*, *Shigella sonnei*) bacteria were assessed through minimum inhibitory concentrations < 70 μ g/ml of *Ec-βG*. In addition, the antibiofilm activity and bacterial viability of *Ec-βG* at 100 μ g/ml was confirmed by light and confocal laser scanning microscopy. Furthermore, *Ec-βG* inhibits α -amylase and α -glucosidase at an IC₅₀-value of 1.23 and 1.42 μ g/ml, respectively. Superoxide anion scavenging activity at IC₅₀-1.4 μ g/ml and DPPH radical scavenging activity at IC₅₀-1.2 μ g/ml showed that *Ec-βG* had potential antioxidant property. The *in vitro* hemolysis assay for biocompatibility of *Ec-βG* at 200 μ g/ml showed 0.06 \pm 0.09%. Therefore, *Ec-βG* has the potential to act as a suggestive agent for antibacterial, antidiabetic, and antioxidant activity.

1. Introduction

A sufficient quantity of nutrients is important for the development of a healthy environment. Nutraceuticals are the healthy consumption of foods, which act as the beneficial effects of physical condition and prevention of diseases in human beings [1]. For this reason most of the research has been carried out from the natural molecules preferable for the health benefits. The oversized molecules obligatory for existence, that are built from smaller organic molecules are called biological macromolecules. A macromolecule is a huge molecule made up of smaller subunits. These include polysaccharides, peptides, nucleic acids and lipids. A polysaccharide is one type of macromolecule made up of monosaccharides. An example of a polysaccharide is starch, or cellulose

and glucans [2]. In the intervening time, the benefits of glucan molecules in pharmaceutical antibiotics has the root of emergence the bacterial resistance exertion in the earlier period. For this concern the use of natural molecules has turned out to be an appealing researchable phase [3]. In particular, a macromolecule such as betaglucan (β G) are non-starch homopolysaccharides that are composed of glucose molecules [4]. The polysaccharide the model linkage and block oligomer formation of the β G is essential to the derivation of β -glucans. Cereal β G is encompassed with β -D-glucopyranose, which is associated with a mixture of β -(1–3), (1–4) glycosidic bonds [5]. The physical properties of cereal β G is firmness because of its main structure, molecular size, and macromolecular conformation. To a large extent, the use of cereal β G owing to its various benefits and the possibility of industrial

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.micpath.2019.103955>

Received 25 October 2019; Received in revised form 14 November 2019; Accepted 27 December 2019

Available online 30 December 2019

0882-4010/ © 2019 Elsevier Ltd. All rights reserved.



Antibiofilm and anticancer potential of β -glucan-binding protein-encrusted zinc oxide nanoparticles

Mani Divya^a, Marimuthu Govindarajan^{b,c}, Sivashanmugam Karthikeyan^d, Elumalai Preetham^e, Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f, Taghreed N. Almanaa^f, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608 002, Tamil Nadu, India

^c Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^d School of Bioscience & Technology, VIT University, Vellore, 632014, Tamil Nadu, India

^e School of Ocean Science and Technology, Kerala University of Fisheries and Ocean Studies, Panangad, Kerala, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:

β -Glucan-binding protein
Biofilm
Confocal laser scanning microscopy
Pseudomonas aeruginosa
Cell line

ABSTRACT

β -Glucan-binding protein (β GBP) is important for the rational expansion of molecular biology. Here, zinc oxide nanoparticle (ZnONP) was synthesized using β GBP from the crab *Scylla serrata* (Ss- β GBP-ZnONP). Ss- β GBP-ZnONP was observed as a 100 kDa band on sodium dodecyl sulfate polyacrylamide gel and characterized with UV-vis spectroscopy at 350 nm. X-ray diffraction analysis displayed values consistent with those for zincite. Fourier transform infrared spectroscopy revealed the presence of functional groups, including amide, alcohol, alkane, alkyl halide, and alkene groups. The zeta potential (-5.36 mV) of these particles indicated their stability, and transmission electron microscopy revealed the presence of 50 nm nanocones. Ss- β GBP-ZnONPs were tested at 100 μ g/mL against the gram-positive *Enterococcus faecalis* and gram-negative *Pseudomonas aeruginosa* using confocal laser scanning microscopy and the bacterial viability assay was also performed. The growth of MCF7 breast cancer cells was inhibited following treatment with 75 μ g/mL Ss- β GBP-ZnONPs. Thus, Ss- β GBP-ZnONPs have the ability to control the growth of pathogenic bacteria and inhibit the viability of MCF7 breast cancer cell lines.

1. Introduction

Nanotechnology has widespread applications in the field of science and medicine [1–6]. Nanoparticles (NPs) are used as imaging and therapeutic agents and their surfaces are often coupled with biological molecules to reduce surface energy [7]. Metallic and metal oxide NPs, including gold, silver, copper, titanium dioxide (TiO₂), and zinc oxide (ZnO) particles, play vital roles [8–11]. The ZnONPs have been used for various applications, including cell imaging, drug delivery, and nanomedicine [12]. The ZnONPs are also used in goods such as personal care products and are known to exhibit excellent antibacterial and UV blocking properties. *In vitro* analyses have revealed the biocompatibility of ZnONPs [13]. The ZnONPs serve as non-toxic, biocompatible, and safe drug carriers. Zinc is an important trace element present in all body tissues, including the brain, muscle, bone, and skin [14]. Zinc is also

important for eukaryotes, as it is known to modulate several physiological functions [15]. In biological systems, zinc has important roles in metabolic pathways such as the carbohydrate, lipid, nucleic acid, and protein syntheses [16]. The NPs may interact with biomolecules such as nucleic acids, proteins, and lipids, owing to their nano size [17]. Biological synthesis of NPs is more labor- and time-intensive and expensive than physical and chemical methods [18].

Cancer is an emerging disorder that is severely affecting human health. It is the second leading cause of mortality worldwide. The prevalence of breast cancer is higher among women [19] and corresponds to one-fifth of all cancer cases worldwide. Studies have confirmed the cytotoxicity of ZnONPs on cancer cells, including HepG-2, MCF-7, HeLa, U87, and S91 cells [20]. To improve their targeting ability, these NPs are often coupled with antibodies or peptides that bind to the receptors over expressed on cancer cells. Today, different

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).



Curcumin-encased hydroxyapatite nanoparticles as novel biomaterials for antimicrobial, antioxidant and anticancer applications: A perspective of nano-based drug delivery

Saleth Sebastiammal^{a,b}, Arul Sigamani Lesly Fathima^{a,*}, Sandhanasamy Devanesan^{c,d},
Mohamad S. AlSalhi^{c,d,***}, Johnson Henry^e, Marimuthu Govindarajan^{f,g},
Baskaralingam Vaseeharan^{h,**}

^a Department of Physics, Holy Cross College (Autonomous), Nagercoil, 629004, Tamil Nadu, India

^b Reg.No:17213042132003, Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli, 627012, Tamil Nadu, India

^c Research Chair in Laser Diagnosis of Cancers, Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh, 11451, Saudi Arabia

^d Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh, 11451, Saudi Arabia

^e Department of Physics, Manonmaniam Sundaranar University, Abishekapatti, Tirunelveli, 627012, Tamil Nadu, India

^f Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, 608 002, Tamil Nadu, India

^g Unit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam, 612 001, Tamil Nadu, India

^h Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Science Campus, Karaikudi, 630 004, Tamil Nadu, India

ARTICLE INFO

Keywords:

Nanotechnology
Curcumin
Hydroxyapatite
Anticancer
Biological applications

ABSTRACT

Hydroxyapatite (HAp) is an important biological molecule and principal mineral component found in bones that is used in various clinical applications to significantly enhance the biological activity and biocompatibility of man-made biological molecule. Several analyses have been performed to control the structural properties of HAp by varying the processing parameters. This study described the synthesis of HAp nanoparticles (NPs) via the sol-gel method, that would be effective for biomedical applications, such as bone graft substitution. It also examined the effects of pure HAp, surfactant (PEG and CTAB)-assisted HAp NPs, and curcumin-mediated HAp NPs. The synthesized HAp NPs were analyzed using XRD, FTIR, RAMAN, FESEM, TEM, EDAX, UV-DRS, and PL analysis. From the XRD analysis, it was found that the prepared HAp NPs have a hexagonal structure with nanosize distribution. From FESEM and TEM analyses, it was found that the synthesized HAp NPs were rod-like in shape and the elemental analysis was conducted using EDAX. In addition, the biological applications were investigated, viz., antibacterial, antifungal, antioxidant, and anticancer activity. The synthesized HAp nanoparticles possessed excellent antibacterial, antifungal, antioxidant, and anticancer activities. Overall, curcumin-encased HAp nanoparticles can play a vital role in a wide range of fields, including water treatment, food preservation, wound dressing, nanomedicines, and cosmetics as biocidal and disinfecting agents.

1. Introduction

Hydroxyapatite (HAp) nanoparticles (NPs) have received considerable attention over the last decade due to their significant biological applications [1–5]. HAp plays a vital role in biomaterial engineering due to its enhanced biocompatibility and bioactivity [6–9]. The

stoichiometric ratio of calcium and phosphorous in HAp is 1.67, which resembles that of human bones, and calcium phosphate (CaP) is the major mineral constituent of vertebrae bones and teeth [10–12].

HAp [$\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$] is a CaP-based biomaterial primarily used in dental and orthopedic applications because of its structural and chemical similarity to bones, teeth, and enamel [13–15]. HAp acts as an

* Corresponding author.

** Corresponding author.

*** Corresponding author. Research Chair in Laser Diagnosis of Cancers, Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box -2455, Riyadh, 11451, Saudi Arabia.

E-mail addresses: leslysat@gmail.com (A.S. Lesly Fathima), malsalhi@ksu.edu.sa (M.S. AlSalhi), vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jddst.2020.101752>

Received 26 March 2020; Received in revised form 10 April 2020; Accepted 12 April 2020

Available online 18 April 2020

1773-2247/ © 2020 Elsevier B.V. All rights reserved.



Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com

Original article

Swift production of rhamnolipid biosurfactant, biopolymer and synthesis of biosurfactant-wrapped silver nanoparticles and its enhanced oil recovery

Vadivel Tamil Elakkiya^a, Periyasamy SureshKumar^{a,*}, Naiyf S. Alharbi^b, Shine Kadaikunnan^b, Jamal M. Khaled^b, Marimuthu Govindarajan^{c,d,*}^aDepartment of Biotechnology, BIT Campus, Anna University, Tiruchirappalli 620024, India^bDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cDepartment of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 24 February 2020

Revised 10 March 2020

Accepted 2 April 2020

Available online 10 April 2020

Keywords:

Nanotechnology

Pseudomonas aeruginosa TEN01

Interfacial tension

Sand packed column

Enhanced oil recovery

ABSTRACT

Microbial enhanced oil recovery (MEOR) is a kind of enhanced oil recovery (EOR) development, often used as a tertiary stage where oil recovery is no longer possible utilizing primary and secondary conventional techniques. Among a few potential natural operators valuable for MEOR, biosurfactants, biopolymers and biosurfactant based nanoparticles assume key jobs. Biosurfactant which are produced by microorganisms' act as are surface active agents that can be used as an alternative to chemically synthesized surfactants. *Pseudomonas aeruginosa* TEN01, a gram-negative bacterium isolated from the petroleum industry is a potential biosurfactant (Rhamnolipid) producer using cassava waste as the substrate. This work focuses on production and characterization of rhamnolipid from *P. aeruginosa* TEN01 and its use in enhanced oil recovery. The effectiveness of Chitosan that is deacetylated form of chitin which is a biopolymer that provides density and viscosity to the fluids is not known in enhanced oil recovery yet and so it is studied. Moreover, the fabrication of biosurfactant-mediated silver nanocrystals and its application in enhanced oil recovery is also studied. Sand-Pack column was constructed and the mechanism of oil recovery in the column was studied. While incubating the crude oil containing sand packed column with Biosurfactant-biopolymer and brine flooding in the ratio of 1:2, and Biosurfactant incubation - flooding with 3 g/l of biopolymer was found to be 34.28% and 44.5% respectively. The biosurfactant based silver nanoparticles are non-toxic and have better stability when compared to chemically synthesized silver nanoparticles. The oil recovery percentage by chemical based Ag NPs and biosurfactant based Ag NPs are 14.94% and 14.28% respectively.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

From the earliest starting point of civic establishments, we are constantly subject to energizes, for example, wood, whale-oil or petroleum derivatives - coal, and raw petroleum related items (lamp oil, diesel, oil). Among each one of those diverse sort of energizes, non-renewable energy source assumed a major job in mechanical goals (Silva et al., 2014). Increase in oil costs based on market interest disparities are observed as of late resulted a decrease which is relied upon to remain at 30–60 \$/bbl oil for a prolonged duration (Schulz, 2016; Banat, 1995; McInerney et al., 2005). Current situation of such an uncertain market at oil costs is a very tough path ahead for oil ventures and those nations which are exceptionally reliant on oil-based economy. Nevertheless, the

* Corresponding authors at: Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India (M. Govindarajan). Department of Biotechnology, BIT Campus, Anna University, Tiruchirappalli 620024, India (P. SureshKumar).

E-mail addresses: drsureshbiotech2003@gmail.com (P. SureshKumar), drgovind1979@gmail.com (M. Govindarajan).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Contents lists available at ScienceDirect

Saudi Journal of Biological Sciences

journal homepage: www.sciencedirect.com

Original article

Identification of a novel antibacterial protein from hemolymph of freshwater zooplankton *Mesocyclops leuckarti*



Varadhan Praveena^a, Sournamanickam Venkatalakshmi^{a,*}, Naiyf S. Alharbi^b, Shine Kadaikunnan^b, Jamal M. Khaled^b, **Marimuthu Govindarajan^{c,d,*}**

^aCentre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

^bDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^cDepartment of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 29 February 2020

Revised 5 April 2020

Accepted 3 May 2020

Available online 11 May 2020

Keywords:

Mesocyclops leuckarti

Zooplankton

Hemolymph

Antimicrobial proteins

SDS-PAGE

ABSTRACT

Bacterial infections are the most important problem of health care worldwide. The hemolymph antibacterial proteins of *Mesocyclops leuckarti* was isolated for the first time and its antibacterial efficacy was evaluated against four different human pathogenic microbes viz., *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumonia* and *Shigella flexneri*. The antibacterial potential of the antimicrobial proteins of hemolymph samples from plankton cultured in water enriched with Cow Urine Distillate (CUD) was compared with normal ones. The results indicated that the hemolymph proteins were more potential against Gram negative bacteria than Gram positive bacteria. *Klebsiella pneumonia* was more susceptible to the hemolymph proteins exhibiting a zone of inhibition measuring 27 mm. The supplement of CUD to the culture media further enriched the antibacterial activity of the hemolymph proteins (29 mm). The SDS-PAGE analysis indicated two different types of clear bands representing proteins of 53 kDa and 19 kDa. Overall, this investigation signified that the microcrustaceans have a defence mechanism hemolymph of *Mesocyclops leuckarti* have a potential agent for novel antibiotics.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Copepods represent about 20% of the mean annual zooplankton biomass. They are found in abundance in many aquatic ecosystems (Huys and Boxshall, 1991) and conspicuous member of aquatic biota that have adapted to various habitats such as damp moss, interstitial sand, subterranean localities and parasitic living. The cyclopoid copepods are successful inland water group animals and *Mesocyclops leuckarti* is the important planktonic cyclopoid copepods. In evolution, invertebrates have not developed acquired immune systems. However, they have been bestowed with non-

specific immune mechanisms. Crustaceans show many antibacterial proteins in their hemolymph. However, the literature does not show any record of anti bacterial proteins in microcrustaceans including zooplankton (Iskratsch et al., 2009; Iwanaga, 1993; Kawabata et al., 1995; Mori and Stewart, 1978; Jayasankar and Subramoniam, 1999) (Table 1).

Hemolymph is the type of blood found in Arthropod's open circulatory system. It contains many bioactive molecules which have functional roles in the defence system. The molecules include lectins, complement, clotting factors, antimicrobial peptides (Vazquez et al., 2009).

Among them the antimicrobial peptides are the prime factors that give immunity to the animal. There are two types of antimicrobial peptides identified in hemolymph. They are high molecular weight large antimicrobial proteins (>100 amino acids) and low molecular weight small antimicrobial proteins. The high molecular weight antimicrobial proteins target the disrupt microbial biomolecules and small antimicrobial proteins disrupt the structure and/or the function of microbial cells (Aspan et al. (1995); Stabili et al. (1999); Fujimoto et al. (1995); Hall et al. (1995)). These antimicrobial peptides are secreted in

* Corresponding authors at: Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India (M. Govindarajan).

E-mail address: dr.s.venkatalakshmi@gcwk.ac.in (S. Venkatalakshmi).

Peer review under responsibility of King Saud University.





Original article

Seed dispersal by ungulates in the point calimere wildlife sanctuary: A scientific and perspective analysis



Govindaraju Chandru^a, Jeganathan Pandiyan^a, Vikramathithan Durga^a, Marimuthu Govindarajan^{b,c,*}, Naiyf S. Alharbi^d, Shine Kadaikunnan^d, Jamal M. Khaled^d, Chellasamy Panneerselvam^e, Kaliyamoorthy Krishnappa^{a,*}

^aDepartment of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai 609305, India

^bUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^cDepartment of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

^dDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

^eDepartment of Zoology, School of Life Sciences, Bharathiar University, Coimbatore 641 046, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 5 March 2020

Revised 22 June 2020

Accepted 23 June 2020

Available online 30 June 2020

Keywords:

Exotic woody weed plants

Prosopis juliflora

Ungulates

Seed dispersal

Germination test

ABSTRACT

Exotic woody weed plants are a very serious threat to seed dispersed by ungulate in the tropical forest of Asia. The ungulates in Point Calimere Wildlife Sanctuary (PCWS) are a significant role in native indigenous seed dispersal. The exotic woody weed tree *Prosopis juliflora* prevalence distributed in the PCWS and they might potentially alter the native medicinal plant species. In the present investigation, we have assessed the seed dispersal by ungulates in PCWS from January to March 2017. Four different ungulate species were selected to understand their seed dispersal rate of different plant species in selected sanctuary. This investigation was planned to confirm the seed dispersal by ungulates of blackbuck, spotted deer, wild boar and feral horse. Among the four different ungulates tested, the maximum numbers of pellets collected from blackbuck and no seed found in their pellets. The low quantities of pellets were collected from wild boar and this study has recorded medium-sized ungulates which dispersed variety of plant. However, the dispersal of the seed of medicinal plants were not considerably high and relatively moderate percentage of seeds dispersal occurred in medium-sized ungulates like wild boar and spotted deer. *P. juliflora* had 100% seed germination rate were observed from the faecal samples of wild boar and feral horse. The control seed achieved maximum seedling rate than the ungulates seeds.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Plants are very less mobility in consequently their diverse of seeds dispersal are highly influencing through variety of biotic vectors. The majority of seeds are dispersing through animals by their

digestive system, excreted seeds which fall on the ground and growing it naturally. Usually, that excreted seeds are getting higher potential chances to form new plants which could be far away from the mother plant (Jordano et al., 2010; Chapman et al., 2017; Susan Harrison, 2017; Chaves et al., 2018). Many birds are likely to eat diverse of fruits varieties which help to dispersing seeds to different locality by their dropping. The animals are doing seed dispersal and play a major role towards, success and wealth of environmental (Ingle, 2003; Lozada et al., 2007; Pejchar et al., 2008; Patrick David et al., 2015). The avian fauna are doing long-distance distribution of seed dispersal and accidentally these seeds are dispersed in uneven and critically important flora filling wild and epizoochoary (Gonzales et al., 2009; Chimera et al., 2010; Balasubramanian et al., 2011). Globally, the past three decades forest fire becomes major threat to wild ecosystem in various parts of the continents. As a result of forest fire, most of the indigenous/endemic flora species will become extinct as possible (Satendra and Kaushik, 2014). These problems are recently highlighted in many

* Corresponding author at: Department of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai 609305, India.

** Corresponding author at: Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India.

E-mail addresses: drgovind1979@gmail.com (M. Govindarajan), krishnappapروفessor@gmail.com (K. Krishnappa).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier



Original article

A novel approach to assess the heavy metal content in the feathers of shorebirds: A perspective of environmental research

Jeganathan Pandiyan ^{a,*}, Shahid Mahboob ^b, Rajendran Jagadheesan ^a, Kuppusamy Elumalai ^c, Kaliyamoorthy Krishnappa ^a, Fahad Al-Misned ^b, Zahid Ali Kaimkhani ^d, Marimuthu Govindarajan ^{e,f,*}

^a Department of Zoology and Wildlife Biology A.V.C. College (Autonomous), Mannampandal, Mayiladuthurai 609305, Tamil Nadu, India

^b Department of Zoology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^c Department of Advanced Zoology & Biotechnology, Government Arts College for Men (Autonomous), Chennai 600035, Tamil Nadu, India

^d Department of Anatomy, College of Medicine, King Saud University, Riyadh, Saudi Arabia

^e Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^f Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 8 July 2020

Revised 25 July 2020

Accepted 26 August 2020

Available online 3 September 2020

Keywords:

Pollution
Heavy metals
Wintering grounds
Shorebirds
Habitats

ABSTRACT

Metals are major toxic elements and cause various diseases and damage shorebirds. The study envisaged the heavy metals content in the feathers of shorebirds, prey, water, and sediment from Pichavaram Mangrove Forest (PMF). Primary feathers of shorebirds species such as Curlew Sandpiper, Eurasian Curlew, and Painted Stork were collected and analyzed the following metals viz., Cd, Cu, Cr, Co, Pb, Hg, Ni, and Zn. The levels of Hg, Cr, Pb and Ni varied significantly among the metals studied ($p < 0.05$). The metal contamination pattern of shorebirds followed in the following order: Painted stork > Eurasian Curlew > Curlew sandpiper. However, in the habitat, Cd, Co, Pb, Hg, Ni and Zn varied significantly between water and sediment ($p < 0.05$) and Cd, Cu, Hg, Pb, Ni, and Zn differed significantly among the prey species ($p < 0.05$). Overall, except for Cd, whereas, other seven metals showed significant differences between the feathers of shorebirds and the environment such as water, sediment, polychaetes, mollusc, crabs, prawns and fishes ($p < 0.05$). Nevertheless, the resident bird (Painted stork) had a higher level of metal accumulation than migratory species, showing that PMF is under threat and requires proper monitoring, management and conservation strategies to sustain organisms that depend on it.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Heavy metals are major toxic elements and cause various diseases and damage wildlife when their load exceeds normal levels. Higher content of Cd in feathers of birds can wreck their flying mechanism and lead to poor development of bones (Spahn and

Sherry, 1999). Pb mainly deposits in the tissues of feathers of birds (Jayakumar and Muralidharan, 2011), and a higher content of it could destroy their thermoregulation, growth of nestlings, and recognition of their siblings (Burger and Gochfeld, 2000). The toxicity of Cr had several impacts on birds, such as the development of embryo and hatching success of eggs in Mallard (Kertész and FánCSI, 2003). Even in lower concentrations, Ni can affect pigment colouring of feathers during moulting (Honda et al., 1990). Zn at higher concentrations can affect reproduction and increase kidney toxicity (Carpenter et al., 2004). Rising Hg levels in birds can affect their breeding success (Gochfeld, 1997), and above 5 ppm adversely affects reproduction (Evers et al., 2007). Co is considered as a significant element necessary for metabolism but can negatively affect it in excessive concentrations (Roginski and Mertz, 1977). The role of metals in benthic organisms is also significant since they are involved in littoral trophic mechanisms and benthic organisms, including fishes, are essential prey for the majority of shorebirds (Wilson, 1989).

* Corresponding authors at: Department of Zoology and Wildlife Biology, AVC College (Autonomous), Mannampandal 609 305, Mayiladuthurai, Tamilnadu, India. (J. Pandiyan); Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India (M. Govindarajan).

E-mail addresses: dunlinpandiyan@gmail.com (J. Pandiyan), drgovind1979@gmail.com (M. Govindarajan).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<https://doi.org/10.1016/j.jksus.2020.08.014>

1018-3647/© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Influences of dietary inclusion of algae *Chaetomorpha aerea* enhanced growth performance, immunity, haematological response and disease resistance of *Labeo rohita* challenged with *Aeromonas hydrophila*

Govindharajan Sattanathan^{a,*}, Thanapal Palanisamy^b, Swaminathan Padmapriya^c,
Vijaya Anand Arumugam^d, Sungkwon Park^e, In Ho Kim^b,
Balamuralikrishnan Balasubramanian^{e,1,*}

^a Department of Life Science, MASS College of Arts and Science, Kumbakonam, 612 501, Tamil Nadu, India

^b Department of Animal Resource and Science, Dankook University, Cheonan -, 31116, South Korea

^c Department of Zoology, Dharmaparam Gnanambigal Government Arts College for Women, Mayiladuthurai, 609 001, Tamil Nadu, India

^d Department of Human Genetics and Molecular Biology, Bharathiar University, Coimbatore, 641 046, Tamil Nadu, India

^e Department of Food Science and Biotechnology, College of Life Science, Sejong University, Seoul-05006, South Korea



ARTICLE INFO

Keywords:

Aeromonas hydrophila
Chaetomorpha aerea
Growth performance
Labeo rohita
Lysozyme activity
Respiratory burst activity

ABSTRACT

The present study was investigated to the sequel of dietary supplementation of algae (*Chaetomorpha aerea*) on growth, exploitation of food, haematological, immunological parameters and disease resistance against *Aeromonas hydrophila* in *Labeo rohita* for 28-days trial period. A total of 180 fishes with a body weight of 55 ± 0.9 g initially, were selected and allotted to one of five dietary treatments, in which the basal diet supplemented with 0, 0.25, 0.5, 0.75, and 1.0 g kg⁻¹ (w/w) of *C. aerea*. The fishes were subjected to challenge with *Aeromonas hydrophila* and relative percentage of survival (%) was recorded for 14 days after the feeding trial. Accordingly, the fishes fed with *C. aerea* increased the growth rate, feed conversion rate, specific growth rate, haemoglobin, neutrophil and lysozyme activity, when compared to the control group ($P < 0.05$). Moreover, *C. aerea* supplementation (0.25 g kg⁻¹) showed that the enhanced body weight, respiratory burst and lysozyme activity and resistance against *A. hydrophila* challenge. On the whole, the obtained results showed that this feeding approach can productively increase the growth and it could be considered as a probable immunostimulant in the organic aquaculture.

1. Introduction

About 15–20% for animal protein intake by humans, provided by the aquaculture and hence the efficient production is the way for ensuring the long lasting as well as efficient industry (FAO, 2010). The major carp in India, *Labeo rohita* is one of the most chosen species in the Asian sub-continent and it includes about 35% of the entire carp manufacture, due to the increased speedy growth along with the delicious taste, acceptable nature towards the feeds provided and the raised value in market (Jhingran and Pullin, 1998; Mishra and Samantaray, 2004; Dey et al., 2005; Rahman et al., 2007; Vutukuru et al., 2007; Kumar et al., 2009). *Aeromonas hydrophila*, has been found to be a predominant pathogenic organism for lower vertebrates, includes fishes as well as amphibians (Vivas et al., 2004; Karunasagar and Rosalind,

1991). The *A. hydrophila* affected fish shows the features like, necrosis, hemorrhagic septicaemia, ulcer formation in skin and exophthalmia that may cause increased death rate, which results in significant drawback in the economy of industries according to the Food and Agriculture Organization that involves the freshwater aquaculture (FAO, 2010; Parker and Shaw, 2011; Shen et al., 2013). The fish may be infected by the pathogens like bacteria due to many factors like intensive farming, pollution in the waters or changes in the external environment (Kumari and Sahoo, 2006). Apart from these factors, the nutritional eminence of the fish may also be a significant factor for the infection due to the inter-relationship with production of energy and the immunity towards the stress caused by the bacterial infection.

Seaweeds are edible and contain vital nutrients such as proteins, essential fatty acids, vitamins, and minerals for human growth. The




* Corresponding authors.

E-mail addresses: sattanathanphd@gmail.com (G. Sattanathan), hmla.mk@sejong.ac.kr, geneticsmurali@gmail.com (B. Balasubramanian).

¹ Equally contributed as first author.

Article

Effect of Green Algae *Chaetomorpha antennina* Extract on Growth, Modulate Immunity, and Defenses against *Edwardsiella tarda* Infection in *Labeo rohita*

Govindharajan Sattanathan ^{1,†}, Vairakannu Tamizhazhagan ², Swaminathan Padmapriya ³, Wen-Chao Liu ^{4,*} and Balamuralikrishnan Balasubramanian ^{5,*}

¹ Department of Life Science, MASS College of Arts and Science, Kumbakonam 612 501, Tamil Nadu, India; sattanathanphd@gmail.com

² Department of Zoology, Syed Ammal Arts and Science College, Ramanathapuram 623 513, Tamil Nadu, India; tamilzoon@gmail.com

³ Department of Zoology, Dharmapuram Gnanambigai Government Arts College for Women, Mayiladuthurai 609 001, Tamil Nadu, India; spsrijan23@gmail.com

⁴ Department of Animal Science, College of Coastal Agriculture Science, Guangdong Ocean University, Zhanjiang 524088, China

⁵ Department of Food Science and Biotechnology, College of Life Science, Sejong University, Seoul 05006, Korea

* Correspondence: liuwc@gdou.edu.cn (W.-C.L.); bala.m.k@sejong.ac.kr or geneticsmurali@gmail.com (B.B.)

† Equally contributed as first author.

Received: 23 September 2020; Accepted: 2 November 2020; Published: 4 November 2020



Simple Summary: Global demand for macroalgal and microalgal foods is growing, and algae are increasingly being consumed for functional benefits beyond the traditional considerations of nutrition and health. The study was undertaken to know the effect of methanolic extract of *Chaetomorpha antennina* in fish. The results demonstrated that the betterment of growth, immune system, and resistance to disease against *Edwardsiella tarda* in Indian major carp, rohu, and *Labeo rohita*. These findings are useful for development of new feed additive in aquaculture sectors.

Abstract: The current study focused on assessing the outcome of methanol extract of *Chaetomorpha antennina* (MECA) on the growth performance and immune modulation in both specific and non-specific immune responses through the assessment of neutrophil, serum lysozyme, serum myeloperoxidase, antiprotease, ceruloplasmin, reactive oxygen species (ROS), and reactive nitrogen species (RNS) activity in *Labeo rohita* (rohu) at 28 days post treatment along with assessment of the disease resistance capacity against *Edwardsiella tarda* at 30 days post immunization. Fishes ($n = 144$; average weight 50.0 ± 0.23 g) were evenly divided into four treatments, with 12 fishes per tank in triplicates. The MECA was injected intraperitoneally in the fishes at different doses as 0, 25, 75, and 150 mg/kg of the body weight. The results demonstrated that fish treated with MECA have an increased body weight, specific growth rate, and feed conversion ratio ($p < 0.05$) with respect to the control group. Results suggested that the MECA inclusion can significantly enhance ($p < 0.05$) the levels of serum lysozyme, neutrophil function, serum antiprotease activity, cellular RNS, and ROS production. Exposure to MECA of 75 mg/kg showed a significantly higher survival percentage against *E. tarda* disease infection. These results indicate MECA as a stimulant of immunity in *L. rohita* against *E. tarda*. The results suggested that MECA is a potent immunostimulant in finfish aquaculture and can offer higher economic welfare.

Keywords: MECA; growth performance; immunostimulant; rohu; challenge test

Efficacy Of Cow Urine Distillate To Enhance The Freshwater Copepod Population Density

Praveena Varadhan, Subhhasri Varadharajan, Priya Muthukirushnan, Venkatalakshmi Sournamanikam

Abstract—Zooplankton has high nutrition value because of some vitamins, amino acids and fatty acids in them. Cow urine is known for its medicinal properties and therapeutic value in india. There are many reports which revealed the application of cow urine in agriculture, poultry, animal health and human health. However, there are no reports for its application in live feed culture. Hence, in the present investigation, the effect of Gir Cow Urine (GCU) was studied on zooplankton for its impact on biomass and biochemical composition. The study revealed that the biomass of zooplankton significantly increased in Gir cow urine distillate treated group. The experiment was carried out one month period. At the end of study period, the biochemical composition of cow urine treated and untreated zooplankton cultures were analyzed.

Key Words—Zooplankton, Gir, Cow urine distillate, Aquaculture, Biochemical analysis, Live feed

1. INTRODUCTION

The word "Plankton" derived from Greek mean drifters and the word was first coined by Victor Hensen in 1887. Plankton commonly falls under two categories namely Phytoplankton and Zooplankton. Phytoplankton are primary producers and minute photosynthetic plants. Zooplankton are secondary producers and animal groups of the freshwater ecosystem. Copepods and Rotifers are the largest, most diversified group in freshwater ecosystem. Copepods are considered as ecological indicators of water quality and global climate change. Alterations in their presence and abundance can be resulted in the low fisheries production. As the major secondary producers of the ecosystem, they fed on phytoplankton and act as food for many major commercially important fishes. For example, a herring stomach contains 60,000 calanoid copepods (1). The studies on copepod ecology in different habitats like estuaries, mangroves, seagrass beds, seaweeds, coral reefs, lagoons and open seas is important to understand the ecosystem health and fish production. Copepod is one of the most abundant live feed organisms which are widely present in almost all the freshwater bodies. Copepods pass through very distinct life stages. They emerge from an egg as a nauplius, usually 100-150µm in length. After six nauplius stages (referred as stages N1 to N6), with growth between each stage, the body shape changes and a series of usually six copepodid stages follow (referred to as stages C1 to C6).

The last of these stages is the adult in which different sexes can be identified. Copepods used as natural food are either cultured or collected from natural water bodies (2) Freshwater finfish seed production often faces a problem of an inadequate food supply. Artificial feeds are widely used, but planktonic animals are very important, especially rotifers, cladocerans, and copepods. Virtually all fish feed on plankton, especially in their early life phases. Planktivorous fish depend on small invertebrates throughout their entire lives. Copepods of the order Cyclopoida are the most important food items in freshwater aquaculture, and their nauplii are especially valuable for feeding fry (3). The importance on mass culture of copepods begins due to the unsatisfactory performance of traditional live feeds *Artemia* and *Rotifer* in larviculture. In mangroves and estuarine environment, harpacticoid copepods are the major food for larvae and juveniles (4). The key problem in culture of copepods is high yield in the sense of hatchery level on a continuous basis. Depletion of DHA (Decosahexanoic acid) and EPA (Eicosapentanoic acid) in the food of the larvae leads to low survival, structural deformities and malpigmentation (5). So that fish and larvae rely mainly on the live feeds which contain high nutrition. The medicinal usage of cow urine is practiced in India from ancient days. Hence cow urine could be expected as a good immunostimulant and water quality enhancer. Cow urine contains various inorganic compounds including silver, Na-K ratio of 4:1 (36%:9% in dried urine), apart from about 3% urea. Fresh cow urine also contains 50-100 mg oestrogens/100 ml; 20- 200 µg of corticosteroids/100 ml and 0.05-0.15 mg of 17-keto-steroids/100 ml (6). Now days, a lot of emphasis has been given on the medicinal use of cow urine in India. Recently the cow urine has been granted U.S. Patents (No. 6,896,907 and 6,410,059) for its medicinal properties, particularly for its use along with antibiotics for the control of bacterial

• Varadhan Praveena, Subhhasri Varadharajan, Priya Muthukirushnan, Venkatalakshmi Sournamanikam

• * Centre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam – 612 001.

• *Corresponding Author E.mail id: dr.s.venkatalakshmi@gcw.ac.in



Original article

Identification of a novel antibacterial protein from hemolymph of freshwater zooplankton *Mesocyclops leuckarti*Varadhan Praveena^a, Sournamanickam Venkatalakshmi^{a,*}, Naiyf S. Alharbi^b, Shine Kadaikunnan^b, Jamal M. Khaled^b, Marimuthu Govindarajan^{c,d,*}^aCentre for Animal Studies, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India^bDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia^cDepartment of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India^dUnit of Natural Products and Nanotechnology, Department of Zoology, Government College for Women (Autonomous), Kumbakonam 612 001, Tamil Nadu, India

ARTICLE INFO

Article history:

Received 29 February 2020

Revised 5 April 2020

Accepted 3 May 2020

Available online 11 May 2020

Keywords:

Mesocyclops leuckarti

Zooplankton

Hemolymph

Antimicrobial proteins

SDS-PAGE

ABSTRACT

Bacterial infections are the most important problem of health care worldwide. The hemolymph antibacterial proteins of *Mesocyclops leuckarti* was isolated for the first time and its antibacterial efficacy was evaluated against four different human pathogenic microbes viz., *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumonia* and *Shigella flexneri*. The antibacterial potential of the antimicrobial proteins of hemolymph samples from plankton cultured in water enriched with Cow Urine Distillate (CUD) was compared with normal ones. The results indicated that the hemolymph proteins were more potential against Gram negative bacteria than Gram positive bacteria. *Klebsiella pneumonia* was more susceptible to the hemolymph proteins exhibiting a zone of inhibition measuring 27 mm. The supplement of CUD to the culture media further enriched the antibacterial activity of the hemolymph proteins (29 mm). The SDS-PAGE analysis indicated two different types of clear bands representing proteins of 53 kDa and 19 kDa. Overall, this investigation signified that the microcrustaceans have a defence mechanism hemolymph of *Mesocyclops leuckarti* have a potential agent for novel antibiotics.

© 2020 The Author(s). Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Copepods represent about 20% of the mean annual zooplankton biomass. They are found in abundance in many aquatic ecosystems (Huys and Boxshall, 1991) and conspicuous member of aquatic biota that have adapted to various habitats such as damp moss, interstitial sand, subterranean localities and parasitic living. The cyclopoid copepods are successful inland water group animals and *Mesocyclops leuckarti* is the important planktonic cyclopoid copepods. In evolution, invertebrates have not developed acquired immune systems. However, they have been bestowed with non-

specific immune mechanisms. Crustaceans show many antibacterial proteins in their hemolymph. However, the literature does not show any record of anti bacterial proteins in microcrustaceans including zooplankton (Iskratsch et al., 2009; Iwanaga, 1993; Kawabata et al., 1995; Mori and Stewart, 1978; Jayasankar and Subramoniam, 1999) (Table 1).

Hemolymph is the type of blood found in Arthropod's open circulatory system. It contains many bioactive molecules which have functional roles in the defence system. The molecules include lectins, complement, clotting factors, antimicrobial peptides (Vazquez et al., 2009).

Among them the antimicrobial peptides are the prime factors that give immunity to the animal. There are two types of antimicrobial peptides identified in hemolymph. They are high molecular weight large antimicrobial proteins (>100 amino acids) and low molecular weight small antimicrobial proteins. The high molecular weight antimicrobial proteins target the disrupt microbial biomolecules and small antimicrobial proteins disrupt the structure and/or the function of microbial cells (Aspan et al. (1995); Stabili et al. (1999); Fujimoto et al. (1995); Hall et al. (1995)). These antimicrobial peptides are secreted in

* Corresponding authors at: Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India (M. Govindarajan).

E-mail address: dr.s.venkatalakshmi@gcwk.ac.in (S. Venkatalakshmi).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

An Analysis of Cropping Pattern and Crop Combination Regions of Thiruvarur District (2011-2016), Tamilnadu, India

S Geetha* and R. Maniyosai#

*Asst. Professor of Geography, Govt. College for Women (A), Kumbakonam

#Asst. Professor of Geography, Govt. Arts College (A), Kumbakonam

(Received 30 September, 2019; accepted 4 January, 2020)

<https://doi.org/10.36224/ijes.120404>

Abstract

India is an Agricultural Nation. Its primary occupation is Agriculture. Today, India ranks second worldwide in farm output. Agriculture and allied sectors like forestry and fisheries accounted for 13.75% of the GDP in 2013, about 50% of the workforce. The crops are generally grown in combination and a particular crop occupies a position of total isolation from other crops. The studies of crop combination regions constitute an important aspect of agricultural geography as it provides a good basis for agricultural regeneration. Through different regions may have different climate and soil conditions. But each region is Individual to crops. For a comprehensive and better understanding of agricultural system the study of crop combination is of great significances and is essential for agricultural planning. The present study enlightens to the crop combinations in Thiruvarur District. Agricultural Geography is very need for rural planning, because high density of population and increasing annual growth rate of population. It is necessary to producer cultivate more crops in the same field. So diversification and suitable combination of crops with short duration is necessary for an increase in the production. The study of crop combination is also helpful for the study of the comprehensive area development planning particularly for the rural areas.

Key words: crop combination, crop diversification, maximum deviation, minimum deviation

1. Introduction

The study of cropping pattern constitutes an important aspect of agricultural geography as it provides a good basis for agricultural regionalization. The crops occupies a position of total isolation other crops in a given area at a given time. The physical factors determine the shape of the areas of crop combination is also helpful for the study of the comprehensive area development planning particularly for the rural areas. The importance of adoption of suitable cropping patterns in a developing country like India cannot be overemphasized. The horizontal expansion of agriculture is not possible without heavy capital investments. Only judicious utilization of land by adopting more remunerative cropping patterns scientific rotation of crops and multiple cropping may help in overcoming the food and raw material problems of the country.

1.1 Aim

The aim of the study is to find out the cropping pattern of Thiruvarur District.

*Corresponding author

Email address: geosgeetha@gmail.com (S Geetha)

ISSN 0976 – 6693. ©2020 SCMR All rights reserved.

ASSESSMENT OF RAINFALL AND RUNOFF ESTIMATION IN SOME PARTS OF THANJAVUR DISTRICT, SOUTH INDIA USING GEO-SPATIAL TECHNIQUES

A. Sumatra¹ and Dr R. Neelakantan²

1. *Research Scholar, Department of Industries and Earth Sciences, Tamil University, Thanjavur, Tamil Nadu, India.*
2. *Professor, Department of Industries and Earth Sciences, Tamil University, Thanjavur, Tamil Nadu, India.*

**Corresponding Author's email: sumanjal@gmail.com*

Abstract

The importance of hydrology in the assessment, development, utilization and management of the water resources of any region is being increasingly realized at all levels. Thanjavur district is one of the worst affected areas during the entire monsoon season. Geographical Information Systems (GIS) is a tool which is used for assessing of Rainfall and Runoff Estimation in some parts of Thanjavur District.

1. INTRODUCTION

Hydrology is the science that deals with the occurrence and distribution of the water of the earth in time and space, both above and below the land surface, including their chemical, biological and physical properties, and their interaction with the physical environment (WMO, 1992). The importance of hydrology in the assessment, development, utilization and management of the water resources of any region is being increasingly realized at all levels. The main scope of hydrology is to study (a) the maximum probable flood that may occur at a given site and its frequency; this is required for the safe design of drains, culverts, dams, reservoirs, channels and other flood control structures, (b) the water yield from an area, its occurrence, quantity and frequency are necessary for the design of dams, municipal water supply, water power and river navigation, (c) the groundwater development for which a knowledge of the hydrogeology of the area, i.e., the formation soil, recharge facilities like streams and reservoirs, rainfall pattern, climate, cropping pattern are required and (d) the maximum intensity of storm and its frequency are required for the design of a drainage project in an area.

2. THE STUDY AREA

Thanjavur district lies between 9° 50' and 11° 25' North latitude and 78° 45' and 79° 25' East longitude (Fig.1). It is bounded on the North by Thiruchirapalli and Cuddalore districts, on the East by Tiruvarur and Nagapattinam districts, on the South by Palk Strait and Pudukottai district and on the west by Pudukkottai district and Tiruchirapalli districts. Total geographical of the study area is 1697 sq.km. The mean maximum temperature was 37.48°C during May - July. Similarly, the mean minimum temperature was 20.82°C during November-January. The north east monsoon provides much rainfall with 545.7 mm and 953.2 as normal and actual rainfall respectively, while southwest monsoon provides 342 and 303.1 mm as normal and actual rainfall respectively. The total population of Thanjavur district is 22,16,138.

A Survey on Load Balancing Techniques in Cloud Computing Servers

E.Suganthi ¹ **Dr.F.KurusMalaiSelvi²**

Research Scholar,PG&Research Department of Computer Science ,Bharathidasan University,India

E-mail:esuganthics@gmail.com

Associate Professor & Head ,PG&Research Department of Computer Science ,Bharathidasan University,India

E-mail:fkmsjoe@gmail.com

Abstract:

Cloud Computing is an recent landmark development in the area of computing. It offers huge power to next generation computing.. Dynamic provisioning forms the base for cloud computing and facilitates and supports the network services. Its primary focus is on making the vision of utility computing a reality, with an additional “pay-as-you-go” feature. It generates immense potential to in development of applications and products that specialize in greater resource utilization and scalability. This paper presents the survey on basic cloud computing fundamentals and therefore the concepts of load balancing i.e. Scheduling of load within the cloud. It elaborates the prevailing load scheduling algorithms with their merits/demerits and suitability within the cloud and heterogeneous computing environment and proposes a replacement perspective for better results as per desired parameters.

Keywords: Load balancing, Cloud, scheduling, response time, scheduling Algorithms.

1.INTRODUCTION

Load balancing ensures that workload is equally distributed across all processors, in order to ensure that no processor is overloaded. A load balancer may be a physical device, running on a specialized hardware or software process, and it accepts multiple requests from users and distributes them evenly across servers. [20] Load balancing increases throughput and thereby reduces reaction time. Load balancing in cloud, distributes the surplus or excess dynamic local workload evenly across all the nodes. It ensures that there is better service provisioning and a near ideal resource utilization ratio is achieved. It is effective in improving improves the general performance of the system. Incoming tasks which are received from different location are received by the load balancer and is then distributed to the data center for improved and proper load distribution.

Geospatial Technology for Assessment of Soil Erosion Probability Zone and Sediment Yield Estimation in Parts of Thanjavur District, South India

Sumatra A^{1*} and Neelakantan R²

1. Research Scholar, Department of Industries and Earth Sciences, Tamil University, Thanjavur, Tamil Nadu, India.
2. Professor, Department of Industries and Earth Sciences, Tamil University, Thanjavur, Tamil Nadu, India.

*Corresponding Author Email - sumiangal@gmail.com

Abstract

Soil erosion level assessment is essential for coastal and delta areas. In the present study, soil erosion probability zone and sediment yield assessment were carried out in Thanjavur district, Tamil Nadu, India using Geographical Information System (GIS). The main topography parameters considered in soil erosion mapping include slope, geography, land use / land area, drainage density and elevation. These parameters are integrated with ArcGIS under various thematic maps. A probability weighting approach was adopted that allowed a linear combination of the probability weights of each thematic map (Wt) and a ranking was assigned to different features in the thematic maps and final scores (Sc) were obtained for their role in soil erosion. The Integrated Soil Erosion Map is classified into five zones as very low, low, moderate, high and high erosion-prone zones. Landslides associated with moderate slopes, high altitudes, high drainage density, flood plains and agricultural land are more prone to erosion. Drainage, drainage density, average annual rainfall, temperature and flow, vegetation cover factor (FC) and slope maps were used to estimate sediment yield. Sediment yield from the district is low and not in a very dangerous condition. The results revealed that soil conservation measures in this area may be considered for sustainable development.

Keywords: Soil Erosion, Sediment Yield, Remote Sensing, GIS, Thanjavur District.

1. INTRODUCTION

Soil erosion is a serious problem worldwide due to its adverse economic and environmental impacts (Ande et al., 2009). Soil erosion probability mapping is one of the most important requirements for planning management, and conservation (Cleaver and Schreiber 1995). Quantitative data on soil erosion rates in the nationwide area are essential for the development of soil conservation and management plans and for the assessment of environmental implications (Alexakis et al., 2013). Different onsite effects of soil erosion like loss of topsoil, change of soil structure, loss of soil organic matter content which leads to reduction of productivity and on the other hand offsite events like reduction of channel depth, water holding, water discharge and transport capacity of a stream caused increase of flood intensity and frequency (Pimentel 2006; Sinha and Joshi 2012; Zhou and Wu 2008). Over the past decades many studies and empirical models have been developed by many researchers to estimate soil erosion in different parts of the world (Morgan et al. 1998; Kim and Julien 2006; Darbral et al., 2008; Melesse et al., 2011).

In hilly and steep sloppy regions, soil erosion may initiate landslides as a short-term and degradation of soil quality as a long-term effect (Michael and Samanta 2016). The erosion in upper catchment areas depends on amount of rainfall, vegetation cover, soil characteristics, slope and is accelerated by human activity like mining, agriculture and deforestation (Pal et al., 2012). Arekhi et al., (2012) have investigated the spatial distribution of annual soil loss and sediment yield in Cham Gardalan watershed, Ilam Province, Iran, using Revised Universal Soil Loss Equation (RUSLE) model. Yildirim (2011) conducted research to apply Universal Soil Loss Equation (USLE) with



ISSN: 0975-833X

Available online at <http://www.ijer.com>

INTERNATIONAL JOURNAL
OF CURRENT RESEARCH

International Journal of Current Research
Vol. 12, Issue, 08, pp.13129-13131, August, 2020

DOI: <https://doi.org/10.24941/ijer.39290.08.2020>

RESEARCH ARTICLE

GEOMATIC BASED SOIL AND GEOMORPHOLOGICAL CLASSIFICATION OF KUMBakonam TOWN, THANJAVUR DISTRICT, TAMILNADU

¹Kalyani, V. and ²Neelakandan, R.

¹Department of Geography, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India

²Department of Industries & Earth Sciences, Tamil University, Thanjavur- 613010, Tamil Nadu, India

ARTICLE INFO

Article History:

Received 15th May, 2020

Received in revised form

21st June, 2020

Accepted 24th July, 2020

Published online 30th August, 2020

Key Words:

Soil, Geomorphology, GIS Technology.

ABSTRACT

In the present study, an attempt has been made to classify the different soil and geomorphic units of Kumbakonam town Thanjavur district, Tamilnadu. The drainage, contour and delineated geological units have been overlaid on IRS-ID LISS III satellite imagery (bands 2, 3 and 4) the system to delineate and characterize different geomorphological units and analysis of their processes based on the field observations. The study area is basically of sedimentary in origin with different geomorphological formations and is influenced by the various processes. Based on the satellite data analysis, the distinct geomorphological units viz., alluvial plain and deltaic plain fills have been delineated and characterized. The information generated from satellite data in the form of vector layers has been used in GIS to generate soil and geomorphological maps of the study area. The present study demonstrates that IRS-ID LISS-III data in conjunction with geomorphology, soil, river and drainage and parameters to enable detailed evaluation of different geomorphological units and analysis of their processes based on the field observations. The delineated geomorphological units can be utilized for evaluation and management of natural resources and geo-environment on sustainable basis at river catchment level.

Copyright © 2020, Kalyani and Neelakandan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Kalyani, V and Neelakandan, R. 2020, "Geomatic Based Soil and Geomorphological Classification of Kumbakonam Town, Thanjavur District, Tamilnadu", International Journal of Current Research, 12, (08), 13129-13131.

INTRODUCTION

The soil and geomorphological pattern using Remote Sensing has been attempted in a number of previous investigations. Roy and Raina (1973) studied hydrogeomorphology of Kotepalli Catchment area of Hyderabad. Chatarjee et al., (1978) studied the geomorphology of central Luni Basin of Western Rajasthan. Raju and Vaidyanadhan (1984) also used Remote Sensing techniques in the study of Sarada River Basin. Similar investigations were also attempted by some more workers (Raviprakash and Mishra 1993, Mangrulkar et al., 1993; Thomas et al., 1995, Saini and Nathawat, 1996). Recently Jaisankar et al., (2001) have undertaken hydromorphological and Remote Sensing studies for groundwater exploration in Agnigundala area, Andhra Pradesh. In this investigation, geomorphology and land use pattern of Visakhapatnam urban industrial area has been taken up using IRS-IB and SPOT imageries with subsequent field checks.

Study Area: Kumbakonam is a special grade Municipal Town and second biggest town in terms of administrative status to Thanjavur District. It is situated 10°57' North Latitudes and 79°23' East Longitude Kumbakonam is located 313 km away from Chennai on the South, 90 km away from Trichy on the East and 40 Km away from Thanjavur on the North-East. The town is bound by River "Cauvery" on the North and "Aresalar" on the South with a gentle slope from north to south. Civil Administration was looked by a town level committee (Municipal Committee) which was formed in 1866 with an extent of 7.68 Sq.km. At present, the Kumbakonam town extends over an area of 12.58 Sq.km and the local planning area extends over an area of 64.02 Sq.Km.

Soil: The town is not having any commercially exploitable minerals. The Alluvium and Regur are the major type of soils found in the region. More than 65% of the areas are covered by alluvium soil due to the presence of Cauvery delta. This is fertile and suited for agricultural purpose. Rice is the principle crop: Coconut, Mango and Tamarind are the major types of trees found in the region. The Adanur soil type mostly found in the western part and the Alattur soil type found in the eastern part of the study area. The Padugai soil type mostly found in the northern part along the edges.

*Corresponding author: Kalyani, V.,

Department of Geography, Government College for Women (Autonomous), Kumbakonam 612001, Tamil Nadu, India.

PHYTOCHEMICAL CHARACTERIZATION OF *Naringi crenulata* (Roxb) LEAF EXTRACT BY GC- MS method

B. Bhavani and S. Vasantha

*Department of Botany, Government College for woman (Autonomous), Kumbakonam,
(Affiliated to Bharathidasan University), Thanjavur, Tamil Nadu, India.*

*P.G and Research Dept. of Botany and Microbiology, A.V.V.M Sri Pushpam College (Auto),
(Affiliated to Bharathidasan University), Poondi, Thanjavur, TamilNadu, India.*

ABSTRACT

The bioactive compounds of *Naringi crenulata* (Roxb) leaves have been evaluated using GC-MS. The active chemical compounds were identified by using Gas Chromatography-Mass Spectrometry. The phytochemical compounds found in the *N. crenulata* leaves with methanolic extract were matched with NIST library. GC-MS analysis of leaves extract *Naringi crenulata* in characterization of photochemical sixteen components were identified. Phytochemical compounds of *N. crenulata* leaves with methanolic extract was 16 specified compounds identified including hexadecocenoic acid performed.

Keywords: Methanol extract, *Naringi crenulata*, GC-MS analysis and photochemical.

INTRODUCTION

Herbal medicine is one of the most remarkable uses of plant based diversity. As many as 75 to 90% of the world's rural people rely on herbal medicine. The success of any health care system depends on the availability of suitable drugs on a sustainable basis. Natural medicine improves strength of the body. The knowledge medicinal plant has been accumulated in the course of many centuries based on different medicinal systems such as Ayurveda, Unani and Siddha. In india, it is reported that traditional healers uses 2500 plants species 100 species

Phytochemical analysis and Biological activities of fruit extract of *Naringi crenulata* (Roxb) Nocols.

Bhavani B and Vasantha S

*P.G and Research Department of Botany and Microbiology
A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur (D.t)
Tamil Nadu, India.
[Email:bhavani1493@gmail.com](mailto:bhavani1493@gmail.com)
Mobile:919843662674*

ABSTRACT

*This study was carried out with an objective to investigate the phytochemical analysis, antimicrobial, antidermatophyte and antihelminthic activity potentials of fruit of *Naringi crenulata* (Roxb).The phytochemical screening and antimicrobial activities with different solvent aqueous, methanol, chloroform and benzene extract and antidermatophytes, antihelminthic activity aqueous and methanol fruit extract treated against microbes and earthworms. The results showed antibacterial activity of maximum zone of inhibition of *Enterococcus aeromonas* with aqueous extract and antifungal activity maximum zone of inhibition *Penicillium sp* in aqueous fruit extract respectively. The antidermatophyte activity maximum zone of inhibition of *T. rubrum* activity fruits extract and dose dependent activity was observed in aqueous and methanol leaf extracts exhibit, respectively dose-dependent action and death mortality (%) compared to others. The present study qualitative and quantitative phytochemical confirm the presence of alkaloid, carbohydrate, flavonoids, glycosides, phlobatannins, protein, saponin, steroid, tannin, terpenoids and teriterpenoids compound in fruit extract. The presence of phytochemicals, may be of use for developing plant based drugs for various ailments.*

Keywords: antifungal activity, secondary metabolites, *Naringi crenulata*, antibacterial activity.



Structural, spectral and nonlinear optical analysis of Sodium bis(2-methylactato)borate: a new semiorganic crystal for photonic applications

G. Gokila¹ · R. Aarthi¹ · C. Ramachandra Raja¹

Received: 15 December 2019 / Accepted: 6 October 2020
© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

Semiorganic single crystals of Sodium bis(2-methylactato)borate (NaMB) have been grown using solvent evaporation process. NaMB crystal comprises a bis(2-methylactato) borate anion and a sodium cation. Powder XRD pattern reveals the good crystallinity. Linear optical transmittance window is found to be 215–1100 nm. Na···O–C intermolecular interaction is observed in the grown crystal and it is established by observed vibrational frequencies. Functional group associated with borate is well addressed through the vibrations observed in FTIR and FT-Raman spectroscopy. ¹³C NMR spectroscopy enunciates the position of carbon atoms in the crystal structure. Intermolecular interactions such as Na1···O6–C5 and Na1···O1–C1 where O6 and O1 are uncoordinated oxygen atoms of carboxylato groups which pulls the electrons from C5 and C1 and causes deshielding effect resulting in shifting of C5 and C1 signals towards higher ppm. These interactions are also established through the shifts observed in ¹H NMR spectroscopy. The third order nonlinear susceptibility (χ^3) value is 1.07911×10^{-6} esu and this larger value is due to electron delocalization in carboxylato group and charge transfer due to intermolecular interaction present in the crystal structure. The self-focussing and saturable absorption nature of the crystal may be utilized in optical sensors and Q-switching applications.

Keywords Crystal structure · Uncoordinated oxygen atoms · NMR spectroscopy · Z-scan technique

1 Introduction

In recent years, there has been significant interest in the study of nonlinear optical (NLO) crystals with better properties because of their wide applications in the areas of laser technology, optical communication, optical information processing and optical data storage technology colors displays, electro optic switches, frequency conversion etc., (Chenlin et al. 2002; Arivuselvi and Ruban Kumar 2016). The majority of the organic crystals

✉ C. Ramachandra Raja
crrajaphy@gmail.com

¹ Government Arts College (Autonomous), Affiliated to Bharathidasan University, Kumbakonam, Tamilnadu 612002, India



Contents lists available at ScienceDirect

Optik

journal homepage: www.elsevier.com/locate/ijleo

Original research article

Interpretation of structural, spectral and nonlinear optical properties of a new semiorganic crystal: Rubidium bis(2-methylactato) borate monohydrate



Gokila G, Aarthi R, Ramachandra Raja C*

Government Arts College (Autonomous), Kumbakonam 612002, Tamilnadu, India

ARTICLE INFO

Keywords:

Crystal growth
Crystal structure
NMR spectroscopy
Z-scan analysis

ABSTRACT

Rubidium bis (2-methylactato) borate monohydrate (RbMB) was developed by adopting slow solvent evaporation technique. Good crystalline nature of RbMB is confirmed by powder X-ray diffraction analysis. The optical transparency window is observed from 190 nm to 1100 nm. FTIR and FT-Raman spectral analysis endorses the formation of RbMB crystal. Thermal stability of RbMB extends upto 134 °C. Molecular structure of RbMB crystal is interpreted through the observed chemical shifts through ^1H NMR and ^{13}C NMR spectroscopy. The intermolecular interactions between organic and inorganic moieties are well established. These intermolecular interactions and electron delocalization enhances the third order nonlinear susceptibility (χ^3) value and is found as 3.75354×10^{-6} esu. The self-defocusing and reverse saturable absorption nature gives negative non-linear refractive index (n_2) and positive nonlinear absorption coefficient (β). This made RbMB a suitable candidate for using in Q-switching, optical pulse shorteners and optical energy limiters applications.

1. Introduction

The fast developments in the field of optoelectronics and photonics creates a demand in developing new and better nonlinear optical (NLO) crystals. These crystals finds greater applications in various fields [1–5]. The structure property relationship is the most important factor to develop an efficient NLO crystal. Combining the organic materials nonlinear optical efficiency and inorganic materials physical stability by choosing a proper precursor results in the formation of favourable semiorganic NLO crystal. Based on this view, authors made an attempt to develop 2-methylactic acid based crystals. Four new 2-methylactic acid based crystals were developed and the solved structures were reported by the authors [6–9]. Among the structures reported, Rubidium bis(2-methylactato)borate monohydrate crystal is characterized using various characterization techniques and the structure-property relationship is reported in this article. The crystal structure consists of one 2-methylactato borate anion, one rubidium cation and a water molecule. O–Rb...O and O–H...O hydrogen bonded intermolecular interactions present in the RbMB structure ensures enhanced (χ^3).

2. Materials and methods

RbMB was prepared by reacting 2-methylactic acid, boric acid and rubidium carbonate in 4:2:1 molar ratio by using deionized water as solvent. The mixture is stirred well and filtered to remove impurities. The filtered solution is kept closed in a beaker using

* Corresponding author.

E-mail address: crrojaphy@gmail.com (R.R. C).<https://doi.org/10.1016/j.ijleo.2020.164324>

Received 19 December 2019; Accepted 27 January 2020

0030-4026/ © 2020 Elsevier GmbH. All rights reserved.



Structural, Spectral, and Optical Characterization of Potassium bis(2-methylactato)borate Hemihydrate Crystal

G. GOKILA,¹ R. AARTHI,¹ and C. RAMACHANDRA RAJA^{1,2}

1.—Government Arts College (Autonomous), Kumbakonam, Tamilnadu 612002, India. 2.—e-mail: crrajaphy@gmail.com

Potassium bis(2-methylactato)borate hemihydrate (KMB) was grown by a solvent evaporation technique. Crystal structural analysis revealed that the potassium cation was pseudo-octahedrally coordinated by five O atoms from four bis(2-methylactato)borate (MB) ligands and half-occupied water. The sharp peaks in the powder x-ray diffraction (PXR) pattern confirmed the perfect crystalline nature of the KMB. Transparency was observed in the range from 215 nm to 1100 nm, and the bandgap of KMB was determined to be 2.15 eV. The molecular structure of KMB was established by interpreting the functional group vibrations through vibrational spectroscopy. The molecular structure was further confirmed by the nuclear magnetic resonance (NMR) spectral technique. The peak followed by valley nature observed is due to the self-focusing behavior of KMB. The minimum transmission near the focus in the open aperture curve is attributed to the reverse saturation absorption nature of KMB. The third-order susceptibility ($\chi^{(3)}$) of KMB is estimated to be 4.17694×10^{-5} esu. These results suggest that KMB can be used in optical sensors as well as other photonic and optoelectronic applications.

Key words: Crystal structure, solvent evaporation technique, NMR analysis, Z-scan technique

INTRODUCTION

Progress in the area of technology and communications has resulted in greater demand for the development of suitable nonlinear optical (NLO) materials.¹ The generation of different frequencies results in different laser sources, in turn enabling advances in the fields of photonics, optoelectronics, and optical communications.^{2–5} To achieve these advances, current research is focusing on the development of efficient NLO crystals. Moreover, many researchers have developed and reported on several organic and inorganic NLO crystals. Organic crystals such as 4-*N,N*-dimethylamino-4'-*N'*-methyl-stilbazolium tosylate (DAST)⁶ and 3-methyl-4-methoxy-4'-nitrostilbene (MMONS)⁷ exhibit high

NLO efficiency but lack mechanical and thermal stability. Meanwhile, inorganic crystals such as $\text{KBe}_2\text{BO}_3\text{F}_2$ (KBBF),⁸ $\text{K}_3\text{B}_6\text{O}_{10}\text{Cl}$ (KBOC),⁹ $\text{Ba}_4\text{B}_{11}\text{O}_{20}\text{F}$ (BBOF),¹⁰ etc. exhibit good physicochemical stability and deep-ultraviolet (UV) generation but poor NLO efficiency. To overcome these shortcomings of organic and inorganic crystals, semiorganic crystals have emerged over the past few years. The functionalization of organic materials with suitable inorganic acids and salts leads to the formation of new semiorganic crystal structures with hydrogen bonding.^{11,12} The formation of hydrogen bonds in the structure promotes the NLO efficiency of such crystal as well as their physical properties.^{13,14} Our research group focuses on the development of new semiorganic-based NLO materials, e.g., using 2-methylactic acid. Among the various organic moieties, 2-methylactic acid is chosen due to its coordinating ability with aqueous media.¹⁴ Cammas et al. established that, when

(Received February 11, 2020; accepted July 23, 2020)



Elucidation of the properties of Lithium bis(2-methylactato) borate monohydrate crystal for laser applications

G. Gokila¹ · R. Aarthi¹ · C. Ramachandra Raja¹

Received: 16 December 2019 / Accepted: 17 March 2020
© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

Lithium bis(2-methylactato) borate monohydrate (LiMB) a semiorganic crystal is grown by solvent evaporation process. Sharp peaks present in powder XRD (PXR) pattern reveals the good crystallinity. Linear transmission window is observed from 220 to 1100 nm. Functional groups such as CH₃, OH and B–O present in the crystal structure are established through FTIR and FT-Raman vibrational analyses. ¹H NMR and ¹³C NMR spectroscopy confirms molecular structure of LiMB crystal. The organic part of LiMB is elucidated through the observed chemical shifts. The influence of intermolecular interactions was well identified and established. Thermal stability of LiMB extends upto 218 °C. The presence of electron delocalization and intermolecular interaction enhances the third-order susceptibility (χ^3) and it is 4.5517×10^{-5} esu. Nonlinear refractive index (n_2) and nonlinear absorption coefficient (β) are also found. The self-defocussing and reverse saturable absorption behavior of grown crystal made it useful for Q-switching, optical pulse shorteners and optical energy limiters applications.

1 Introduction

Nowadays materials research society focusses on developing semiorganic nonlinear optical (NLO) crystals for electronics and photonics applications because of their ability to combine the salient features of organic and inorganic moieties [1–3]. Nonlinear optical crystals such as L-Valium picrate, Triphenylmethane DAST and MMONS gives good NLO efficiency and can be utilized in the field of logic devices, biological imaging and optical switching [4, 5]. Semiorganic NLO crystals find more significance in the field of laser technology. The demand for lasers with various wavelength is increasing in the industries, medical field, communication, defence etc. [6–8]. Developing laser with various wavelength can be achieved through developing a good NLO crystal. Centrosymmetric crystal grown in polymeric structure with several intermolecular interactions gives better third-order NLO parameters [9]. Also, the centrosymmetric third-order nonlinear optical crystals gain more importance in the field of optical bistability, ultra fast optical switching, and optoelectronic devices [10–13]. The NLO crystals with negative third-order nonlinear refractive index and positive nonlinear

absorption coefficient finds applications in the field of optical sensors and Q-switching applications. Joshua et al. synthesized a centrosymmetric Lithium bis(2-methylactato) borate monohydrate, reported its structure [14] and is given in Fig. 1. In this article, the molecular structural confirmation of LiMB crystal through vibrational and NMR spectroscopies is reported. Also, the thermal stability, linear optical transmittance window and third-order nonlinear efficiency of LiMB crystal are reported. The reported O—Li—O intermolecular interactions were well acknowledged through the various characterization techniques and also it is responsible for the enhanced χ^3 value.

2 Experimental

Slow solvent evaporation method is utilized to grow LiMB crystal and was grown by reacting 2-methylactic acid, boric acid and lithium carbonate in 4:2:1 molar ratio. 4.1644 g of 2-methylactic acid, 1.236 g of boric acid and 0.7389 g of lithium carbonate were appropriately dissolved in 10 ml, 50 ml and 10 ml of deionized water and these solutions were mixed together using magnetic stirrer until to get the clear solution. The clear solution was filtered in a beaker and covered using a perforated paper. Then the beaker was left to stand at a room temperature. Slow evaporation of the solvent leads to growth of crystals in one month.

✉ C. Ramachandra Raja
crrajaphy@gmail.com

¹ Government Arts College (Autonomous), Kumbakonam,
Tamil Nadu 612002, India

Soft Multi Pre Compact and Soft Multi Pre Lindelof Spaces via Soft Multi Pre open Sets

R. GOWRI¹ and **G. SAHITHYABHARATHI²**

¹ Assistant Professor, Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Trichirappalli
E-mail gowrigck@rediffmail.com

² Research Scholar, Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Trichirappalli
E-mail gsahithyabharathi@gmail.com

Abstract

The purpose of this paper is to study the concept of topological structure formed by soft multi sets. Also, the notion of four sorts of generalized soft multi compact spaces namely, soft multi pre compact, soft multi pre Lindelof, almost soft multi pre compact, almost soft pre Lindelof spaces via soft multi pre open sets are to be introduced and their properties are analysed.

AMS Subject Classification: 54A40, 03E70

Keywords: soft multi topological spaces, soft multi pre open sets, Soft multi pre compact spaces, soft multi pre Lindelof spaces, almost soft multi pre compact spaces, almost soft multi pre Lindelof spaces.

1 Introduction

In 2012, Zorlutuna et al [13] first studied the compactness for soft topological spaces. In 2011, Alkhazaleh et al. [1] as a generalization of Molodtsov's soft set we introduce the definition of a soft multi set, its basic operations such as complement, union and intersection etc. Soft pre open sets in soft topological spaces were introduced by Gananambal Ilango and Mirudula Ravindran [6] in the year of 2013. In 2018 Al-Shami and El-Shafei [3] introduced and studied the concept of soft compact and soft Lindelof spaces via soft pre open sets. In 2019, Gowri and Sahithyabharathi introduced the concept of soft multi b-compact spaces by using the concept of soft b-open sets and its properties.

The purpose of this study is to employ soft multi pre open sets to define the concepts of four kinds of generalized soft multi compact spaces namely, soft multi pre compact, soft multi pre Lindelof, almost soft multi pre compact and almost soft multi pre Lindelof spaces, we provide various examples to elucidate the relationships among these spaces and to point out some properties of them. Moreover, we offer some soft topological concepts such as soft multi pre hyperconnected and pre partition spaces and discuss some properties which connect them with introduced soft multi spaces. The sufficient conditions for the four initiated soft multi spaces to be soft multi hereditary properties are investigated.

Applications of Soft Multi Topological Spaces in Multi Criteria Decision Making

R. GOWRI¹ and **G. SAHITHYABHARATHI²**

¹ Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Trichirappalli.
E-mail gowrigck@rediffmail.com

² Research Scholar, Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Trichirappalli.
E-mail gsahithyabharathi@gmail.com

Abstract

In this paper we use Analytic Hierarchy Process (AHP) to conclude that the Soft Multi Sets (SMS) of Soft Multi Topological spaces (SMTS) gives better results in Multi Criteria Decision Making.

AMS Subject Classification: 54A40, 03E70

Keywords: soft multi sets, soft multi topological spaces, Decision Making, Multi Criteria Decision Making, Analytic Hierarchy Process.

1 Introduction

To deal with uncertainties, many theories have been recently developed, including the theory of probability, the theory of fuzzy sets, the theory of rough sets and so on. However, difficulties are still arising due to the inadequacy of parameters. In 1999, Molodtsov [6] introduced the concept of soft set theory as a general mathematical tool for dealing with uncertainty. The solution of such problems involve the use of mathematical principles based on uncertainty and imprecision.

In the definition of Molodtsov the soft set is a mapping from a set of parameters to a power set of the universe. This means, we have one set of parameters to the power set of one universe, so the user can make a decision where an element in the universe having a property. But all the problems does not consists of one universe. To solve the problem with multi set of universes and as a generalization of Molodtsov's soft sets we introduce the definition of a soft multi set, its basic operations such as complement, union and intersection are introduced by Alzhazaleh et. al. [1] in 2011.

In 2011, Shabir and Naz[11] introduced and studied the concept of soft topological spaces and related concepts. In 2013, Anjan Mukherjee et. al. [2] introduced the concept of topological structure formed by soft multi sets and soft multi compact spaces.

Multi criteria decision making (MCDM) plays a critical role in many real life problems. It is not an exaggeration to argue that almost any local or federal government, industry, or business activity involves, in one way or the other the evaluation of a set of alternatives interms of a set of decision criteria.[12]



Interplay of three-body and higher-order interactions on the modulational instability of Bose–Einstein condensate

SABARI SUBRAMANIYAN,^{1,2,6} OLIVIER TIOKENG LEKEUFACK^{3,4,*} ID RAMASWAMY RADHA,^{1,7}
AND TIMOLEON CREPIN KOFANE^{5,8}

¹Centre for Nonlinear Science, Department of Physics, Government College for Women, Kumbakonam 612001, India

²Department of Physics, Bharathidasan University, Tiruchirappalli 620024, India

³Pure Physics Laboratory: Group of nonlinear physics and complex systems, Department of Physics, Faculty of Science, University of Douala, P.O. Box 24157, Douala, Cameroon

⁴Centre d'Excellence Africain des Technologies de l'Information et de la Communication (CETIC), Université de Yaoundé I, Yaoundé, Cameroon

⁵Department of Physics, Faculty of Science, University of Yaoundé I, P.O. Box 812, Yaoundé, Cameroon

⁶e-mail: ssabari01@gmail.com

⁷e-mail: vittal.cnls@gmail.com

⁸e-mail: tckofane@yahoo.com

*Corresponding author: lekeufackolivier@gmail.com

Received 20 April 2020; revised 9 June 2020; accepted 9 June 2020; posted 10 June 2020 (Doc. ID 395007); published 7 July 2020

We investigate the modulational instability (MI) of trapped Bose–Einstein condensates with three-body and higher-order interactions employing both semi-analytical and numerical methods. Using the time-dependent variational approach, we derive variational equations for the time evolution of the amplitude, phase of modulational perturbation, and effective potential of the system. By means of an effective potential, we retrieve the corresponding MI condition of the dynamical system under consideration. The interplay between three-body interaction and higher-order interaction is discussed in detail. The semi-analytical predictions are confirmed through numerical simulations. © 2020 Optical Society of America

<https://doi.org/10.1364/JOSAB.395007>

1. INTRODUCTION

In the ultracold regime where the temperature is much lower than the critical temperature for condensation, a Bose gas may obey the $T = 0$ formalism. Most experiments in Bose–Einstein condensates (BECs) are reproduced and described by the theoretical model based on the nonlinear mean-field Gross–Pitaevskii (GP) equation with two-body interaction. The effect of the interatomic interaction leads to a nonlinear term in the GP equation, which is proportional to the s -wave scattering length a_s and the condensate density [1]. One can change the sign and strength of the scattering length by using the Feshbach resonance technique [2]. This indicates that the strength of the interaction can be restrained by using diverse experimental devices. In the one-dimensional (1D) homogeneous limit, the GP equation takes the form of a nonlinear Schrödinger equation that supports a spectrum of exact soliton solutions. Experimentalists approach this mathematically ideal scenario by confining the condensate in an elongated and prolate trap, typically with tight radial confinement. However, this quasi-1D geometry is usually accompanied by the presence of weak axial

harmonic trapping, which removes the integrability of the system [3].

It is understood that at low densities, where the interatomic distances are significantly greater than the distance scale of atom–atom interactions, the two-body interaction can be described by a scattering length where the effects of the higher-order interactions are negligible [1,4]. But in certain experiments, the density of the BECs is considerably high. In particular, the evolution of BECs on the surface of atomic chips and in atomic waveguides involves a strong compression of the traps, and this results in the enhancement of densities in BECs [5–8]. Consequently, the simple GP equation (with two-body interaction alone) becomes inadequate. Hence, the dynamics of the BEC needs a better description of atom–atom interaction. Such a system comprises three-body interactions due to higher densities and shape-dependent potential due to strong confinement. Higher-order terms in the expansion of the phase shifts at low momenta, determined by the effective range, the shape parameter, etc., give corrections to the simple GP equation. It has been shown that the critical number of condensed atoms needed for stability, the chemical potential, the condensate



Dynamics of solitons in the fourth-order nonlocal nonlinear Schrödinger equation

T. A. Gadzhimuradov · A. M. Agalarov ·
R. Radha · B. Tamil Arasan

Received: 12 April 2019 / Accepted: 5 November 2019 / Published online: 14 November 2019
© Springer Nature B.V. 2019

Abstract We consider the fourth-order nonlocal nonlinear Schrödinger equation and generate the Lax pair. We then employ Darboux transformation to generate dark and antidark soliton solutions. The highlight of the results is that one ends up generating a two-soliton solution characterized by one spectral parameter alone, a property which has never been witnessed so far.

Keywords Nonlocal nonlinear equations · Lax pair · Darboux transformation · \mathcal{PT} symmetry · Soliton

1 Introduction

The advent of femtosecond pulse propagation through optical fibers [1] prompted the necessity to go beyond the classical integrable nonlinear Schrödinger (NLS) equation [2,3] to study higher-order effects on ultra short pulse transmission. In fact, this resulted in the investigation of several higher-order integrable NLS

equations like Hirota equation [4], quintic NLS equation [5], etc. However, all these integrable equations are local equations, i.e., the evolution of the solution depends on local space and time.

On the other hand, optical systems exhibiting parity-time (\mathcal{PT}) symmetry have gained the attention of the researchers in the last few years [6–10]. The optical beam dynamics in such systems is described by the Schrödinger equation with a complex potential which can be realized through a judicious inclusion of refractive index and gain/loss regime. If the dependence of the refractive index on electric field intensity is introduced [11,12], then the dynamical system can be mathematically modeled by the celebrated \mathcal{PT} symmetric NLS equation [13]

$$i\partial_t q(x, t) + \partial_x^2 q(x, t) + 2\alpha q(x, t)q(x, t)\bar{q}(-x, t) = 0 \quad (1)$$

In the above equation, the potential is both \mathcal{PT} symmetric and nonlinear simultaneously. In a single word, the above equation does not have a direct application in optics; it can describe prospective meta-magnetic materials by the virtue of its gauge equivalent magnetic spin systems [14]. In fact, rich dynamics exhibited by Eq. (1) [15–19] has proved to be a catalyst to open new vistas in nonlocal integrable equations. In addition, the recent investigation of several \mathcal{PT} symmetric nonlinear partial differential equations (pdes) [20–22] in the field of nonlinear optics, hydrodynamics, and atmospheric sciences has underscored the emphasis to penetrate deep into the domain of nonlocal \mathcal{PT} symmetric

T. A. Gadzhimuradov (✉) · A. M. Agalarov
Department of Theoretical Physics, Institute of Physics
Dagestan Scientific Centre, Russian Academy of Science,
Makhachkala 367003, Russia
e-mail: gta-1987@mail.ru

R. Radha
Centre for Nonlinear Science (CeNSc), Post-Graduate and
Research Department of Physics, Government College for
Women (Autonomous), Kumbakonam 612001, India

B. Tamil Arasan
Department of Physics, Presidency College (Autonomous),
University Of Madras, Chennai 600005, India



Multiple bright and dark soliton solutions in three component spinor Bose-Einstein condensates

V. Rajadurai^a, V. Ramesh Kumar^{a,*}, R. Radha^b

^a Department of Physics, Velammal Engineering College (Autonomous), Ambattur Redhills-road, Chennai, 600066, India

^b Centre for Nonlinear Science, Department of Physics, Government College for Women (Autonomous), Kumbakonam, 612001, India

ARTICLE INFO

Article history:

Received 28 August 2019
 Received in revised form 18 November 2019
 Accepted 20 November 2019
 Available online 27 November 2019
 Communicated by B. Malomed

Keywords:

Spinor Bose-Einstein condensate
 Three component Gross-Pitaevskii equation
 Soliton solution
 Gauge transformation method

ABSTRACT

We investigate a quasi one dimensional spin-1 Bose-Einstein Condensates (BEC) in the absence of an external confinement governed by a system of three coupled Gross-Pitaevskii (GP) equation. Based on the Lax-pair, we construct one soliton solution employing gauge transformation method. In addition, the multiple bright and dark soliton solutions are obtained by properly choosing amplitude dependent parameter in the Lax-pair. The results of the paper emphasizes the richness in the structure of soliton solutions admitted by the spin components, a phenomenon which has never been brought out to the fore. We have also extended the gauge transformation method to generate two soliton solutions.

2019 Elsevier B.V. All rights reserved.

1. Introduction

A large amount of bosons would occupy the lowest energy state when they are cooled close to absolute zero. This novel state, proposed by Bose and Einstein in 1925, is named as Bose-Einstein condensate (BEC). It was experimentally realized by Eric Cornell, Carl Wieman and Wolfgang Ketterle in 1995 [1]. After that in 1998 [2], Tin-Lun Ho and Ohmi group observed BEC in Na^{23} and Rb^{87} using dipole optical trap. They were able to realize BEC with internal degrees of freedom corresponding to the three hyperfine spin states ($m_f = +1, 0, -1$ for $F = 1$). It is known as “spinor BECs”. The important aspect of spinor BEC is that two or more hyperfine states of atoms in the condensates almost have same energy. So, the spin degrees of freedom impacts the evolution of condensates that are not present in the usual single-component BEC.

The dilute gaseous BEC with external trapping is modeled by Gross-Pitaevskii (GP) equation [3]. The nonlinearity in GP equation is governed by inter-atomic interaction which can be controlled by s -wave scattering length using external magnetic, optical or dc-electric field [4]. The possibility of controlling the inter atomic interaction in BECs has been witnessed in many experimental and theoretical studies.

Single component GP equation has been shown to admit bright solitons [5], dark solitons [6], vortices [7], etc., while two-component GP equation admits bright [8], ring dark [9], bright-dark [10], dark-dark and dark-bright solitons [11]. The multi component spinor BEC has been investigated and different types of solitons, like dark solitons [12], gap solitons [13], bright solitons [14], bright-dark soliton complexes [15], magnetic solitons [16], breathing solitons [17], dark-bright solitons [18], rogue waves [19], spin domains and spin textures [20–22] have been reported. The modulational instability of the spinor BEC model has been investigated in [23] and some exact solutions were reported in [24]. The non-autonomous three component GP equation has been analyzed in [25] and the dynamics of $F = 1$ spinor BEC has been investigated employing Hirota method. The three component defocussing nonlinear Schrödinger equation is analyzed in [26] using inverse scattering method. The dynamics of the polar bright soliton under the influence of SOC in spin-1 BECs is reported in [27] and some solutions were presented in Refs. [28–31].

Miki Wadati and co-workers had introduced three component integrable GP equation using 2×2 matrix [32]. They observed that bright solitons in spinor BECs show two spin states such as ferromagnetic (nonzero total spin) and polar (zero total spin) states. A specific

* Corresponding author.

E-mail addresses: rameshkumar@velammal.edu.in, vrama22@gmail.com (V. Ramesh Kumar).

Relief From Stress: It Is In Your Finger Tips

Dr.R.VIDHYA SREE
Director Of Physical Education
Govt.College For Women,Kumbakonam

Introduction:

There's a thing about yoga which is unlike gymming or doing any other form of exercise – it helps you connect with your inner self and to gradually uncover it. Our inner selves are in a perpetual blissful state and yoga can help you reach there. A thing as simple as hand gestures, known as mudras in yoga, can help you relax and unwind. Mudras are done in conjunction with breathing exercises and help regulate the flow of energy throughout your body. The word Mudra has two words. First one is 'muda' and second one is 'dra'. In Samskruth, 'muda' means happiness, 'dra means drawing. That means 'drawing happiness' by practicing mudras as per the Science of mudras given by our ancient sages. These days Yoga, Ayurved and Science of Mudras have been doing the rounds all over the world. When we study and incorporate the same in our daily life, health is not a miracle but can be achieved by adopting simple methods. Since mudras are part of Yoga, they are called 'Yoga Mudras'. Nerves and energy centres keep creating energy and flowing and glowing on the outer parts of human body. That lively vital energy can enter back to the body by practising Yoga Mudras. Universe is made of 5 elements – Fire, Air, Ether/Space, Earth and Water. If we wish to be healthy, we must ensure balance among 5 elements. This is the secret of being healthy.

Electro-magnetic waves are sent to our brain when we join any one of the tips of fingers with thumb. Different mudras are formed with different fingers joining with thumb. As per the Mudra Science, this is called circuit bypass. This energy will reach the brain and incite the nerve centres which will make the energy to reach different organs of the body to activate them in right manner. Five fingers of our hand represent 5 elements. Thumb gives basic foundation to other fingers of the hand to make the body healthy. Fire element helps other elements to keep them in right proportion. We are alive only when there is fire element in the body. Thumb represents fire element. These are not as per science's definition of elements but refer to the five building blocks that go into the formation of any living body. Several ancient health systems are based on the concept of the balance of the five elements. Indian Medical science according to Ayurveda and metaphysics affirm that distortion or impairment of the 5 elements create outer disturbance and inner sickness in the body.

The Five Fingers of the Hands represent these Five Elements:

- The Thumb symbolises the Fire
- The Forefinger symbolises the Wind
- The Middle finger symbolises the Ether
- The Ring finger symbolises the Earth
- The little or small finger symbolises the Water

Keeping specified nerves stretched for specified periods tones up of the nervous system. The fingers of each hand are held folded in certain specific postures and this provides the required tension on the nerves.

The fingers of each and every individual in the world are different in their shapes and sizes. These are determined and provided by nature as a tool to bring the nerves into prime condition when affected adversely. Acupressure where the nerves are influenced by the application of pressure on certain points or Acupuncture, where slight electrical impulses are conveyed through needles inserted in the body. The advantage in Mudras is that the pressure to be applied on the nerves is automatic and controlled by the shape and size of the fingers and not by external agencies. Because the Mudras work on the nerves, they are a NEURAL SCIENCE.



Versatile fabrication and characterization of Cu-doped ZrO₂ nanoparticles: enhanced photocatalytic and photoluminescence properties

Rajesh Gopal¹ · Akilandeswari Sambandam^{1,2} · Thirumalai Kuppulingam³ · Swaminathan Meenakshisundaram⁴ · Mohamad S. AlSalhi⁵ · Sandhanasamy Devanesan⁵

Received: 11 November 2019 / Accepted: 21 March 2020
© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

Pristine ZrO₂ and molar ratios of Cu-doped ZrO₂ nanoparticles were synthesized by chemical precipitation. The XRD pattern of pristine ZrO₂ and Cu-doped ZrO₂ nanoparticles shows the formation of t-tetragonal phase and shifting of tetragonal phase to monoclinic phase. The average crystallite sizes of the pristine ZrO₂ and Cu-doped ZrO₂ nanoparticles were calculated as 2.9, 4.65, 4.76, 5.02, and 4.96 nm, respectively. The Cu (0.06 M)-doped ZrO₂ nanoparticles are present as spherical morphology and high agglomeration was confirmed by FE-SEM and TEM analyses. The XPS spectra affirmed the presence of Cu²⁺, Zr⁴⁺, and oxygen ions in the Cu (0.06 M)-doped ZrO₂ nanoparticles. The Cu-doped ZrO₂ nanoparticles exhibit two energy gaps at 5.30 and 3.05 eV, 5.21 and 2.22 eV, 4.65 and 2.11 eV, and 4.44 and 2.65 eV. The surface defects and oxygen vacancies were analyzed by PL and ESR spectroscopy. The ESR spectra are clearly asymmetric in shape and contain a peak signal related to the presence of the copper species (Cu²⁺) in the distorted tetragonal coordination of ZrO₂. The photocatalytic activities of Cu (0.06 M)-doped ZrO₂ nanoparticles were successfully sought on degradation of the two azo dyes: methyl violet and methyl blue under sunlight irradiation. Cu (0.06 M)-doped ZrO₂ showed complete degradation at 70 min. The reusability of Cu (0.06 M)-doped ZrO₂ nanoparticles shows maximum degradation efficiency for six successive runs.

1 Introduction

Dyes are used in a variety of different commercial enterprises, for products such as leather, textiles, plastics, cosmetics, paper, and petrochemical industry. These different

industries produce both organic and inorganic pollutants that are released into the water sources. The pollution of water has become a major issue and affecting the human beings, animals, and plants. Therefore, major drive to reduce the level of pollution before the release of industrial waste [1–3]. The scientific community has used various techniques to remove the organic/inorganic pollutants from water, like coagulation [4], bio-degradation [5], ultra-filtration [6], processes using membranes [7], reverse osmosis [8], advance oxidation processes (AOP) [9], and so on. Among these techniques, photocatalysis is one of the most important approaches and it can completely degrade the hazardous pollutants to H₂O and CO₂ [1–3].

Photocatalysis occurs when a photon is absorbed onto the surface of a semiconductor-based material, and electron–hole pairs are generated. The generated electron and hole pairs react with H₂O, inducing oxidation and/or reduction. This technique has proved useful for several applications such as air purification [10], wastewater treatment [11], water splitting [12], disinfection [13], CO₂ reduction [14], and in surface cleaning treatments [15]. Many oxide and sulfide materials such as ZnO [16], SiO₂ [17], SnO₂ [18],

✉ Akilandeswari Sambandam
akilaphy2010@gmail.com

✉ Mohamad S. AlSalhi
malsalhi@ksu.edu.sa

¹ Department of Physics, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India

² Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India

³ Department of Chemistry, Government Arts College, Thiruvannamalai, Tamil Nadu, India

⁴ Department of Chemistry, Annamalai University, Annamalai Nagar, Chidambaram, Tamilnadu 608 002, India

⁵ Department of Physics and Astronomy, College of Science, King Saud University, P.O. Box 2455, Riyadh 11451, Kingdom of Saudi Arabia



Enhancement of photocatalytic activity of ZrO₂ nanoparticles by doping with Mg for UV light photocatalytic degradation of methyl violet and methyl blue dyes

G. Rajesh¹ · S. Akilandeswari^{1,2} · D. Govindarajan¹ · K. Thirumalai³

Received: 23 June 2019 / Accepted: 20 January 2020 / Published online: 3 February 2020
© Springer Science+Business Media, LLC, part of Springer Nature 2020

Abstract

In this study, pristine ZrO₂ (600 °C) and Mg (0.02, 0.04, 0.06, 0.08 M)-doped ZrO₂ nanoparticles were effectively synthesized at about 10 nm size and achieved complete degradation of methyl violet and methyl blue dyes under UV irradiation. The calcined products were analyzed by XRD, FTIR, UV-DRS, PL, FESEM-EDX, TEM-SAED pattern and XPS techniques. The tetragonal crystal structure of the pristine ZrO₂ and Mg (0.02–0.08 M)-doped ZrO₂ nanoparticles were affirmed by XRD analysis. The Mg (0.08 M)-doped ZrO₂ nanoparticles were approximately in quasi-spherical morphology and high agglomeration was confirmed by FESEM and TEM results. The surface defects and oxygen vacancies were analyzed by PL spectroscopy. The Mg (0.08 M)-doped ZrO₂ nanoparticles exhibited enlarged photocatalytic activity with 94% and 90% degradation of methyl violet and methyl blue dyes under UV irradiation.

1 Introduction

The various organic dyes are increasingly applied in many industries such as leather tanning, textile, plastic, paper, food, pharmaceutical, cosmetics, photo-electrochemical cells, hair coloring, rubber and agricultural research and so on [1–3]. These types of industries subsequently discharge their colored wastewater containing pigments and dyes. Colored wastewater (dyes) significantly attributed to environmental pollution and health risk [4]. Almost these colored wastewaters are toxic and non-biodegradable. Various physical, biological and chemical techniques have been applied for its removals such as precipitation, activated carbon, ozonation, membrane separation, Fenton, photo-Fenton catalytic reaction, bioremediation, adsorption, reverse osmosis, coagulation and ultra-filtration [1–3, 5]. Among these techniques, photocatalysis is the most important approach

because of its low cost, eco-friendliness and totally oxidizes the pollutants to H₂O and CO₂ [1–3].

There are many materials applied for photocatalytic degradation including TiO₂ [6] ZnO [7], WO₃ [8], ZrO₂ [9], SnO₂ [10], In₂O₃ [11], Fe₂O₃ [12], NiO [13], CuO [14] CdS [15], and PbS [16]. Among these, zirconium oxide (ZrO₂) is a most attractive material. ZrO₂ is a wide band gap (5.0–5.5 eV) semiconductor, and this oxide material had detected broad application in ceramics, catalysts, gas sensors, fuel cell, solid-state electrolytes, barrier coatings and optical devices [17–21]. It exhibits excellent thermal, mechanical, electrical and optical properties, such as low thermal conductivity, high hardness, high fracture toughness, high refractive index, optical transparency, high corrosion resistance and polymorphic nature [17–22]. Zirconium oxide (ZrO₂) exhibits three kinds of polymorphs as follows: Monoclinic (m-ZrO₂), tetragonal (t-ZrO₂) and cubic (c-ZrO₂). The m-monoclinic (m-ZrO₂) phase is stable below at < 1170 °C, t-tetragonal (t-ZrO₂) phase is stable at 1170–2370 °C, and c-cubic (c-ZrO₂) phase is stable at > 2370 °C, respectively [23]. These crystal phases strongly depend on thermal condition and preparation method [24].


Moreover, the t-tetragonal phase (t-ZrO₂) has strongly improved photocatalytic activity compared to c-cubic (c-ZrO₂) and m-monoclinic (m-ZrO₂) phase zirconium oxide (ZrO₂) materials [25, 26]. Pristine zirconia is essentially a poor oxide ion conductor at lesser temperature. Hence,

✉ S. Akilandeswari
akilaphy2010@gmail.com

¹ Department of Physics, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India

² Department of Physics, Government College for Women (Autonomous), Tamil Nadu, Kumbakonam 612001, India

³ Department of Chemistry, Government Arts College, Tiruvannamalai, Tamil Nadu, India



Biosynthesis of silver nanoparticles using *phyllanthus emblica* fruit extract for antimicrobial application

R. Renuka^{a,*}, K. Renuka Devi^b, M. Sivakami^b, T. Thilagavathi^c, R. Uthrakumar^d,
K. Kaviyarasu^{e,1}

^a Department of Physics, Government Arts College for Women (Autonomous), Pudukkottai, Tamil Nadu, 622001, India

^b Department of Physics, H.H The Rajah's College (Autonomous), Pudukkottai, Tamil Nadu, 622001, India

^c Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, 612001, India

^d Department of Physics, Government Arts College (Autonomous), Salem, Tamil Nadu, 636007, India

^e UNESCO-UNISA Africa Chair in Nanosciences/Nanotechnology Laboratories, College of Graduate Studies, University of South Africa (UNISA), Muckleneuk Ridge, PO Box 392, Pretoria, South Africa

^f Nanosciences African Network (NANOAFNET), Materials Research Group (MRG), iThemba LABS-National Research Foundation (NRF), 1 Old Faure Road, 7129, PO Box 722, Somerset West, Western Cape Province, South Africa

ARTICLE INFO

Keywords:

Biosynthesis
Silver nanoparticles
Phyllanthus emblica
UV-Vis
Antibacterial activities

ABSTRACT

Biological synthesis of metal nanoparticles is one of the effortless, cost-effective, environmental-friendly methods and scale down the handling of toxic chemicals. The silver nanoparticles were triumphantly synthesized from the silver nitrate solution through a greener route using Indian gooseberry (*phyllanthus emblica*) fruit extract and the prepared particles are of well-balanced shape and size. The stabilization and the bio-reduction of silver nanoparticles were described by adopting UV-vis spectrometry interprets the surface plasmon resonance peak at 449 nm. The functional groups of *phyllanthus emblica* fruit extract and the silver nanoparticles were diagnosed by using Fourier transform Infrared spectrum (FTIR). The crystalline nature of face-centered cubic (FCC) was examined by X-ray diffraction (XRD) and the crystalline size ranges between 19 nm to 45 nm with an average size 30 nm. Scanning electron microscope (SEM) witnessed the hexagonal shape of silver nanoparticles and Energy-dispersive X-ray spectroscopy (EDX) assured the well-established fabrication of nanocrystalline silver element. The synthesized nanoparticle exhibits significant antibacterial effects and very clear zone of inhibition against *Klebsiella pneumoniae* and *Staphylococcus aureus* bacteria.

1. Introduction

The research on metal nanoparticles is a foremost field of nanotechnology because of their massive applications in the areas of physics, chemistry, medicine and biology (Ponarulselvam et al., 2012). The most essential and specific properties of nanomaterials are having high surface area to volume ratio exhibits remarkable antibacterial properties (Ahmed et al., 2016). Due to their noble properties, silver nanoparticles may have vast applications in the areas of microelectronics, photonics and photo catalyst and lithography (Shankar et al., 2017). Therefore, nanoparticles can be formulated from different physicochemical methods and the drawbacks of physicochemical methods are mainly because of the handling of toxic chemicals, the production of harmful toxic wastes, requires high pressure and tem-

perature (Iyyappa Rajan et al., 2017 & Valan Arasu et al., 2019). Because of these drawbacks, the valuable alternative and attractive biological methods were focused in the present work. The bio-inspired synthesis using plants are the nontoxic, easily available, low-cost, environmentally friendly methods (Ahmed and Ikram, 2015). Not only the greener method is environmentally friendly, but they can also be employed to fabricate large quantities of nanoparticles and free of contamination (Mittal et al., 2013).

Fabrication of nanoparticles by adopting plant extract is also differing from microbial synthesis because they are relatively fast, no need to maintain culture conditions and so this method was appropriate to fabricate nanoparticles of distinct size and shape (Tarannum et al., 2019). Bio-fabrication of silver nanoparticles using *phyllanthus emblica* fruit extract has been reported in the present work. It has been

* Corresponding author.

E-mail address: renuka101281@gmail.com (R. Renuka).



Green synthesis of magnetic nanoparticles via *Cinnamomum verum* bark extract for biological application

M. Sivakami^{a,*}, K. Renuka Devi^b, R. Renuka^b, T. Thilagavathi^c

^a H.H. The Rajah's College, Pudukkottai, 622001, India

^b Government Arts College for Women, Pudukkottai, 622001, India

^c Government Arts College for Women, Kumbakonam, 612001, India

ARTICLE INFO

Editor: GL Dotto

Keywords:

Biosynthesis
Nanoparticles
Cinnamomum verum bark
Antibacterial activity
Scanning electron micrographs
Vibrating sample magnetometer analysis

ABSTRACT

The green synthesis of magnetic Fe nanoparticles (NPs) was prepared via *Cinnamomum Verum* bark extract. The UV–vis analysis explained the formation of Fe NPs with SPR band at 288 nm. The powder X-ray diffraction analysis (XRD) explicated the mean crystallite size was at 36 nm. The Fourier Transform Infra-Red analysis (FTIR) revealed the functional groups in the prepared Fe NPs. The Scanning Electron Microscope (SEM) and High-Resolution Transmission Electron Microscope (HR-TEM) analyses notified the circular and spherical shaped Fe NPs with the size at 20–50 nm. The Energy dispersive X-ray spectroscopy (EDS) and mapping analyses validated the formation of Fe NPs with purity. The Vibrating Sample Magnetometer (VSM) revealed the paramagnetic behavior of prepared Fe NPs. The phytochemical analysis described the phytochemicals compounds which present in *Cinnamomum Verum* bark extract. The Antibacterial assay described the much higher inhibition zone of prepared Fe NPs upon human pathogenic bacteria. The Antioxidant (DPPH) assay described the efficient scavenging behavior of Fe NPs with 89 % at 80 µg/mL concentration. The Anti-inflammatory assay explicated the potential protein denaturation behavior of Fe NPs with 87 % at 500 µg/mL concentration. The Anti-diabetic assay reported the much higher potential efficiency of prepared Fe NPs with 84 % at 500 µg/mL concentration.

1. Introduction

In recent decades, nanotechnology has gained more recognition due to its unique properties associated with the size distribution and morphology of nanoparticles. Nanotechnology was an umbrella term that covers many research areas dealing with objects that are covered in nanometers such as chemistry, physics, biology, engineering and other scientific aspects of nanotechnology [1,2]. Nanoparticles having 1–100 nm possess great impact in the field of chemistry, optics, batteries, physics, environmental remediation, drug delivery and medicine. Nanoparticles exhibit enormous structures which create a different approach in catalytic, physical, chemical and medicinal properties of materials than bulk [3–5].

Nowadays, the researchers concentrate on the metal nanoparticles due to the large surface area, low melting point, and good optical, catalytic, electrical and thermal properties. These distinctive properties of metal nanoparticles create exploitation in the industrial area such as food, agriculture, space, cosmetics, medical and chemical aspects of use

in day-to-day life [6–9].

Recent research in the synthesis of nanoparticles opened a new era in the fast-growing method for the production of nanoparticles. Physical and chemical methods were usually used for the synthesis of nanoparticles, however, based on the toxicity of these methods; the objective of the research has latterly moved towards the biosynthetic method [10–12].

Nanoparticles of magnetic materials have attracted much consideration due to their properties deviate from those of bulk materials and they can be used to make materials and devices with new properties [1]. Magnetic Nanoparticles have important applications in magnetic storage devices, in ferrofluids, i.e. stable suspensions of magnetic nanoparticles, in magnetic beads that are applied in biotechnology, for contrast enhancement in magnetic resonance imaging (MRI), and targeted drug delivery, bioelectrochemical sensing, environmental remediation and as an electrode for supercapacitors and lithium-ion batteries [2]. The magnetic nanoparticles received from the green synthesis method using plants applied in many fields such as semiconductors [13], catalysts [14], optics, chemistry [15] and medicine [16].

* Corresponding author at: Department of Physics, H.H. The Rajah's College, Pudukkottai, 622001, India.

E-mail address: Siva_hhrc@yahoo.in (S. M.).

<https://doi.org/10.1016/j.jece.2020.104420>

Received 11 July 2020; Received in revised form 16 August 2020; Accepted 20 August 2020

Available online 5 September 2020

2213-3437/© 2020 Elsevier Ltd. All rights reserved.

SP-50

PHYTOCHEMICAL CHARACTERIZATION OF *NARINGI CRENULATA* (ROXB) LEAF WITH METHANOLIC EXTRACT BY GC-MS METHOD.

¹B. BHAVANI AND S. VASANTHA²

¹Department of Botany, Government college for woman (Autonomous), Kumbakonam, Thanjavur, Tamil Nadu, India. Affiliated to Bharathidasan University, Trichirappalli

²P.G and Research Department of Botany and Microbiology, A.V.V.M Sri Pushpam College (Autonomous), Poondi, Thanjavur, Tamil Nadu, India. Affiliated to Bharathidasan University, Trichirappalli

Corresponding author Email: bhavani1493@gmail.com
Mobile: +91 9843662674

ABSTRACT

The bioactive compounds of *Naringi crenulata* (Roxb) leaves have been evaluated using GC-MS. The active chemical compounds were identified by using Gas Chromatography-Mass Spectrometry. GC-MS analysis of leaves extracts *Naringi crenulata* in the characterization of photochemical sixteen components were identified. GC-MS results were Octane, 2,4,6-trimethyl, 1-Iodo-2-methylundecane, Hexadecane, Methoxy acetic acid, Pentadecanoic acid etc., the existence of other major components was also presented.

KEYWORDS: Methanol extract, *Naringi crenulata*, GC-MS analysis and photochemical.

INTRODUCTION

Herbal medicine is one of the most remarkable uses of plant-based diversity. As many as 75 to 90% of the world's rural people rely on herbal medicine. The success of any health care system depends on the availability of suitable drugs on a sustainable basis. Natural medicine improves the strength of the body. The knowledge medicinal plant has been accumulated in many centuries based on different medicinal systems such as Ayurveda, Unani and Siddha. In India, it is reported that traditional healers use 2500 plants species 100 species plant serves as regular sources of medicine during the last decades there had been an increasing interest in the study of medicinal plants and their traditional use in different parts of the world. According to the reports of the world health organization (WHO), as many as 80% of the world's people depend on traditional medicinal for their primary health care needs due to the considerable economic benefit in their development and All parts of this tree viz. root, stem, bark, leaf and fruit has been used as folk medicine¹. It has been used as folk medicine¹. The root extract of this plant is used for curing vomiting, dysentery, and colic disorders.^{2,3} bioactive components of leaves and bark of *Naringi crenulata* using GC-MS⁴. Analysis of methanolic extract stem³. Various parts of this plant have been employed in indigenous medicine and it is used as antiepileptic, purgative, sudoferic, colic trouble and cardialgia.⁵ Leaves are used as a remedy for epilepsy. Bark is aromatic and cooling and is useful in vitiated conditions of Pitta⁷. Crenulatine along with twenty known indole alkaloids were isolated from the stem of the plant⁸. GC-MS is a combination of two different analytical techniques Gas chromatography (GC) and Mass Spectrometry (MS), used to analyze biochemical and organic samples. GC can separate semi-volatile and volatile compounds present in the sample with great resolution, but it cannot identify them. While the MS can provide detailed structural information so that they can be identified but cannot be quantified. Application of GCMS is to monitor and clean the environment, criminal forensics, law enforcement, security, food, beverage, and perfume analysis. It can also be used in astrochemistry and medical field⁹.

MATERIALS AND METHODS

Collection of plant materials

The fresh and healthy leaves of the plant *N. crenulata* were collected from around Thanjavur, Tamil Nadu, India., during the 2017 and identified with Rapinat Herbarium by Principal Scientific Officer Dr.S.John Britto, St. Joseph's College, Tiruchirappalli, Tamil Nadu, India.

SP-49

FT-IR SPECTROSCOPIC ANALYSIS OF LEAF EXTRACTS OF *NARINGI CRENULATA* (ROXB.) NICOLS.

B. BHAVANI AND S. VASANTHA

*Department of Botany, Government college for women (Autonomous), Kundabalam, Thanjavur, Tamil Nadu, India.

Affiliated to Bharathidasan University, Trichirappalli

*P.G and Research Department of Botany and Microbiology, A.V.V.M Sri Pushpam College

(Autonomous), Poomdi, Thanjavur, Tamil Nadu, India.

Affiliated to Bharathidasan University, Trichirappalli

Corresponding author Email: bhavani1493@gmail.com

Mobile: +91 9543662674

ABSTRACT

Medicinal plants play a major role in the life of human beings throughout the globe. These plants are commercially important because of the presence of various chemical substances with the power of curing different diseases and healing several ailments affecting the normal activity of human beings. These economically important plants are found growing in remote forests, hills, and mountains where human invasion and intervention are minimum. Special knowledge about the location, usage and benefits. The Traditional Aboriginal Knowledge (TAK) is prevalent among the members of certain tribal communities inhabiting these geographical locations. The importance of medicinal plants was known to the outside world with the advent of communication technology and messages published in mass media. Increasing commercial pressure on these medicinal plants and competition among the drug manufacturers reflected on the purity and genuineness of the drugs. This ultimately resulted in several side effects in the physiological functions of the persons using these drugs. Hence pharmacognostical study of such wild plants is needed to safeguard the users. The native tribes Irulars and Malasars in the Vallangiri Hills of the Western Ghats use different parts of *Naringi crenulata* (*N.crenulata*) (Roxb.) Nicols (Plate 1) to cure different kinds of ailments. From the analysis of data of FTIR, strong absorption bands at alcohols, phenols, alkyis, amines, and some functional groups presented.

KEYWORDS: *Naringi crenulata* methanolic extract, FT-IR, analysis, photo components.

INTRODUCTION

The Traditional medicinal plant for several thousand years'. Medicinal plants are a large number of users as an alternative medicine for diseases of humans and animals since most of them are without side effects when compared with synthetic drugs. The chemical compounds' identification of the chemical nature of present in the medicinal plants provided some information on the different functional groups responsible for their medicinal properties. Studying the in-vitro efficacy of bioactive extracts of fifteen medicinal plants against multidrug resistant microbes¹. The plants are used as remedies for human disease photochemical components of therapeutic value that produce physiological action on the human body². The mainly used to various human diseases, treatment of ailment and health of affected organs from the period immemorial³. The world's population- 30% traditional remedies for their healthcare. Today, about 70,000 to 80,000 plants are used for medicinal. This is because of some biological active and naturally occurring phytochemical present in the various parts of plants. The plant '*Naringi crenulata*' (Roxb.) Nicolson belongs to the Rutaceae family, commonly known as 'kannarakkam', 'Malanarakkam', in Malayalam, 'Maharivum' in Tamil, a widespread species of the genus "*Naringi*". It has been used as folk medicine⁴. The root extract is used for vomiting, dysentery and colic disorders⁵. It is reported that its methanolic extract showed significant anthelmintic activity⁶. The photochemical compounds major groups as the most activity in plant extracts by infrared spectroscopy screened the bioactive group of chemicals in the dry leaf powder by FTIR analysis⁷. Numerous species of *Naringi crenulata* are known to possess a variety of biological activities including anti aging⁸. leaves are used for curing mental disorders⁹ folk



Molecular docking and network pharmacology-based approaches to explore the potential of terpenoids for *Mycobacterium tuberculosis*

K. Kalaimathi^a, G. Thiyagarajan^b, S. Vijayakumar^c, K. Bhavani^d, K. Karthikeyan^b, J. Maria Jancy Rani^e, K. Dass^f, J. Sureshkumar^g, S. Prabhu^{g,*}

^a Department of Chemistry Government College of Engineering, Sengipatti, Thanjavur-613 402, Tamil Nadu, India

^b Department of Botany Government Arts College, (Autonomous), Affiliated to Bharathidasan University, Kumbakonam-612 002, Tamil Nadu, India

^c PG and Research Department of Botany V. V. M Sri Pushpam College (Autonomous), Affiliated to Bharathidasan University, Poondi-613 503, Thanjavur, India

^d Department of Zoology Government College for Women (Autonomous), Affiliated to Bharathidasan University, Kumbakonam-612 001, Tamil Nadu, India

^e Department of Chemistry Annai Vailankanni Arts and Science College (Affiliated to Bharathidasan University), Bishop Sundaram Campus, Thanjavur-613 007, Tamil Nadu, India

^f Department of Zoology Annai Vailankanni Arts and Science College (Affiliated to Bharathidasan University), Bishop Sundaram Campus, Thanjavur-613 007, Tamil Nadu, India

^g Department of Botany Annai Vailankanni Arts and Science College (Affiliated to Bharathidasan University), Bishop Sundaram Campus, Thanjavur-613 007, Tamil Nadu, India



ARTICLE INFO

Keywords:

Racemosol

Pacharin

Mtb

Molecular docking

Network pharmacology

ABSTRACT

Objective: *Mycobacterium tuberculosis* (Mtb) infection is quickly countered by the host immune system; but, the pathogen is never eradicated. In this infection, 10% of infections are considered as open tuberculosis, while the remaining 90% are becoming latent, a state that can persist in the host for decades until immune control is lost. To date, more than 350 plant species have been studied in mycobacterial research.

Methods: In this research, we used a mixture of software and database explorations to find the anti-mycobacterial possibilities of phytoconstituents. Maestro v12.7 and Cytoscape v3.8.0 were applied as software. In addition, the String v11.0, KEGG, Reactome and Pass Server databases assisted us in understanding TB cellular processes and phytoconstituent drug possibilities.

Results: The racemosol and pacharin have registered higher docking scores of 7.823 and -7.715 respectively. Furthermore, these compounds have substantial binding affinities in the active pocket of Mtb, and their docking metrics have consisted close to that of an approved drug (Rifampin). The network pharmacological exploration is also revealed possible connections with all of the TB-associated genes and other important signaling pathways.

Conclusions: This computational study found that racemosol and pacharin were the safest anti-mycobacterial drugs for tuberculosis. Following comprehensive research of the pharmacological properties of racemosol and pacharin, we decided that this would be the first study to focus on anti-mycobacterial activity utilizing those compounds. Also, we believe that racemosol and pacharin compounds will reveal potential actions when used in mycobacterial research. Furthermore, the derivatives of these compounds will be acted as possible drugs for tuberculosis and associated diseases. More pharmacological and toxicological explorations are required to have a better understanding of these compounds.

1. Introduction

Mycobacterium tuberculosis (Mtb) is the bacteria that cause Tuberculosis (TB). It is considered to be the world's lethal infectious disease agent and leads to cause high fatality rate in developing countries including South-East Asia, India, Africa, Western Pacific, Indonesia, China, Eastern Mediterranean, Philippines, Pakistan, Nigeria, Bangladesh, South Africa, America and Europe [1]. This bacterium

spreads extremely faster when there are a huge number of people present, as well as in situations of starvation and poverty [2]. It primarily infects the respiratory system of people. In the case of respiratory tract infections, it reveals the signs such as persistent cough, chest discomfort, hemoptysis, weakness or weariness, weight loss, fever, and night sweats [3].

Since 1996, the World Health Organization has issued a global TB prevalence report every year [1]. In 2020, data from 198 countries and territories represented more than 99% of the world's population and

* Corresponding author.

E-mail address: prabhushbotany@gmail.com (S. Prabhu).

<https://doi.org/10.1016/j.prmcm.2021.100002>

Received 28 June 2021; Received in revised form 8 August 2021; Accepted 11 August 2021

Available online 13 August 2021

2667-1425/© 2021 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>)




GOVERNMENT COLLEGE FOR WOMEN (AUTONOMOUS)

KUMBAKONAM – 612 001

Affiliated to Bharathidasan University

DST - CURIE Sponsored Institution

IV Cycle of Accreditation

 **0435 – 2401391**

 **principal@gwk.ac.in**



CRITERION III – RESEARCH, INNOVATIONS AND EXTENSION

FRONT PAGE OF PUBLISHED PAPERS

2019-2020

The Pharma Innovation



Haematological status of common carp, *Cyprinus carpio* L. exposed to sublethal dose of organophosphorus pesticide, monocrotophos

Sharmila Gunasekaran and Kavitha Arumugam Vellaichamy

Abstract

The introduction of organophosphorus insecticide into the aquatic ecosystem through agricultural runoffs, rainwater seepage adversely affects many non-target organisms including fishes. The chronic effect of organophosphorus pesticide, Monocrotophos on the haematological parameters of freshwater fish, *Cyprinus carpio* has been investigated in the present study. *Cyprinus carpio* were procured and their fish, *Cyprinus carpio* has been investigated in the present study. *Cyprinus carpio* were procured and their concentrations (T2) and (T3) were selected to which the fishes were exposed for 30 days. Control set up without monocrotophos (T1) was run simultaneously. Blood from anaesthetised fishes were collected after 30 days of exposure period. Hemoglobin (Hb), Red Blood Cells (RBCs) and hematocrit (Hct) significantly decreased with the increasing concentration of monocrotophos during the experimental periods. But White Blood Cells (WBCs) significantly increased in the treated fishes when compared to the control depicting negative effects of monocrotophos on the haematological parameters of *Cyprinus carpio*.

Keywords: *Cyprinus carpio*, chronic effect, haematology, monocrotophos, sublethal concentration

1. Introduction

Applications of pesticides have been increasing ever since the onset of green revolution to feed the overgrowing population of the world. This phenomenon is evident over the last few decades especially in tropical countries like India, where majority of the population is reliant on agriculture for their livelihood. Now the use of pesticides has become inevitable in current agriculture practices to ensure increased productivity through the pest control. However, the indiscriminate use of pesticides ultimately leads to pollution of aquatic environment and becomes hazardous to the non-target aquatic life. Among the innumerable aquatic organisms, fishes occupy an important position in the field of aquatic toxicology (Giulio and Hinton, 2008) [7] not only because of their position at topmost trophic level of the aquatic food chain but also due to the high accumulation of pesticides in their body by bioaccumulation and biomagnification processes. Hence different concentrations of insecticides and pesticides present in water bodies found their way to fishes affecting these non-target species (Talebi, 1998; Uner *et al.*, 2006; Banaee *et al.*, 2008) [18, 20, 1].

Various stressors and pollutants generally cause rapid changes in biological characteristics of fish (Sahan *et al.*, 2007) [15]. These changes can be measured and used as indicators or biomarkers for effects of such toxicants. These biomarkers enable the rapid assessment of the health of the organisms and warn about environmental risks associated with the toxicants. Among biological changes, haematological parameters are considered potential biomarkers of exposure to toxicants, since the latter can induce an increase or decrease in the various haematological parameters (Oost *et al.*, 2003) [12]. Blood is the most important and abundant body fluid acting as a vehicle for quickly mobilizing defense against trauma and diseases. Its composition often reflects the total physiological condition of the body. The main route of entry for any pesticide is through the gills from which it gets transported to various parts of the body via the blood stream. Henceforth blood provides an ideal medium for toxicity studies and haematological variables of fish under stress are of great significance in assessing the impacts of pollutants in the biota of a particular ecosystem.

Among the various pesticides used in farming practices, the organophosphate pesticide, Monocrotophos often referred to as the "farmer's benevolent pesticide" is most preferable by farmers because of their low cumulative ability, high insecticidal property, low mammalian

ISSN (E): 2277- 7695
ISSN (P): 2349-3242
NAAS Rating: 5.03
TPI 2019; 8(3): 178-182
© 2019 TPI
www.thepharmajournal.com
Received: 18-01-2019
Accepted: 22-02-2019

Sharmila Gunasekaran
Assistant Professor, PG &
Research Dept. of Zoology,
A.V.C. College (Autonomous),
Mannampandal, Mayiladuthurai,
Nagapattinam, Tamil Nadu,
India

Kavitha Arumugam Vellaichamy
Assistant Professor, Dept. of
Zoology, Govt. College for
Women (Autonomous),
Kumbakonam, Thanjavur,
Tamil Nadu, India

Correspondence

Sharmila Gunasekaran
Assistant Professor, PG &
Research Dept. of Zoology,
A.V.C. College (Autonomous),
Mannampandal, Mayiladuthurai,
Nagapattinam, Tamil Nadu,
India

Haematological variables of *Oreochromis mossambicus* against *Aeromonas hydrophila* infection by using dissimilar types of gaumutra distillate

Varadhan Praveena, Ravichandiran Vinitha, & Sournamanickam Venkatalakshmi*

Centre for Animal Studies, Post Graduate and Research, Department of Zoology, Government College for Women (Autonomous),
Kumbakonam- 612 001, Tamilnadu, India

[E-mail: dr.s.venkatalakshmi@gcw.ac.in]

Received 04 May 2018; revised 12 June 2018

In Aquafarming, gaumutra has been used to recuperate broad healthiness of a specific person. An analysis was conceded out to assess the consequence of different breeds of cow and bull urine (distillate) was used for the analysis of haematological parameters of *Oreochromis mossambicus* fish against *Aeromonas hydrophila* bacterial infection. Average weight (15.6 ± 0.2 g) of healthy fishes was maintained in 70 litres aquaria with most favourable temperature range from 25.8 ± 28.8 °C. Furthermore, fishes were exposed to different cow urine distillate (CUD) namely, C (without cow urine), T1 (Gir calf), T2 (Gir cow), T3 (Gir Bull calf) and T4 (Gir Bull) on 0.1 % concentration in the medium for seven days. *O. mossambicus* injected with heat-killed *A. hydrophila* (1×10^4 cells) during post stimulation phase. Meanwhile, blood samples were collected to determine the different haematological parameters viz. total RBCs, WBCs, haemoglobin concentration and mean corpuscular haemoglobin. One-way analysis of variance exhibited significant value ($P < 0.01$) of haematological parameters and as well as haematological contents. Based on the above findings, the present study revealed that T2 (Gir cow) urine distillate has the potential antimicrobial activity of *A. hydrophila* and also enhanced the health status of *O. mossambicus*.

[Keywords: *Aeromonas hydrophila*; Cow urine; haematology; *Oreochromis mossambicus*; T2- Gir cow]

Introduction

Aquafarming is a unique fast upward nourishment's creating sectors of the universe and intended to enlarge productivity per unit space. Fish farming is considered as an outstanding innovation in the use of organic waste. The manure (cow urine) and supplementary feeding play an imperative role in various types of fish farming practices. The feed and fertilizers cost could be reduced significantly by integrating fish farming with livestock farming which ultimately is economical with respect to the cost of production¹. The augmented escalation of aquaculture has managed a great number of pathogens, the major cause of the outbreak of diseases in fishes. Therefore, distant predictable methods such as the use of disinfectants and antibacterial medications have been inadequate achievement in the avoidance or therapy of fish infections. The enormous use of antimicrobials for disease resistor and development promotion in fish proliferations the selective pressure used on the microbial world and boosts the usual emergence of bacterial resistance². The use of inoculations in fish farming to avert the bacterial diseases is successful³,

but the main safeguard is species (pathogens) specific and expensive⁴.

Recently, cow urine has been reported as a bio enhancer source of allopathic antibiotics and anticancer drugs¹. It enhanced their effect and reduced the toxic substances and others including adverse effects of their synthetic drugs. As per Ayurveda, cow urine is needed to purify and detoxify in many drugs. Cow urine distillate (CUD) known as "Kamadhenu ark", exhibited many biological activities including immune modulatory and anti-potential anti-microbial effect of various living beings⁵. Freshwater fish *Oreochromis mossambicus* (Tilapia) is often used as a good experimental model and is extensively used in biological, genetic and physiological studies in relation to pollution, stress, or growth promoters^{6,7}. Tilapia is an excellent experimental model for haematological studies and microbial infection because of its worldwide economic importance and also tolerates poor water quality conditions. Therefore, the current work was intended to examine the haematological changes and antimicrobial effect of cow urine distillate of T2 (Gir cow) in *Oreochromis mossambicus*.



Impact of pesticide monocrotophos on microbial populations and histology of intestine in the Indian earthworm *Lampito mauritii* (Kinberg)

Vaithyanathasamy Kavitha^{a,b,*}, Ramachandran Anandhan^{a,c}, Naiyf S. Alharbi^d, Shine Kadaikunnan^d, Jamal M. Khaled^d, Taghreed N. Almanaa^d, Marimuthu Govindarajan^{a,b,c,*}

^a Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, 608 002, India

^b Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India

^c Department of Zoology, Government Arts College (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India

^d Department of Biotechnology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:
Insecticide
Ecotoxicology
Microbes
Earthworm
Intestine
Histology

ABSTRACT

Soil contamination has enlarged over the decades due to intensive use of pesticides and chemical fertilizers in agronomy. Earthworms are significant organisms in the soil community. Earthworms are the major role in soil fertility in most ecological system and the production of biogenic structures. Moreover, earthworm gut mucus enhances the beneficial soil microorganism potential biological activities. They are used as model organisms for assessing the ecological risks of chemicals. Enrichment of essential nutrients in soil through earthworm is a cost-effective and eco-friendly approach. In India, the organophosphorus pesticide monocrotophos is commonly used to control agricultural pests. Hence, it is important to study the effect of monocrotophos on the gut microbiota in *Lampito mauritii*. A 15-day exposure to a low (1/10th of the LC₅₀ after 96 h i.e., 0.093 ppm kg⁻¹) and high sublethal concentration (1/3rd of the LC₅₀ after 96 h i.e., 0.311 ppm kg⁻¹) of monocrotophos led to reduced proliferation of the gut microbiota in *L. mauritii*. However, exposure for 30 days led to a recuperation of the microbial populations to near control values. Among the eight bacterial and five fungal species that inhabit the gut of *L. mauritii*, only six bacterial and three fungal species were able to survive after exposure to monocrotophos. In addition to the study, histopathological changes were observed in the intestine of *L. mauritii* after application of lower sublethal concentration of monocrotophos. Severe pathological changes such as vacuolization, degenerated nuclei, damaged villi and congestion of the blood sinuses were noticed in the intestine on 1st and 5th day of the experiment. But in 30th day the damages were slowly recovered due to degradation of monocrotophos by the presence of some pesticides degrading bacterial and fungal species and regenerative capability of chloragogenous cells in the intestine. The results suggested that reduced microbial populations and pathological damages in intestine were observed during the application of monocrotophos. So, the monocrotophos have several harmful impacts on earthworms.

1. Introduction


There is a growing concern for soil contamination due to the widespread and global use of agropesticides. Knowledge about the effect(s) of pesticides in the soil environment is imperative since the resident soil organisms may be adversely affected. Depending on the type of soil organisms affected, this may have damaging consequences for soil fertility. A large proportion (80%) of the biomass of terrestrial invertebrates comprises earthworms that are important for structuring and increasing the nutrient content of soil. Therefore, they are suitable bioindicators for the chemical contamination of soil in terrestrial

ecosystems that can be used to provide an early warning for deteriorating soil quality [1,2]. This is important for protecting the natural environments and is of increasing interest in protecting human health [3] as well as other terrestrial vertebrates that prey on earthworms. Earthworms are suitable bioindicators of soil toxicity since they ingest large quantities of decomposed litter, manure, and other organic matter deposited on soil, thereby aiding the conversion into rich topsoil [4]. The pesticides in soil may accumulate in worms via ingesting contaminated soil or the absorption from soil water. Monocrotophos (dimethyl (E)-1-methyl-2-(methylcarbamoyl) vinyl phosphate) is an organophosphorus pesticide that is used to control a broad spectrum of

* Corresponding author. Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu, 608 002, India.

** Corresponding author. Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, 612 001, India.

E-mail addresses: kavitha@annamalai.edu (V. Kavitha), drjgovind1979@gmail.com (M. Govindarajan).



Anti-cancer, anti-biofilm, and anti-inflammatory properties of hen's albumen: A photodynamic approach



Justin Sherly Carolyn^a, Daniel Selva Raj^a, Balasubramanian Malaikozhundan^b,
Marimuthu Govindarajan^c, Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Jamal M. Khaled^{e,*},
Mohammed N. Al-Anbr^e, Ahmed S. Alobaidi^e, Baskaralingam Vaseeharan^{b,*}

^aDepartment of Zoology, Scott Christian College, Nagercoil 629 003, Tamil Nadu, India

^bBiomaterials and Biotechnology in Animal Health Lab., Department of Animal Health and Management, Alagappa University, Science Campus 6thFloor, Karaikudi 630004, Tamil Nadu, India

^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^dDepartment of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^eDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

ARTICLE INFO

Keywords:

SDS-PAGE
Light microscope
CLSM
Clinical bacteria
Biofilm
Hepatic carcinoma

ABSTRACT

The albumen plays a major role in the protection of eggs against microorganisms. It contains an arsenal of natural antimicrobial molecules and antibacterial proteins, including the well-known ovotransferrin and lysozyme, which exert their activities against a range of bacteria. In the present study, the hen's albumen extract treated with the dried insect body of blister beetle *M. pustulata* was assessed for antibacterial, antibiofilm, anti-inflammatory and anti-proliferative activity. The zone of inhibition against Gram positive *E. faecalis* and *S. aureus* was 10.8 mm and 12.1 mm respectively at 100 $\mu\text{g mL}^{-1}$. However, it was 13.6 mm and 15.3 mm for Gram negative *P. aeruginosa* and *P. vulgaris* respectively. The biofilm of tested bacteria was significantly inhibited at 100 $\mu\text{g mL}^{-1}$. The hydrophobicity of bacterial biofilms was considerably condensed after treatment with the hen's albumen extracts at 100 $\mu\text{g mL}^{-1}$. The anti-inflammatory activity of hen's albumen extracts was confirmed by the inhibition of cyclooxygenase (COX) enzyme to 84.91% at 100 $\mu\text{g mL}^{-1}$ with the relative IC₅₀ of 8.26 $\mu\text{g mL}^{-1}$. The albumen extract effectively inhibited the viability (23.61%) of HepG2 hepatic cancer cells at 100 $\mu\text{g mL}^{-1}$. The anti-proliferative activity of the albumen extracts was further revealed by the induction of HepG2 apoptotic cell morphology. This study concludes that the hen's albumen extract treated with *M. pustulata* is a natural therapeutic agent to treat biofilm associated clinical bacteria, inflammations and human hepatic cancer cells.

1. Introduction

Biofilms are bacterial exopolysaccharides that are formed on solid surfaces [1–10]. Inflammation is a biological defense against infection or injury [11]. It helps us to eliminate the foreign bodies or injurious agents. Furthermore, it removes damaged tissues and assists in healing process. However, many acute and chronic human diseases are due to uncontrolled inflammation [12].

Hepatocellular carcinoma (HCC) is the second leading cause of deaths worldwide [13–18]. Hence, the progress of an efficient anti-cancer agent is a vital need for administration of HCC. The study on the potency of animal-born drugs to treat hepatocellular carcinoma has received considerable attention due to their safety compared to

chemotherapeutics.

Since ancient times, insects have been used to cure diseases. In traditional medicine, insect bodies, eggs and secretions was used for different ailments. Recent scientific studies have shown the pharmaceutical applications of insect extractions including anti-bacterial, anti-inflammatory and anti-tumor activities [19]. The blister beetles commonly called as "oil beetles" belong to the family Meloidae. They secrete yellow coloured, pungent smelling oily droplets of haemolymph from the leg joints when disturbed. The exudation contains the toxic material 'cantharidin' or cantharidin [20–24]. Cantharidin and its derivatives have been found to contain anti-tumoral properties against breast, colorectal, melanomas, bladder, pancreatic, hepatic cancers and leukemia [25]. The blister beetle of this study *M. pustulata* are black and

* Corresponding authors.

E-mail addresses: gkhaled@ksu.edu.sa (J.M. Khaled), vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.pdpdt.2019.07.026>

Received 27 June 2019; Received in revised form 23 July 2019; Accepted 29 July 2019

Available online 30 July 2019

1572-1000/ © 2019 Elsevier B.V. All rights reserved.



Facile synthesis of haemocyanin-capped zinc oxide nanoparticles: Effect on growth performance, digestive-enzyme activity, and immune responses of *Penaeus semisulcatus*

Ramachandran Ishwarya^a, Rengarajan Jayakumar^b, Muthukumar Abinaya^a, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Jamal M. Khaled^e, Mohammed N. Al-Anbr^e, Baskaralingam Vaseeharan^{a,*}

^a Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma Colony, Karaikudi 630 004, Tamil Nadu, India

^b Mandapam Regional Centre, ICAR - Central Marine Fisheries Research Institute, Mandapam, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

ARTICLE INFO

Article history:

Received 25 April 2019

Received in revised form 30 July 2019

Accepted 30 July 2019

Available online 31 July 2019

Keywords:

Penaeus semisulcatus

Antioxidants

Zinc nanoparticles

Haemocyanin

ABSTRACT

ZnO nanoparticles (NPs) synthesized using haemocyanin (Hc-ZnONPs) purified from haemolymph of *Penaeus semisulcatus* were characterized using various techniques. HR-TEM and SEM microscopy indicated Hc-ZnONPs had a typical size of 20–50 nm and were spherical. The objective of current investigation was to assess the effects of dietary supplementation of Hc-ZnONPs on the development and activity of digestive and metabolic enzymes, as well as the antioxidant levels in *P. semisulcatus*. Trial basal diets were supplemented with Hc-ZnONPs at rates of 0, 10, 20, 40, 60, and 80 mg kg⁻¹ (dry feed weight) and were fed to *P. semisulcatus* for 30 d. For 60 mg kg⁻¹ Hc-ZnONPs-supplemented feed, significantly ($P < 0.05$) enhanced endurance, development, and activity of the digestive enzyme were observed. The enzymatic antioxidants and metabolic enzymes activities in the muscle exhibited no significant changes when 10–60 mg kg⁻¹ Hc-ZnONPs-supplemented feed was fed to *P. semisulcatus*. Conversely, feeding the *P. semisulcatus* with 80 mg kg⁻¹ Hc-ZnONPs produced a harmful outcome, with significant increase in the enzymatic antioxidants and metabolic enzymes. Consequently, 80 mg kg⁻¹ Hc-ZnONPs was identified as lethal to *P. semisulcatus*. Hence, it is proposed that the diet of *P. semisulcatus* can be supplemented with up to 60 mg kg⁻¹ Hc-ZnONPs for improving the endurance, development and immunity.

© 2019 Published by Elsevier B.V.

1. Introduction

Aquaculture is the most important major food manufacturing sector, after agriculture. The green tiger prawn *Penaeus semisulcatus* is the major species supporting the marine fishery along the southeast shore of India [1]. Micronutrients play crucial roles in the physiological, biological, and immunological functions of organisms. Zn is a necessary mineral for stabilizing cellular membranes and is a trace element in the organs [2]. It is important in numerous metabolic pathways and essential for the development, protein synthesis, carbohydrate metabolism, cell division, and immune systems of all organisms [3–5]. Zn dietary requirements differ among species, and the purposes of Zn for endurance, development, muscle composition, antimicrobial action, immune response, and stress tolerance in several aquatic animals have

been described [6–9]. Zn insufficiency may restrain the role of pointing molecules related to the replication of DNA and organism repair. The assimilation of Zn restrains the reproductive recital, development, and mineral consumption in fish and shellfish [10–12].

Nanoscience is a widespread and multidisciplinary field of study that is currently expanding. The productivity of fish and shellfish businesses can be increased through nanotechnology, by using novel devices to improve the capacity of cultivable animals to uptake nutrients, vaccines, and hormones [13]. Se, Al, Fe, ferrous oxide, Zn, and zinc oxide nanoparticles (NPs) play essential roles in aquaculture processes. Conversely, the surfeit of dietary NPs may be poisonous to aquatic animals [14,15]. The crucial achievement of aquaculture is resolute through the endurance, development and health of the cultivable species. Still, there is requiring of information pertaining to ZnONPs on the fundamental physiology and biochemistry of crustaceans. Green synthesis approaches are ahead attention to avoid the high expenses and practice of toxic chemicals and cruel circumstances for reduction and

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).



New insecticides and antimicrobials derived from *Sargassum wightii* and *Halimeda gracillis* seaweeds: Toxicity against mosquito vectors and antibiofilm activity against microbial pathogens

S. Suganya^a, R. Ishwarya^a, R. Jayakumar^b, M. Govindarajan^{c,d}, N.S. Alharbi^e, S. Kadaikunnan^e, J.M. Khaled^e, M.N. Al-anbr^e, B. Vaseeharan^{a,*}

^a Nanobiosciences and Nanopharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block 6th floor, Burma Colony, Karaikudi 630 004, Tamil Nadu, India

^b Mandapam Regional Centre, Central Marine Fisheries Research Institute, Mandapam, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry, and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

ARTICLE INFO

Article history:

Received 4 April 2019

Received in revised form 23 July 2019

Accepted 2 August 2019

Available online 22 August 2019

Edited by WA Stirk

Keywords:

Arbovirus
Halimeda gracilis
Sargassum wightii
Larvicides
Pesticides
Zika virus

ABSTRACT

Phytochemicals derived from seaweeds offer a novel source of compounds to develop new insecticides and antimicrobials. In the present study, two seaweed species, the brown *Sargassum wightii* and the green *Halimeda gracilis*, were investigated. Extracts were obtained using five solvents of varying polarity (acetone, chloroform, methanol, ethanol, and water). The total phenolic content, total flavonoid content and the antioxidant activities of the extracts were estimated. The ethanol extract from *S. wightii* exhibited more potent antioxidant activity compared to that of the *H. gracilis* extracts. The ethanol extract from *S. wightii* possessed higher values of total phenolics, flavonoids, and antioxidant activity compared to those of the *H. gracilis* ethanol extract. The total phenolic and flavonoid content for ethanol extracts of *S. wightii* and *H. gracilis* was 5.5 mg gallic acid equivalent (GAE)/g, 3.8 mg GAE/g, 10 mg quercetin equivalent (QE)/g, and 8.3 mg QE/g, respectively. The total antioxidant activity of the seaweed extracts was 47 mg ascorbic acid equivalent (AAE)/g for *S. wightii* and 35.9 mg AAE/g for *H. gracilis*. Additionally, the bioactive compounds present in these seaweeds were characterized by FTIR spectroscopy and GC–MS analysis, and the antimicrobial activity of these extracts was then tested against the biofilm formation capabilities of three pathogenic Gram-negative bacteria. The ethanol extracts from *S. wightii* and *H. gracilis* presented the highest inhibitory effect with up to a 40–75% significant reduction in biofilm formation by Gram-negative bacteria. Finally, their insecticidal potential was assessed against late third instar larvae of malaria vectors (*Anopheles stephensi*), dengue and Zika virus vectors (*Aedes aegypti*), and the Japanese encephalitis vector (*Culex tritaeniorhynchus*). LC₅₀ values estimated for the *S. wightii* ethanol extract were lower than 50 ppm against all tested mosquito species. Light microscopy revealed that morphological changes were triggered upon exposure to seaweed extracts. Overall, the present study sheds light on the relevant bioactivity of phytochemicals obtained from two widespread seaweeds, suggesting the possibility that these seaweeds can be used to develop novel and biodegradable pesticides and also antimicrobial drugs that can be used to inhibit the development of microbial biofilms.

© 2019 SAAB. Published by Elsevier B.V. All rights reserved.

1. Introduction


For centuries, marine algae and seaweeds have been used by humans for various purposes that range from food sources to drug development (Mohy El-Din and El-Ahwany, 2016). In particular, seaweeds represent an abundant unexplored source of highly diverse phytochemicals (Smit, 2004) that possess various biological functions such as

antimicrobial and anti-tumor activities (Kolanjinathan et al., 2009; Chakraborty et al., 2010; Manivannan et al., 2011). For example, natural products containing energetic biocompounds that possess effective antibacterial and antioxidant activity have recently been used in aquaculture to prevent diseases (Atanasov et al., 2015; Gopinath et al., 2016).

Seaweeds are exposed to elevated concentrations of oxygen and light intensity. To survive in these challenging environments, seaweeds possess an abundance of antioxidants within their tissues to prevent oxidative damage (Gupta and Abu-Ghannam, 2011; Souza et al., 2011; Chakraborty et al., 2015). Consequently, marine seaweeds provide a

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).



Synthesis and characterization of crustin capped titanium dioxide nanoparticles: Photocatalytic, antibacterial, antifungal and insecticidal activities

Ravichandran Rekha^a, Mani Divya^a, Marimuthu Govindarajan^{b,c}, Naiyf S. Alharbi^d, Shine Kadaikunnan^d, Jamal M. Khaled^d, Mohammed N. Al-Anbr^d, Roman Pavela^e, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Nanobiosciences and Nanopharmacology Division, Department of Animal Health and Management, Alagappa University, Karaikudi 630004, Tamil Nadu, India

^b Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India

^c Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^d Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

^e Crop Research Institute, Drnovska 507, 161 06 Prague, Czech Republic

ARTICLE INFO

Keywords:

Crustin
Nanoparticles
Antibiofilm
Photocatalyst
Vector management
Human parasites

ABSTRACT

Current scenario of bio-nanotechnology, successfully fabrication of ultrafine titanium dioxide nanoparticles (TiO₂NPs) using various biological protein sources for the multipurpose targets. The present research report involves synthesis of TiO₂NPs using antimicrobial peptide (AMP) crustin (Cr). Crustin previously purified from the blue crab, *Portunus pelagicus* haemolymph, by blue Sepharose CL-6B matrix assisted affinity column chromatography. Synthesized Cr-TiO₂NPs was physico-chemically characterized by UV-Visible spectroscopy (UV-Visible), X-ray Diffraction (XRD), Fourier transform infrared spectroscopy (FTIR), High-resolution transmission electron microscopy (HR-TEM) and zeta potential examination. X-ray diffraction analysis for crystalline nature and phase identification of titanium dioxide nanoparticles was absorbed. Functional groups were found through FTIR ranges between 1620 and 1700 cm⁻¹. HR-TEM analysis showed that the synthesized Cr-TiO₂NPs tetragonal shape and sizes ranging from 10 to 50 nm. Finally, the surface charge of the Cr-TiO₂NPs was confirmed through zeta potential analysis. Furthermore, the characterized Cr-TiO₂NPs exhibited good biofilm inhibition against GPB - *S. mutans* (Gram Positive Bacteria- *Streptococcus mutans*), GNB - *P. vulgaris* (Gram Negative Bacteria- *Proteus vulgaris*) and fungal *Candida albicans*. Moreover, photocatalysis demonstrated that the Cr-TiO₂NPs was effectively explored the degradation of dyes. The results suggest that Cr-TiO₂NPs is an excellent bactericidal, fungicidal and photocatalytic agent that can be supportively used for biomedical and industrial applications.

1. Introduction

In recent years, pollution has become a major problem, with pollutants such as wastes and pathogens found in various bodies of water, especially from industrial organic dyes [1]. These dyes are used in textiles, printing, paints and sunscreen materials and are extracted from natural sources and also used extensively in biological stains [2]. Nowadays, chemicals are used to prepare different dyes and they are highly demanded by the textile industry [3]. Hence various dyes or pollutants totally change the land and water ecosystem cause severe health problems for society. The continuous discharge and

accumulation of dye effluents in the water ecosystem inhibits the growth of microorganisms by preventing sunlight [4]. There have been many attempts to reduce water pollution, especially dye effluents, through various physical, chemical and mechanical methods, but less impacts has been obtained, with low efficiency, high time consumption and high cost [5]. Such effluents are difficult to remove from bodies of water; major research is therefore needed to minimize dye effluent pollution in bodies of water.

TiO₂ nanoparticles play a vital role in the dye degradation process because of their efficient dye degradation processes with no contaminants. The unique features of titanium have found various usages,

* Corresponding author.


E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.jphotobiol.2019.111620>

Received 3 July 2019; Received in revised form 6 August 2019; Accepted 27 August 2019

Available online 03 September 2019

1011-1344/ © 2019 Elsevier B.V. All rights reserved.



Synthesis of ZnO nanoparticles using insulin-rich leaf extract: Anti-diabetic, antibiofilm and anti-oxidant properties

Viswanathan Vinotha^a, Arokiadhas Iswarya^a, Rajagopalan Thaya^b, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Jamal M. Khaled^e, Mohammed N. Al-Anbr^e, Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi 630004, Tamil Nadu, India

^b Centre for Animal Science Research and Extension Services, Foundation for Innovative Research in Science and Technology, Kelavannavilai, NGO Colony, Nagercoil, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

ARTICLE INFO

Keywords:

Costus igneus
Biosynthesis
XRD
TEM
ZnO nanoparticles
Biological properties

ABSTRACT

Here, we report the novel fabrication of ZnO nanoparticles using the *Costus igneus* leaf extract. Gas chromatography-mass spectrometry (GC-MS) and proton nuclear magnetic resonance (¹H NMR) spectroscopy to determine the bioactive components present in the plant extract. The synthesis of *Ci*-ZnO NPs (*C. igneus*-coated zinc oxide nanoparticles) was accomplished using a cost-effective and simple technique. *Ci*-ZnO NPs were specified using UV-visible spectroscopy, FTIR, XRD, and TEM. *Ci*-ZnO NPs was authenticated by UV-Vis and exhibited a peak at 365 nm. The XRD spectra proved the crystalline character of the *Ci*-ZnO NPs synthesized as hexagonal wurtzite. The FTIR spectrum illustrated the presence of possible functional groups present in *Ci*-ZnO NPs. The TEM micrograph showed evidence of the presence of a hexagonal organization with a size of 26.55 nm typical of *Ci*-ZnO NPs. The α -amylase and α -glucosidase inhibition assays demonstrated antidiabetic activity of *Ci*-ZnO NPs (74 % and 82 %, respectively), and the DPPH [2,2-diphenyl-1-picrylhydrazyl hydrate] assay demonstrated the antioxidant activity of the nanoparticles (75%) at a concentration of 100 μ g/ml. The *Ci*-ZnO NPs exhibited promising antibacterial and biofilm inhibition activity against the pathogenic bacteria *Streptococcus mutans*, *Lysinibacillus fusiformis*, *Proteus vulgaris*, and *Vibrio parahaemolyticus*. Additionally, the *Ci*-ZnO NPs showed biocompatibility with mammalian RBCs with minimum hemolytic activity (0.633 % \pm 0.005 %) at a concentration of 200 μ g/ml.

1. Introduction

Costus igneus Nak (Family: Costaceae), generally identified as insulin plant or spiral flag. Insulin plant, inhabitant to America, has been recently introduced to India. It is believed that consuming these leaves reduces glucose levels in the blood. The leaves of *C. igneus* were traditionally used by the ethnic community from Kolli hills of Tamil Nadu, India for treating diabetes [1]. Shetty et al. [2] reported that the leaves of *C. igneus* condensed the fasting and postprandial blood sugar levels in dexamethasone-induced hyperglycemia in rats. Kalailingam et al. [3] reported that diosgenin isolated from the *C. igneus* leaf extract acts as an effective hypoglycemic compound. It stimulates the renewal of β -cells in pancreas and pancreatic insulin secretion. Further, diosgenin plays a

role in diminishing cardiovascular risk by reducing LDH release and by upregulating the antioxidant enzymes to scavenge glucose induced free radicals. Diosgenin has physiological effects similar to that of insulin. It is proficient at diminishing blood glucose level via oral administration as compared to its intraperitoneal administration [4].

In the history, the fabrication of nanoparticles has been considered as one of the most dynamic fields and highly preferred in the field of nanotechnology due to their enviable characteristics [5–7]. Among them, metal oxide nanoparticles have gained increased interest and are being incorporated into a diversity of products based on their catalytic capacity, electronic and antimicrobial properties [8,9].

Currently, scientist are spotlight on fabrication of metallic nanoparticles from zinc, gold, silver, platinum, palladium, iron, titanium and

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).



Novel and Facile Synthesis of Sea Anemone Adhesive Protein-Coated ZnO Nanoparticles: Antioxidant, Antibiofilm, and Mosquito Larvicidal Activity Against *Aedes aegypti*

Muthukumar Abinaya¹ · Ravichandran Rekha¹ · Shanthini Sivakumar² · Marimuthu Govindarajan^{3,4} · Naiyf S. Alharbi⁵ · Shine Kadaikunnan⁵ · Jamal M. Khaled⁵ · Ahmed S. Alobaidi⁵ · Mohammed N. Al-Anbr⁵ · Baskaralingam Vaseeharan¹

Received: 3 April 2019 / Published online: 8 May 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

Recently, marine bio-products have provided a new insight on biomaterials for pharmaceutical applications. Herein, the use of adhesive protein in the fabrication of bio-inspired nanomaterials noted to the possible realization of vital biomedical applications. This study illustrates the biosynthesis of ZnO nanoparticles (ZnO NPs) using adhesive protein from sea anemone *Stichodactyla haddoni* (*ShAp*). *ShAp*-ZnO NPs were characterized via SDS-PAGE, UV-Visible, XRD, FTIR, TEM and SAED. The molecular weight of the adhesive protein was determined by SDS-PAGE to be between 36 and 42 kDa. The UV-Vis peak of *ShAp*-ZnO NPs was revealed at 210 nm while XRD exhibited the crystalline nature of *ShAp*-ZnO NPs. FTIR revealed that functional group of *ShAp*-ZnO NPs, which exhibited peaks at 3441.31–1073.36 cm⁻¹. HR-TEM revealed that *ShAp*-ZnO NPs obtained structure were hexagonal with 10 nm diameter. The antioxidant properties of *ShAp*, zinc acetate, and *ShAp*-ZnO NPs were noted at 100 µg/mL. Further, microscopic analysis demonstrated that 50 µg/mL of *ShAp*, zinc acetate, and *ShAp*-ZnO NPs efficiently repressed the biofilm formation of both Gram⁽⁺⁾ and Gram⁽⁻⁾ bacteria. In addition, maximal larvicidal effects were noticed at 100 µg/mL of *ShAp*-ZnO NPs against the 3rd instar of *Aedes aegypti*. Overall, the *ShAp*-ZnO NPs could have entomological and pharmaceutical applications.

Keywords *Stichodactyla haddoni* · Adhesive protein · Antibiofilm · Antioxidant · Insecticide

Introduction

Currently, researchers focusing on bioactive molecules have discovered a wide range of sources of organic compounds from marine ecosystems [1, 2]. Few of these

compounds have been reported to have industrial and toxicological applications; however, most of them have not been extensively studied [3, 4]. In invertebrates, the adhesive proteins from marine organisms in the phylum Cnidaria, including sea anemones, play a vital role in

✉ Baskaralingam Vaseeharan
vaseeharanb@gmail.com

Jamal M. Khaled
gkhaled@ksu.edu.sa


¹ Nano Biosciences and Nano Pharmacology Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block 6th Floor, Burma Colony, Karaikudi, Tamil Nadu 630004, India

² Centre for Animal Science Research and Extension Services, Foundation for Innovative Research in Science and Technology, Kelavannanvilai, NGO Colony Road, Nagercoil, Tamil Nadu, India

³ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, Tamil Nadu 608 002, India

⁴ Department of Zoology, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu 612 001, India

⁵ Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia



Toxicology

Chronic exposure of *Oreochromis niloticus* to sub-lethal copper concentrations: Effects on growth, antioxidant, non-enzymatic antioxidant, oxidative stress and non-specific immune responses

Narayanan Gopi^a, Sekar Vijayakumar^a, Rajagopalan Thaya^b, Marimuthu Govindarajan^{c,d},
 Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Jamal M. Khaled^e, Mohammed N. Al-Anbr^e,
 Baskaralingam Vaseeharan^{a,*}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Foundation for Innovative Research in Science and Technology, Kelavannanvilai, NGO Colony Road, Nagercoil, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia



ARTICLE INFO

Keywords:

Fish
 Copper
 Chronic toxicity
 Biomarkers
 Oxidative stress
 Immuno-toxicity

ABSTRACT

Background: Trace elements of copper (Cu) are one of the main forms of ecological noxious waste in freshwater systems that affect the survival and development of organisms. The objective of the current study was to investigate the effects of chronic exposure to Cu on the growth, oxidative stress, immune and biochemical response in the Nile tilapia, *Oreochromis niloticus*.


Methods: Three groups of *O. niloticus* were tested as follows; the first group was used as the control (not treated with Cu in water), while the 2nd and 3rd groups were exposed to (low) 40 µg L⁻¹ and (high) 400 µg L⁻¹ concentrations of Cu added to water, respectively. The duration of the experiment, which was conducted in triplicate, was 60 d. End points were evaluated on days 30 and 60. Following 30 d and 60 d of exposure to Cu, the fish were removed from experimental tanks to determine growth. Consequently, blood samples were collected from caudal veins at the end of the trial period (30 d and 60 d) and serum was separated to evaluate different immunological parameters, such as lysozymes (LYZ), respiratory burst activity (RBA) and myeloperoxidase (MPO). Gill and liver tissues were collected for evaluation of Cu and certain biochemical parameters as follows: antioxidant enzymes such as superoxide dismutase (SOD), catalase (CAT), glutathione peroxidase (GPx) and glutathione-S-transferase (GST); non-enzymatic antioxidants such as glutathione (GSH) and metallothionein (MT), and oxidative stress indicators such as malondialdehyde (MDA) and protein carbonyl (PCO). The results pertaining to treatments and the control were compared using two-way ANOVA and Tukey's HSD test. The level of significance was set at $P \leq 0.05$. Data were expressed as mean \pm SD.

Results: Chronic exposure to Cu did not induce any mortality in fish during the test period. However, following exposure to Cu, growth of fish in the exposed groups was affected more than that in the control group (unexposed to Cu). In addition, accumulation of Cu in the liver tissue was higher than that in the gill tissues of fish exposed to Cu, compared to that in the control. Gill and liver tissues of Cu-exposed fish showed a significant ($P \leq 0.05$) reduction in the activities of the antioxidant enzymes, SOD, CAT, GPx, and GST, compared to those of unexposed fish. Non-enzymatic antioxidants, GSH and MT, in gill and liver tissues were significantly increased ($P \leq 0.05$) in fish exposed to both concentrations of Cu, compared to those in unexposed fish. Oxidative stress indicators, MDA and PCO in gills and liver of Cu-exposed fish was significantly ($P \leq 0.05$) at both tested concentrations, when compared to control group. Non-specific immune response of LYZ, RBA, and MPO activity in serum decreased significantly ($P \leq 0.05$) in Cu-exposed fish, compared with that of unexposed fish.

Conclusion: Overall, the present results highlighted that chronic exposure to Cu ions may exert a strong effect on the antioxidant and immune responses of *O. niloticus*. Changes in antioxidant enzymes, oxidative stress effects and immune parameters during post-chronic metal exposure may indicate the potential of these parameters as biomarkers of metal toxicity in aquatic ecosystems.

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).



Microbial exopolymer-capped selenium nanowires – Towards new antibacterial, antibiofilm and arbovirus vector larvicides?



Muthukumar Abinaya^a, Baskaralingam Vaseeharan^{a,*}, Ravichandran Rekha^a, Sivakumar Shanthini^b, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^{e,*}, Shine Kadaikunnan^e, Jamal M. Khaled^e, Mohammed N. Al-Anbr^c

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma Colony, Karaikudi 630004, Tamil Nadu, India

^b Centre for Animal Science Research and Extension Services, Foundation for Innovative Research in Science and Technology, Vallankumaran vilai, NGO Colony road, Nagercoil 629002, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Keywords:

Dengue
Lymphatic filariasis
Microbial pathogens
Nanobiotechnology
Nanoparticles, Zika virus

ABSTRACT

Arboviral diseases and microbial pathogens resistant to commercially available drugs are on the rise. Herein, a facile microbial-based approach was developed to synthesize selenium nanowires (Se NWs) using microbial exopolymer (MEP) extracted from the *Bacillus licheniformis* (probiotic bacteria). MEP-Se NWs were characterized using UV–Visible, XRD, FTIR, HR-TEM, FE-SEM and EDX. An UV–Visible peak was detected at 330 nm while XRD spectrum data pointed out the crystalline nature of MEP-Se NWs. FTIR spectrum revealed functional groups with strong absorption peaks in the range 3898.52–477.97 cm^{-1} . FE-SEM and HR-TEM revealed that the obtained structures were nanowires of 10–30 nm diameter. Se presence was confirmed by EDX analysis. MEP-Se NWs at 100 $\mu\text{g}/\text{ml}$ highly suppressed the growth of both Gram⁽⁻⁾ and Gram⁽⁺⁾ bacteria. Further, microscopic analysis evidenced that 75 $\mu\text{g}/\text{ml}$ MEP-Se NWs suppressed biofilm formation. Hemolytic assays showed that MEP-Se NWs were moderately cytotoxic. In addition, LC_{50} values lower than 10 $\mu\text{g}/\text{ml}$ were estimated testing MEP-Se NWs on both *Aedes aegypti* and *Culex quinquefasciatus* 3rd instar larvae. Morphological and histological techniques were used to elucidate on the damages triggered in mosquito tissues, with special reference to midgut, post-exposure to MEP-Se NWs. Therefore, based on our findings, MEP-Se NWs can be considered for entomological and biomedical applications, with special reference to the management of biofilm forming microbial pathogens and arbovirus mosquito vectors.

1. Introduction

Mosquito (Diptera: Culicidae) vectors pretense a key threat to millions of people worldwide in tropical and subtropical areas [1–3]. *An. stephensi*, *Ae. aegypti* and *Cx. quinquefasciatus* are the most important vectors of numerous pathogens and parasites clerical for enormous mortality and morbidity worldwide including malaria, dengue and filariasis, respectively [4]. In particular, arboviral vector threads have recently become more common in neglected geographical regions with over 2.5 billion peoples at risk [5]. The WHO estimates 50–100 millions of dengue infections worldwide every year [6]. However, the utilize of synthetic insecticides to control mosquito larvae populations led to

resistance, high operational costs & adverse environmental effects. Therefore, the development of ecofriendly control tools is an important public health challenge [7–9].

In this scenario, the biosynthesis of one-dimensional nanomaterials, such as nanowires, have generated considerable attention because of their distinctive physicochemical properties, which remarkably differ from those of bulk materials, along with their applicability in nanodevices [10–13]. Although metals and semiconductors have been extensively investigated, the potential of metalloids such as selenium (Se) are yet to be analyzed [14]. Selenium is particularly interesting as it is a crucial micronutrient for numerous organisms. It is a fundamental co-factor of glutathione peroxidase, formate dehydrogenase and

* Corresponding authors.


E-mail addresses: vaseeharanb@gmail.com (B. Vaseeharan), nalharbil@ksu.edu.sa (N.S. Alharbi).

<https://doi.org/10.1016/j.jphotobiol.2019.01.009>

Received 13 December 2018; Received in revised form 11 January 2019; Accepted 16 January 2019

Available online 18 January 2019

1011-1344/ © 2019 Elsevier B.V. All rights reserved.



Bioinorganic chemistry

Enhanced antibacterial activity of hemocyanin purified from *Portunus pelagicus* hemolymph combined with silver nanoparticles – Intracellular uptake and mode of action



Ramachandran Ishwarya^a, Baskaralingam Vaseeharan^{a,*}, Sivakumar Shanthini^b,
Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^{e,*}, Shine Kadaikunnan^e, Jamal M. Khaled^e,
 Mohammed N. Al-anbr^e

^aCrustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma Colony, Karaikudi, 630 004, Tamil Nadu, India

^bCentre for Animal Science Research and Extension Services, Foundation for Innovative Research in Science and Technology, Kelavannanvilai, NGO Colony Road, Nagercoil, Tamil Nadu, India

^cUnit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar, Tamil Nadu, 608 002, India

^dDepartment of Zoology, Government College for Women, Kumbakonam, 612 001, Tamil Nadu, India

^eDepartment of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

ARTICLE INFO

Keywords:

Bacterial biofilm
 DNA fragmentation
 Reactive oxygen species
 Malondialdehyde
 Membrane damage
 Nucleic acid leakage
 Peroxidase
 Superoxide dismutase

ABSTRACT

Recently, biogenic nanoparticles have been considered promising candidates for manufacturing antibacterial nanodrugs. Here, we synthesized AgNPs using the crab-borne antibacterial agent hemocyanin and assessed the antibacterial action against several pathogenic bacteria. In this study, the crustacean immune protein hemocyanin (*Pp*-Hc, 78 kDa) purified from *Portunus pelagicus* hemolymph was used to fabricate silver nanoparticles. Characterization of hemocyanin-fabricated AgNPs (*Pp*-Hc AgNPs) were achieved using ultraviolet-visible spectrophotometer, X-ray powder diffraction (XRD), Fourier-transform infrared spectroscopy (FTIR), high-resolution transmission electron microscopy (HR-TEM), and energy-dispersive X-ray spectroscopy. The antibacterial efficacy of AgNO₃, *Pp*-Hc and *Pp*-Hc AgNPs was compared by growth inhibition, antibiofilm and live and dead analyses. Based on the results, *Pp*-Hc AgNPs was more efficient than *Pp*-Hc and AgNO₃ against pathogenic bacteria. Mechanistic analysis revealed membrane damage and reactive oxygen species (ROS) generation, suggesting that *Pp*-Hc and *Pp*-Hc AgNPs rely to similar modes of action. Intracellular protein molecules and nucleic acid leakage confirmed that *Pp*-Hc AgNPs increase membrane permeability, leading to cell death. Based on our results, capping of the exterior surface of nanoparticles with antimicrobial crab-borne peptides, such as *Pp*-Hc, improves their functions as potential agents against bacterial diseases, which may be useful in clinical applications.

1. Introduction

Antibiotics are highly important in the fight against bacterial infections, to reduce the morbidity and mortality of humans, as well as in livestock and pets worldwide. However, in the past few years, an increase in antibiotic resistance of microorganisms – largely due to the overuse of antibiotics – has become a severe problem worldwide [1–3]. An approach for overcoming this problem is to use novel, low-cost, and efficient antimicrobial agents that have biological ability to reduce the rate and spread of resistance development of target microorganisms [4,5]. An increased incidence of infections caused by antibiotic-

resistant microbial strains expanded the attention about antimicrobial properties of silver and its composites [6,7]. Silver has been used as an antimicrobial complex since before the beginning of antibiotics in the 1940s, and is currently used in a broad array of medicinal purposes because of its antibacterial effects and relatively small toxicity to human cells [8–10]. However, the cost burden and current identification of silver-resistant bacteria pointed out the inadequate use of raw silver as possible alternative to broad-spectrum antibiotics [11].

Nanotechnology has rapidly advanced in the current century because of the stunning potential of nanomaterials, with special reference to nanosize, which radically modifies their physical, chemical, and

* Corresponding authors.

E-mail addresses: vaseeharanb@gmail.com (B. Vaseeharan), nalharbil@ksu.edu.sa (N.S. Alharbi).

<https://doi.org/10.1016/j.jtemb.2019.03.005>

Received 27 September 2018; Received in revised form 6 March 2019; Accepted 9 March 2019

0946-672X/ © 2019 Elsevier GmbH. All rights reserved.



Facile and Cost-Effective Ag Nanoparticles Fabricated by *Lilium lancifolium* Leaf Extract: Antibacterial and Antibiofilm Potential

Mysoon Al-Ansari¹ · Noorah Alkubaisi¹ · Kasi Gopinath² · Viswanathan Karthika² · Ayyakannu Arumugam³ · Marimuthu Govindarajan^{4,5}

Received: 8 April 2019 / Published online: 30 April 2019
© Springer Science+Business Media, LLC, part of Springer Nature 2019

Abstract

In the present study one-pot green fabrication of silver nanoparticles (Ag NPs) with aqueous leaf extract of *Lilium lancifolium*. Synthesized Ag NPs were illustrated by UV–Vis, FT-IR, XRD, AFM and TEM analysis. A surface plasmon resonance peak was noticed at 411 nm in UV–Vis analysis. XRD patterns of Ag NPs showed high crystalline nature with face centered cubic structure. AFM and TEM analysis of Ag NPs exhibit special and anisotropic morphology with accurate particle size about 10–70 nm and average mean value of 45 nm. Ag NPs were tested for antibacterial and anti-biofilm activity against Gram positive and Gram-negative bacterial strains. The exposure to 25 µL exhibited better antibacterial activity and 50 µL significantly control the bacterial biofilm growth. Overall, the proposed green synthesis method of Ag NPs is facile, convenient, less time consuming and environmentally safe.

Keywords *Lilium lancifolium* · Green synthesis · Nanoparticle · AFM and TEM · Antibacterial and antibiofilm activity

Introduction

Since the concept of nanotechnology became famous in the early 1990s, scientists have been developing new techniques by incorporating nanoparticles and nanomaterials into various fields of science. The concept of nanomaterials, although new, has been a part of our civilization since a

very long time. Silver nanoparticles (Ag NPs) have been taken attention due to their special properties of size, shape, and self-assembly. They are commonly used in wide range of industries like pharmacology [1, 2], biomedical devices [3, 4], textiles [5, 6], water purification [7, 8] and cosmetics [9, 10]. Many methods are available for synthesis of Ag NPs which include physical, chemical and biological synthesis. In which, biological means of synthesis gained good-attention compared to other two methods due to low cost, simple, environmentally friendly and efficient substitute for the mass production of nanoparticles [11]. In biological methods, Ag NPs are synthesized via using fungus [12, 13], microbes [14, 15] and plants [16–22]. Compare to the fungus and microbes, plants are promising source for nanoparticles production. The advantages of the plant-mediated synthesis of nanoparticles include cost efficient, large-scale production and require low maintenance [23–27]. Important point of using plants in nanoparticles production instead of fungus or bacteria is the lack of pathogenicity [28]. In the past researches, silver nanoparticles are synthesized using extracts from different plant materials such as *Azadirachta indica* [29], *Catharanthus roseus* [30], *Origanum vulgare* [31], *Buddleja*

✉ Mysoon Al-Ansari
soona70@yahoo.com

✉ Marimuthu Govindarajan
drgovind1979@gmail.com

¹ Department of Botany and Microbiology, Female Campus, College of Science, King Saud University, Riyadh, Saudi Arabia

² Department of Packaging, Yonsei University, 1 Yonseidaegil, Wonju-si, Gangwon-Do 26493, South Korea

³ Department of Botany, Alagappa University, Karaikudi, Tamil Nadu 630 003, India

⁴ Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalinagar, Tamil Nadu 608 002, India

⁵ Department of Zoology, Government College for Women, Kumbakonam, Tamil Nadu 612 001, India

**ANALYSE OF SELECTED HEAVY METAL CONTAMINATION IN
THE MUDIKONDAN RIVER NANNILAM THIRUVARUR DISTRICT
TAMIL NADU INDIA**

Kavitha V.¹, Anandhan R.*¹, Sumathi R.², Balasubramanian.R.³ and S. Baskarsanjeevi⁴

¹Assistant Professor, Department of Zoology, Annamalai University, Annamalai Nagar. 608 002. Tamil Nadu, India.

²Research Scholar, Department of Zoology, Thiru Vi Ka Arts College, Thiruvavur. Tamil Nadu, India.

³Assistant Professor, Center Advanced Study in Marine Biology, Faculty of Marine Sciences, Annamalai University, Parangipettai- 608 502, Tamil Nadu, India.

⁴Assistant Professor, Department of Zoology, Government Arts College, Kumbakonam. 612 002.

ABSTRACT

The present short-review briefly summarizes the status of heavy metal in Cauvery River and its sources and the status and effect of heavy metals in the river sediments and water. The difference in the heavy metals in different parts of the pathways of the river Cauvery is provided in addition to the pollution level and the effect of polluted water or enriched heavy metals on living systems. The pollution status and heavy metal contaminants level varies in water and in sediments with reference to different locations. The rapid population growth along the Mudikondan River has necessitated proper conservation and efficient utilization of freshwater bodies for sustainable development. This is necessary because there has been accelerated deterioration of

water quality and also because of increased domestic, municipal and agricultural activities. Effluent discharge, urbanization and deforestation are the main causes of environmental degradation within the catchment.

KEYWORDS: Cauvery River, Contamination, Heavy metals, Pollution and Sediments.

Article Received on
21 May 2019,

Revised on 12 June 2019,
Accepted on 01 July 2019,

DOI: 10.20950/wjpr20198-15384

***Corresponding Author**

Dr. Anandhan R.

Assistant Professor,

Department of Zoology,

Annamalai University,

Annamalai Nagar.608 002.

Tamil Nadu, India.



Assessment of Flood hazard prone areas in parts of Thanjavur District, South India using FIGUSED Method

Sumatra A¹ and ²Neelakantan R

¹Research Scholar, ²Professor and Head,

Department of Industries and Earth Sciences,

Tamil University, Thanjavur, Tamil Nadu, India

Abstract: Thanjavur district is primarily an agricultural area enclosed with irrigation system of the Cauvery in Tamil Nadu. But during the monsoon period flooding has been disrupted the areas and directly effected the soil erosion in those areas. The flood hazard assessment is one of the important task to understand the soil erosion of the area due to natural hazards. In this context, the present study to assess flood hazard prone areas using FIGUSED method adopted with remote sensing data and GIS technology. In FIGUSED method, seven parameters has been used such as flow accumulation, distance from the drainage network, elevation, and use, rainfall intensity and geology. The relative importance of each parameter for the occurrence and severity of flood has been connected to weight values. These values are calculated following a normalized weight parameter rating method. According to their weight values, information of the different parameters is superimposed, resulting to flood hazard mapping. The Flood Hazard Index (FHI) has been defined and a spatial analysis in a GIS environment has been applied for the estimation of its value. The historical flood events, accuracy and sensitivity analysis not examined in this study and it is only understand the flood hazard area. The study revealed that about 12% of the area falls under very high flood hazard and 26% high to moderate flood hazard zones and indicates that immediate attention have to be taken and protect the agricultural lands for the sustainable development of those areas.

Index Terms – Normalized weight, flood prone area, GIS analysis, flood hazard area, South India

1. INTRODUCTION

Flood is a major natural hazard with often immeasurable impact, affecting annually 170 million people (Kowalzig, 2008). Therefore, flood risk management needs to overcome national borders, geographic location and socio-economic limitations (Degiorgis et al., 2012). Flood risk management is usually divided into flood risk assessment and flood risk mitigation (Schanze et al., 2006). From sustainable development point of view, the flood hazard management is very essential for future (Schober et al., 2015). Tehrany et al., 2013 have studied 10 parameters with the relative importance of each parameter defined following a statistical analysis. During the JAL cyclone event (November to December 2010), severe floods, occurred in Thanjavur district and Thanjavur taluk which spreads at the margin of Cauvery River was one of the affected places. Cyclones ravage the district once in 3-5 years, during north east monsoon, resulting in flood and crop damage. During 1982-83, 1990-91 and 1992-93 cyclones of high intensity have affected the district. Every year monsoon cyclone flood and drought situation may occur during Rabi season which may also considerably affect the paddy production in Thanjavur district.

The application of GIS-based multi-criteria analysis in the context of flood risk assessment was rare until 2000. Black and Burns (2002) have studied the changes in the estimation of flood risk on Scottish rivers with time by re-analyzing flood records. An early attempt to use GIS on water-related hazards has been presented in Meja-Navarro et al. (1994). The present article deals with the first element of flood risk management, i.e. the definition of flood hazard areas in a specific region. The present study methodology adopted based on the Kazakis et al (2015) FIGUSED methods to identify flood hazard zones in parts of Thanjavur and the output of the results is very useful to the farmer and planners for the agricultural purposes in the study area.

2. STUDY AREA

Thanjavur district lies between 9° 50' and 11° 25' North latitude and 78° 45' and 79° 25' East longitude (Fig 1). It is bounded on the North by Thiruchirapalli and Cuddalore districts, on the East by Tiruvarur and Nagapattinam districts, on the South by Palk Strait and Pudukkottai district and on the west by Pudukkottai district and Tiruchirapalli districts. Total geographical of the study area is 1697 sq.km. The mean maximum temperature was 37.48°C during May – July. Similarly, the mean minimum temperature was 20.82°C during November-January. The north east monsoon provides much rainfall with 545.7 mm and 953.2 as normal and actual rainfall respectively, while southwest monsoon provides 342 and 303.1 mm as normal and actual rainfall respectively. The total population of Thanjavur district is 22,16,138. Thanjavur district stands unique from time immemorial for its agricultural activities and is rightly acclaimed as the granary of South India lying in the deltaic region of the famous river Cauvery and criss-crossed by lengthy network of irrigation canals. Therefore this coastal district abounds in green paddy fields, tall coconut groves, vast gardens of mango and plantain trees and other verdant vegetation.

AN EVALUATION OF MATERNAL HEALTH CARE SERVICES IN PRIMARY HEALTHCARE CENTERS (PHC) IN THANJAVUR DISTRICT TAMIL NADU

S. GEETHA & R. MANIYOSAI

Assistant Professor, Department of Geography,

Government College for Women, Kumbakonam, Tamilnadu, India

ABSTRACT

This study covers the services of Primary Healthcare Centers in Thanjavur District. It examines the visitors' view of how they utilized these PHC Services. The study concentrates on How PHC helps to the rural population. Moreover, what are factors controlling PHCs. To identify the challenges that prevent PHCs from utilizing resources, the study has been conducted in All PHCs of Thanjavur district. A survey was done on the PHC users and asking about their services. The main purpose of this study is to evaluate the current services of PHCs in Thanjavur district from the perspective of patients. A Questionnaire created and used to collect the information using by direct investigation method. The questions asked to mother and collect feedback directly. From the Findings, the PHCs of Thanjavur district is good in providing health services. It observed that the people effectively utilizing the services of PHC and Health sub-centers for vaccination and emergency basis. They have awareness about the utilization of PHC services, drugs, vaccinations and Government schemes. The contribution of village nurses is to improve the people's knowledge about PHCs Services in their area. Frequent awareness programs is required to reach government schemes among users.

KEYWORDS: *Primary Health Center, Maternal Health, Patient, Village Nurses & Respondents*

Received: Nov 24, 2018; **Accepted:** Dec 14, 2018; **Published:** Jun 21, 2019; **Paper Id.:** IJEEFUSAUG20192

INTRODUCTION

Nowadays, society is decided to create awareness between people like "A Healthy Society is an indication of Wealthy Nation." A primary health center provides health care services for economically backward people. Primary health centers started from the ideal of healthcare that was received in the declaration of the International Conference on primary health care held in, "Alma Ata," Kazakhasta in 1978. According to WHO (World Health Organization) goal is "Health for all." The ultimate goal of PHC is to give better health services to all. PHC is a federally qualified center that provides primary health care services, maternal, and healthcare, including family planning, education on health problems. It also explores how to control them and adequate and safe water supply and sanitation.

STUDY AREA

Thanjavur district is located between 10° 47' to 13.19" North latitude and 79° 8' to 16.17" East longitude. It is bounded by the districts on the north by Thiruchirapalli and Cuddalore, on the east by Tiruvarur and Nagapattinam, on the south by palkstrait and Pudukkotai and Thiruchirapalli districts. The total geographical area of Thanjavur district is 3,602.86 sq.km. Thanjavur district is divided into Eight taluks for administrative purposes

Urban Population Growth Using GIS in Kumbakonam Municipality, Tamilnadu, India

¹Kalyani V, ²Rajakokila K, ³Neelakantan R,

¹ Asst Professor, ² Guest Lecturer, ³ Professor and Head,

¹ Department of Geography, Govt. College for women(A), Kumbakonam, India

² Department of Geography, Bharathidasan University Tiruchirappalli, India

³ Department of Industries and Earth Sciences, Tamil University, Thanjavur, India.

¹ Email: kalyanisatheesh@gmail.com, ² Email: k.rajakokila@gmail.com, ³ Email: neels2004@gmail.com

Abstract

Urbanization is the processes by which villages turn into towns and towns develop into cities. It is a process and can be interpreted that there is an increasing economic specialization and advancing technology which results in the accumulation of people. Kumbakonam Municipality, located in South India has experienced rapid urbanization in the past two decades. Urban population growth occurs in the study area with major processes led to a slow by slow transformation of agricultural land into urban land-uses leading to increase in more built-up area. This study attempt to measure and analyze urban growth in the study area using temporal data changes for years 2000 and 2015 using ARC GIS.

Keywords: Urban Population Growth, Built-up area Mapping, Buffer Analysis and GIS

1.0 Introduction

Urbanization is the processes by which villages turn into towns and towns develop into cities. Two major causes are responsible for the growth of urbanization and its rapid spread from the beginning of the 21st century. The first is development of technology and rapid industrialization, and the second is unplanned migration to large cities from the rural areas. Urbanization is two way processes because it involves not only movement from village to cities and change from agricultural activities to commercial and service activities but it also involves change in the attitudes and behavior patterns of migrant peoples. Man has always tried to improve his lifestyle everybody, because the new invention of instruments helps to reduce the working time which in turn brought changes in human lifestyle change. Change detection analysis is an important key to understanding changes took place over time. Towns and cities have acted as focal points in the cultural landscape of world. Areal differentiation of any phenomenon became not only complex but also dynamic posing challenges to geographical research.

1.1 Literature Review

A STUDY ON MATERNAL AND SOCIO - ECONOMIC FACTORS INFLUENCING INFANT BIRTH WEIGHT IN THANJAVUR DISTRICT, TAMIL NADU

S.Geetha and R.Maniyosai

1. Research Scholar, Department of Geography, Government Arts College (A), Kumbaonam.
2. Asst.Professor of Geography, Government Arts College (A), Kumbaonam.

ABSTRACT

Birth weight of a baby is important for their health. Low birth weight is one of the main reasons for infant mortality. Birth weight of less than 2500g is defined as Low birth weight (WHO). A study conducted on how socio-economic and demographic factors affecting the birth weight of a baby. It was conducted on Primary Health care centers of Thanjavur District. The study has taken into 805 new born babies. 198 low birth weight babies, 482 Normal birth weight and 125 above normal weight babies taking into consideration. Statistical Analysis and Chi Square tests used for Analysis. Occupation of Husband, Income of Husband, Young age marriage, low income and employment of pregnant women are taken as the main factors for low birth weight. From the study economic, social and demographic factors have not significant effect on birth weight. Maternal health care factors have an significant effect on birth weight.

Key words: Birth weight, maternal factors, health of mothers.

1. INTRODUCTION

WHO defines Birth weight of less than 2500g is Low birth weight. Birth weight is a major indicator for reproductive health of a woman. Low weight babies can face many health issues like infection, breathing and feeding, inability to maintain body temperature and so on. To grow a healthy baby, mothers need good nutrition and rest, adequate antenatal care, and a clean environment (UNICEF). UNICEF estimates almost every third new born baby in India is Low birth weight. According to NFHS 4 Report 20% of low birth weight babies births to mothers from the poorest families. A. F. Roche and S. S. Sun et.al (2003) Socio-economic state is a concept devised to measure some aspects of education, occupation, and social prestige of a person or a social group. Ashraf Mohammadzadeh et.al. (2010). The variables which show the most significant association were birth weight, sex, economic status, and educational level of the

parents. In this study, maternal and paternal education has significant influence on weight although this effect on height was not significant.

2 AIM

The Aim of the study to identify what are the factors affecting the birth weight of the new born babies.

3. REVIEW OF LITERATURE

Birth weight of the newborn is believed to be influenced by a number of factors. The maternal age and parity have a significant effect on birth weight, as shown by Dhall & Bagga (1995). Jolanta Dickute et al (2004), low education, unstable marital status and low income are the major factors for low birth weight. Maternal unfavorable socio-economic factors increased the risk to deliver low birth weight baby. Joshi H.S et.al (2010). The main factors significantly associated with LBW were maternal education, stature, age at delivery; short inter pregnancy interval, inadequate antenatal care, and per capita income of family. Shabbeer Ahmed et.al (2016) The main factors which were significantly associated with LBW were education, community, burden of labor and per capita income of family. Rashidul Alam Mahumud et.al (2016). explained that delayed conception, advanced maternal age, and inadequate ANC visits had independent effects on the prevalence of LBW. Prerna Bansal et.al (2018) Socio-cultural and maternal risk factors like rest received in the afternoon during pregnancy, dietary intake during pregnancy and period of gestation were found to be significantly associated with low birth

RISK MANAGEMENT IN EQUITY INVESTMENTS – A STUDY WITH REFERENCE TO THANJAVUR DISTRICT

K. KULANJIAMMAL

Ph.D Research Scholar & Assistant Professor,
Dept. of Commerce, Govt. College for Women (Autonomous), Kumbakonam

Dr. R. REVATHI

Assistant Professor, Dept. Of Commerce,
Govt. College for Women (Autonomous), Kumbakonam

ABSTRACT

The researcher studied risk management of investors in equity investments in Thanjavur district. The sample size was 384 and primary data were collected from them using well-structured questionnaire. The researcher applied simple percentage and correlation analysis as statistical tools to analyze the data. The study found that respondents in the study area management risk while making investment decisions efficiently. The respondents considered risk before making investment decisions. Majority of the respondents had wider portfolio in order to reduce risk, they were cautious about sudden ups and downs in the market, they held shares when prices were increasing, they kept distance when the market was not good, they avoided shares which did not perform good and they disinvested when they determine that bearish market begins. But majority of the respondents did not consider inflation rate and did not discuss with experts about risk before making investment decisions. Educational level of the respondents had significant positive relationship with risk management. Similarly monthly trading volume traded by them, investment frequency and investment experience had significant positive relationship with risk management in the study area. Percentage of total investment to stock market investment did not have significant relationship with risk management.

Key terms: Stock market, equity, investment, risk and inflation.

INTRODUCTION

People who have excess of income over their expenditure are intended to save their money. For this purpose they seek for best investment type among availability of many number of investment avenues. Type of investment is chosen by the investor is based on their need and their attitude towards investments. The rate of return on investments varies based on risk vested in the investments. Generally risk and returns have direct relationship, i.e, if there is high risk in an investment there is high rate of return on it and if the risk level is low in an investment there is low rate of return on such investment. Investors, who have high risk taking capacity, will go for type of investments which have more risk. In recent decades stock market is becoming as one of the most attractive investment

A STUDY ON FACTORS INFLUENCING STOCK MARKET INVESTMENTS IN THANJAVUR DISTRICT – A SPECIAL REFERENCE TO RETAIL INVESTORS

K. KULANJIAMMAL

Ph.D Scholar

Dr. R. REVATHI M.Com.,M.Phil.,Ph.D

Assistant Professor of Commerce

Govt. College for Women (Autonomous)

Kumbakonam- 612 001

Abstract

Investments in stock market by retail investors are keeping on increasing during recent years. Many factors are influencing the investors to make investments in stock market. Hence the researcher studied the selected factors influencing stock market investments in Thanjavur district. For this purpose the researcher collected primary data from 384 respondents in Thanjavur district using well structured questionnaire. The researcher applied simple percentage and chi-square test as statistical tools to analyse the data. The researcher found that the factors influence of brokers / sub-brokers, expectation of high returns, continues bullish market, having risk taking capacity and influence of advertisements influenced more on the investments in stock market. It was also found that economic and financial factors and market factors influenced the respondents more on the investments in stock market. The study also found that there was significant association in factors influencing investments in stock market with gender, age, annual savings of the respondents and experience of the respondents in stock market investments. No significant association was found in factors influencing stock market investments with education, income and occupation of the respondents.

Key words: Stock market, equity, factors, return and investments.

Introduction

Stock market is a place where shares and securities are traded. Stock markets are exchanging the securities from a person to another person which are already listed in their stock exchange. In other words, stock markets are dealing with second securities. Stock market is playing as a most preferable investment avenue for all type of investors such as retail investors, domestic institutional investors and foreign institutional investors. Investments in stock market gives high rate of return compared to other type of investments in India. Apart from this feature many other features of stock market investments such liquidity, capital appreciation etc are also attracting the investors towards investment in stock

A Proposed Methodology for the Recognition and Classification of the Ancient Scripts from Epigraphy Images Using Histogram of BRISK Orientation Sign Code

S. Sundareswari, Research Scholar, Research and Development Centre, Bharathiar University, Coimbatore, Tamil Nadu, India.
E-mail: profresearch2014@gmail.com

Dr. I. Kaspar Raj, Director, Computer Centre, The Gandigram Rural Institute Deemed University, Gandigram, Dindigul, Tamil Nadu, India. E-mail: kasparraj@gmail.com

Abstract— The extraction of ancient Tamil scripts from epigraphy images is a challenging problem in the pattern recognition area. Now-a-days, modern readers are faced with many difficulties to interpret ancient script from epigraphy images. In this paper, we present a proposed methodology to extract ancient scripts from epigraphy images and also a comparative study of previous work has been done. In this work, we introduce Histogram of BRISK (Binary Robust Invariant Scalable Key points) orientation sign code to extract the ancient scripts from epigraphy images easily and accurately.

Keywords—BRISK (Binary Robust Invariant Scalable Key Points), Histogram of BRISK Orientation Sign Code, Zernike Moment Descriptor, LOMO (Local Maximal Occurrence), Random Forest, K-Nearest Neighbor.

I. Introduction

An extraction of characters from document image is an easy process. Many techniques are available for the extraction of characters from document image. It is very difficult to extract characters from epigraphy images. The ancient script in epigraphy images is not having the same shape and similarity. Their writing style and size of the characters are different.

For these reasons, it is not easy to extract ancient scripts from epigraphy images. i) In preprocessing, Gabor filter and active contour are used to enhance the images. ii) After the preprocessing, BRISK (Binary Robust Invariant Scalable keypoints) is used to identify the valid points in the ancient character. Next, the proposed technique Brisk histogram of sign code is used to identify the horizontal and vertical valid points in each and every ancient characters.

The combinational approach of Zernike moment descriptor, LOMO (Local Maximal Occurrence) and Histogram of BRISK orientation sign code are used as feature descriptors. iii) In classification, KNN (K-Nearest Neighbor) and RF (Random Forest) are used as classifiers and to select best classifier for this research work. This proposed method helps us to get good results and extracts ancient scripts from epigraphy images accurately and clearly.

This paper contains the following section 1 presents the overview of the work, section 2 explains the details of the BRISK technique, section 3 presents the details of the algorithm, section 4 discuss the details of feature extraction techniques, section 5 discuss the various types of classifiers, section 6 contains results and discussion, section 7 concludes the performance of the research.

II. BRISK (Binary Robust Invariant Scalable Key Points)

BRISK (Binary Robust Invariant Scalable Key points) proposed by Stefan Leutenegger [1] is used to detect the corners in scale space. It involves basically two steps: i) Scale Space Key point detection ii) Key point description. Key point detection means it detects the interest points in image and scale dimensions using a prominent criterion. The computation is done by identifying the key points in octave layers of the image pyramid as well as in layers in-between. Quadratic function fitting is used to calculate the location and scale of each key point in the continuous domain [2].

An Extraction Of Ancient Tamil Scripts From Epigraphy Images Using Gabor Filter, Active Contour And Lomo

S.Sundareswari, Dr.I.Kaspar Raj

Abstract : The extraction of Ancient script from epigraphy images is a tough task. Many techniques are available to extract the characters from the document image. Most of the authors have given their ideas to extract the characters from the document image. The extraction of ancient Tamil script from epigraphy images is a challenging problem in pattern recognition area still. In this paper, we present an efficient approach to extract the ancient Tamil script from epigraphy images. Recognizing the ancient Tamil script from epigraphy images is a very tough task. The proposed methodology consists of three stages: preprocessing, segmentation and feature extraction and classification. Experimental results show that the proposed methodology is more effective and efficient and also it extracts the ancient Tamil script from epigraphy images accurately.

Keywords: Tamil Ancient Characters , Gabor Filter and Active Contour, Feature Extraction(Zernike and Lomo), Classification(KNN and RF)

1 INTRODUCTION

Epigraphy is the study of the science of epigraphs or inscriptions, especially of ancient inscriptions. An epigraph is any kind of text from a single character to a lengthy document. Texts in epigraphy images are different from written texts of each culture [1]. In this work, we have used Ancient Tamil Script epigraphy images as input images. Segmentation of text or character from document image is an easy process. In epigraphy images, the engraved characters are different in size, color, and writing style. Mostly the epigraphy images have broken characters, uneven character intensity, touching characters each other and direction variations within symbols [2]. In preprocessing Gabor filter and active contour are used to enhance the image clearly without any noise. Then the combinational approach of two different feature extractions like Zernike moment and Lomo are used. Finally, in classification, two classifiers such as KNN(K-Nearest Neighbor) and RF(Random Forest) are used and tested. This paper is made as follows: Section 1 contains the initiation of epigraphy images, Section 2 explains the details of proposed methodology through various steps involved, Section 3 presents the details of combinational approach of feature extraction techniques, Section 4 discuss the various types of classifiers, Section 5 discussion on results, Section 6 concludes research performance.

PROPOSED METHODOLOGY

This proposed methodology involves the following steps: i) Image Acquisition: Read an input image and convert into suitable format. ii) Preprocessing: Gabor filter and Active contour used to remove the noise. iii) Feature Extraction:

The combinational approach of Zernike moment and Lomo are used to extract the ancient Tamil scripts from epigraphy images. iv) Classification: In this proposed methodology two classifiers like K-Nearest Neighbor, Random Forest are used and tested. In this proposed methodology, Gabor filter and Active contour are used for preprocessing and for feature extraction techniques combinational approach of Zernike Moment and Lomo features are used. Hence the combinational of these two feature extraction techniques may improve the overall detection rate of the characters in epigraphy images. Next, we have used two different classifiers like RF (Random Forest) and KNN (K Nearest Neighbor) is used and tested to select the best classification algorithm for this research work

2.1 Proposed Algorithm :

Step 1: Read the following inputs like Image, θ (theta), f (frequency), threshold value (g_m thr) Step 2: Read the testing image as Img . Step 3: Initial parameter for Gabor filter GF as $\theta \leftarrow 90$ and frequency value is 0.2 and threshold value is 50. Step 4: Apply Gabor filter using the equation 4.1. 4.1) $[Img_{gFC}, Img_{gFI}] \leftarrow GF(Img)$ where $\theta = 90, f = 0.2$. Step 5: for each pixel in image $\leftarrow Img$ $Img_{gF mag} = \text{SQRT}(Img_{gFC}^2 + Img_{gFI}^2)$ End Step 6: Find Gabor filter magnitude binary mask as $GF_{BW} \leftarrow Img_{gF mag} > g_m$ thr. Step 7: Extract the Contour of the image Img using GF_{BW} as initial mask in Active Contour mode. $Img_{ACBW} \leftarrow ACmode(Img, GF_{BW})$. Step 8: for each connected label $\leftarrow Img_{ACBW}$ Extract the characters by using the following feature extraction techniques. $Z_{feat} \leftarrow Zernike(BW_c)$ $L_{feat} \leftarrow Lomo(Img(BW_c))$ end Step 9: Feature vector = $[Zernike(BW_c), Lomo(Img(BW_c))]$ Step 10: Predict Characters \leftarrow Classifiers(Training feature, feature extraction). In this proposed methodology, first character samples are trained and tested. In preprocessing Gabor filter and Active contour are used. In Gabor filter, first read the input image, θ (ω), frequency (f) and threshold value (g_m thr). Next, read the testing image as Img . Then set the initial parameters for Gabor filter is θ is 90 and the frequency value is 0.2 and the threshold value is 50. Applying Gabor filter using the equation $[Img_{gFC}, Img_{gFI}] \leftarrow GF(Img)$ and get real and imaginary part of each pixel value using the above parameters. Then find out the Gabor filter magnitude binary mask of images as $GF_{BW} \leftarrow Img_{gF mag} > g_m$ thr This value should be greater than the threshold value. Next, extract the contour of the image Img using GF_{BW} as initial mask in Active Contour mode as $Img_{ACBW} \leftarrow ACmode(Img, GF_{BW})$. Finally, for each connected

- S.Sundareswari, Research Scholar, Research and Development Centre, Bharathiar University, Coimbatore, TamilNadu, India. profresearch2014@gmail.com
- Dr.I.Kaspar Raj, Director, Computer Centre, The Gandhigram Rural Institute Deemed University, Gandhigram, Dindigul, Tamil Nadu, India
- kasparraj@gmail.com
- (This information is optional: change it according to your need.)

Multi-Level Credit Card Fraud Detection

V. Sobanadevi, G. Ravi

Abstract: *Fraud detection in credit card transactions is one of the major requirements of the current business scenario due to the huge amount of losses associated with the domain. This work presents a multi-level model that can provide highly effective fraud detection in credit card transactions. The model is based on the amount for which the transaction is committed. The proposed MLFD model identifies the nature of the transaction and depending on the significance level of the transaction, the appropriate learning model is selected. Experiments were performed with the standard benchmark data and comparisons were performed with existing model in literature. Results shows that the proposed model exhibits high performance compared to the existing model.*

Keywords: *Credit card fraud detection, Decision Tree, Multi-Level Modelling, Naïve Bayes, Random Forest.*

I. INTRODUCTION

The process of fraud detection in credit card transactions is used to determine if a transaction is fraudulent or legitimate. The detection process is governed by the historical transactions. The past labelled transaction data is used to train the model for predicting the current transactions. This is a challenging task mainly due to the change in the spending behaviors of the users. Spending behaviors of users often tend to vary with time, which is considered as seasonal changes and also gradually varies over long time periods, which can be treated as trends in the behaviors. This results in the historical models becoming obsolete due to the absence of recent data. Further, fraudsters are also creating evolving models to compete with the recent advancements in the detection strategies.

Recent research shows that credit card fraud losses have resulted in significant losses for organizations and also for banks. It was observed that the losses due to credit card increased by 300% in five years (2010 to 2015) [1]. By the year 2020, the global loss due to fraud is expected to touch \$31.67 billion. The two major sources of card frauds were identified as counterfeit card based frauds and frauds at point-of-sale regions. Both these span globally and are not confined to bank premises. Hence they inadvertently require effective prediction mechanisms to stop the fraud from happening at those locations itself. This calls for a model that operates in real-time and also accurately. However, when it was viewed from a business perspective, not all predictions are to be considered significant. Every prediction involves a certain cost. Identifying a transaction as fraudulent requires business models to raise fraud alarms to customers, while falsely flagging a fraudulent transaction as legitimate incurs losses. It would be more appropriate to concentrate on higher

Valued transactions, which can help balance the cost factors. This work presents a multi-level fraud detection model that aims to consider the amount involved in the transaction to determine the type of prediction model to be used for prediction. The amount directly corresponds to significance of the transaction. Transactions with low significance are predicted using fast predictors, while transactions with higher significance are predicted with better prediction models. This enables the models to provide cost sensitive and faster prediction.

This remainder of this work is presented as follows: Section II presents a review of literature, Section III presents a detailed description of the working of the proposed model, Section IV presents the results and discussion and Section V concludes the work.

II. LITERATURE REVIEW

Credit card fraud detection is considered to be one of the major domains of analysis in the research world. This is due to the huge amount of losses associated with the domain. This section presents some of the significant works in the fraud detection domain.

Risk based modelling involving cost has become a major factor in the current fraud detection models. A risk based model was proposed by Bahnsen et al. [2]. This method is based on Naïve Bayes to identify the risk levels to be used for the final prediction. The work by Pozollo et al. [3] aims to provide testing mechanisms for models handling concept drift. A cost based model was proposed by Mahmoudi et al. [4]. This technique considers cost as a major factor for the prediction process involving credit card transactions. The method is based on Fisher Discriminant Analysis. A Game Theory based analysis for credit card fraud detection was proposed by Gianini et al. [5]. A Big Data based credit card fraud detection model that utilizes Big Data based techniques was proposed by Vaughan [6]. A customer incentive based model was proposed by Wang et al. [7].

Feature engineering is one of the mostly incorporated methods into the credit card fraud detection strategy. The behavioral change in users has made fraud detection in credit card transactions a complex process. In order to leverage the best rules, it becomes mandatory to mine as much information as possible from the transaction data. Feature engineering strategies perform the exam same process. A feature engineering based model for detecting frauds in credit card transactions was proposed by Lucas et al. [8]. This work aims at using HMM for the prediction process. Multiple perspectives are obtained by identifying several features that constitute behavioral nature of the customer. The method

Revised Manuscript Received on November 19, 2019

V. Sobanadevi, Research Scholar, Department of Computer Science, Jamal Mohamed College, Trichy, Tamilnadu, India.

G. Ravi, Associate Professor and Head, Department of Computer Science, Jamal Mohamed College, Trichy, Tamilnadu, India..

A GEOGRAPHIC STUDY ON REPRODUCTIVE HEALTH CARE OF WOMEN IN THANJAVUR DISTRICT

S.Geetha and R.Maniyosai

1. Research Scholar, Government Arts College (A), Kumbaonam.
2. Asst.Professor of Geography, Government Arts College (A), Kumbaonam.

ABSTRACT

"Health is wealth", Health plays vital role in wealth of a society. Reproductive health determines the health of the future society. According to 2011 census 19.1% of population was young in India. It is estimated to have increased 34.33% of youth in total population by 2020. Reproductive health includes the prenatal and postnatal care of pregnant women. According to WHO (World Health Organization), "Reproductive health is a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity, in all matters relating to the reproductive system and to its functions and processes". Reproductive health is highly affected by socio, economic and psychological factors. Antenatal health care is health support for during pregnancy time. ANC includes the Pregnancy registered within 12 weeks and regular checkups, regular vaccination, hemoglobin test, HIV test, Blood sugar, BP and intake of Iron and Folic Acid (IFA) tablets.

Keywords: *Reproductive health, Antenatal care, Socio economic status.*

1. INTRODUCTION

Maternal/Reproductive Healthcare includes the Prenatal, Postnatal care, family Planning, reduces maternal mortality and child Health care. A maternal health care service provides the safe motherhood. RCH Phase I provide the immunization, vitamin A, antenatal care visits, and Iron prophylaxis for pregnant women, and prevention of pneumonia, safe deliveries and postnatal care. The RCH Programme has given special attention to rural areas, Slums and Tribal areas. RCH phase II (2005), gives importance to essential obstetric care and strengthening referral system.

An Efficient Feature Extraction Approach for Extracting the Ancient Tamil Scripts from Stone Inscription Images

S.Sundareswari¹ Dr.I.Kaspar Raj²

¹ Research Scholar, Research and Development Centre,
Bharathiar University, Coimbatore, Tamil Nadu, India
¹profresearch2014@gmail.com

² Director, Computer Centre, The Gandhigram Rural Institute Deemed University,
Gandhigram, Dindigul, Tamil Nadu, India
²kasparraj@gmail.com

Abstract - Ancient Tamil script recognition is more complex due to enormously large character set and high similarity between characters. The extraction of ancient Tamil script from stone inscription images is a challenging problem in pattern recognition area. In this paper, we present an efficient approach to extract the ancient Tamil script from stone inscription images. Recognizing the ancient Tamil script from stone inscriptions is a very tough task. The proposed approach consists of three stages: preprocessing, segmentation and feature extraction and classification. Therefore this proposed approach helps to extract the characters from stone inscriptions easily and accurately. Experimental results show that the proposed methodology is more effective.

Keywords- Tamil Ancient Characters; Feature Extraction; Classification.

I. INTRODUCTION

Ancient Tamil script have large variations of individual writing style. Ancient script extraction from stone inscription images becomes very complex. It involves the following steps: 1) Collection of sample images. 2) Pre-processing on collected samples in order to get a clear image. 3) Feature extraction technique is required to extract features to form a feature vector. In this work, the combinational approach of three different feature extractions like HOG (Histogram of Gradient), Zernike moment, and Projection label are used. 4) After the feature extraction, a classification algorithm is used to classify the character samples. In this paper, three different classification techniques like SVM (Support Vector Machine), RF (Random Forest), KNN (K-Nearest Neighbor) are used and tested in order to select the best classification algorithm. Some of the stone inscription images with ancient Tamil script are given below in Figure 1, 2, 3.



Fig.1.



Fig.2.



Fig.3.

The *in vitro* hepatoprotective activity of the flower extract of *Senna alata* against Carbon tetra chloride induced liver damage in rats.

Dr. S. Jasmine Mary

Lecturer in Chemistry

Gcw(A), Kumbakonam

Abstract : The present investigation has been carried out to evaluate the *in vitro* hepatoprotective activity of different concentrations of *Senna alata* flower extract (100, 250 and 500µg/ml). The *in vitro* hepatoprotective activity was evaluated using estimation of Malondialdehyde, activity of GOT (AST), activity of GPT (ALT) and determination of reduced glutathione (GSH). Ethanolic extract showed the highest hepatoprotective activity when compared to standard due to the presence of phytochemicals present in the flower extract.

InterTerms: Hepatoprotective activity, *Senna alata*, Malondialdehyde, GOT, ALT.

1. INTRODUCTION

The liver plays an astonishing array of vital functions in the maintenance, performance and regulating homeostasis of the body. It is involved with almost all the biochemical pathways to growth, fight against disease, nutrient supply, energy provision and reproduction. And it functions as a centre of metabolism of nutrients such as carbohydrates, proteins and lipids and excretion of waste metabolites. The bile secreted by the liver has, among other things, plays an important role in digestion. Therefore, maintenance of a healthy liver is essential for the overall well being of an individual (Smuckler, 1975). Liver cell injury caused by various toxicants such as certain chemotherapeutic agents, Carbon tetrachloride, thioacetamide, chronic alcohol consumption and microbes are common. Enhanced lipid per oxidation during metabolism of ethanol may result in development of hepatitis leading to cirrhosis (Agarwal, 2001).

Herbal drugs have gained importance and popularity in recent years because of their safety, efficacy and cost effectiveness. The Indian Traditional Medicine like Ayurveda, Siddha and Unani are predominantly based on the use of plant materials. The association of medical plants with other plants in their habitat also influences their medicinal values in some cases. One of the important and well documented uses of plant products is their use as hepatoprotective agents. Hence, there is an ever increasing need for safe hepatoprotective agent (Ward and Daly, 199). In spite of tremendous strides in modern medicine, there are hardly any drugs that stimulate liver function, offer protection to the liver from damage or help regeneration of hepatic cell. Many formulations containing herbal extracts are sold in the Indian market for liver disorders (Achuthan *et al.*, 2003).


1.1 Liver Diseases and Medicinal Plants:

Liver has a pivotal role in regulation of physiological processes. It is involved in several vital functions such as metabolism, secretion and storage. Furthermore, detoxification of a variety of drugs



Contents lists available at ScienceDirect

Wave Motion

journal homepage: www.elsevier.com/locate/wamot

Inelastic dromions, rogue waves and lumps of (2+1) dimensional long dispersive wave equation



R. Radha^a, C. Senthil Kumar^{b,*}, R. Saranya^{a,c}

^a Centre for Nonlinear Science (CeNSc), Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, India

^b Department of Physics, Vinayaka Mission's Kirupananda Variyar Engineering College, Vinayaka Mission's Research Foundation (Deemed to be University), Salem 636308, India

^c Post-Graduate and Research Department of Mathematics, Government College for Women (Autonomous), Kumbakonam 612001, India

ARTICLE INFO

Article history:

Received 16 December 2017

Received in revised form 11 October 2018

Accepted 26 November 2018

Available online 1 December 2018

ABSTRACT

In this paper, we revisit the (2+1) dimensional long dispersive wave equation employing the truncated Painlevé approach. We then generate the solutions in the closed form in terms of lower dimensional arbitrary functions of space and time. By suitably harnessing the arbitrary functions present in the solution, we then construct localized solutions such as dromions, lumps and rogue waves. We have also explicitly brought out the generality of the localized solutions compared to the localized solutions generated earlier. The collisional dynamics of dromions, lumps and rogue waves is then explored.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

In recent times, there is a profound interest among physicists to study the nature of rogue waves, due to their wide range of applications, arising in multiple physical situations [1–4]. Generally, rogue waves which are also known as freak waves, occur in deep ocean. These waves are like shock waves and appear only for a short duration of time. The important feature of rogue waves is that they come from nowhere and disappear with no trace. Rogue waves find applications in various fields like, hydrodynamics [5], nonlinear optics [6–8], Bose–Einstein condensates [9,10], plasma physics [11], etc. Another interesting class of solutions which occur in (2+1) dimensional integrable systems is Dromions [12–15]. They arise essentially by virtue of coupling the field variable to a mean field/potential, thereby preventing wave collapse. Like dromions, there exists another class of solutions called lumps [16,17]. Unlike dromions, lumps do not interact with each other while rogue waves are found to be unstable in nature. Even though dromions and their collisional dynamics have been investigated in (2+1) nonlinear partial differential equations (PDEs) [18–24], the existence of lumps and rogue waves besides their dynamics has not yet been brought out explicitly in several nonlinear partial differential equations.

In this paper, we consider the (2+1) dimensional long dispersive wave (2LDW) equation introduced by Chakravarthy, et al. [25], by symmetrically reducing the self-dual Yang–Mill's equation. This equation is a generalization of the one dimensional long dispersive wave equation [26–28]. This equation holds significance in plasma physics under specific circumstances. Radha and Lakshmanan have established that the 2LDW equation is Painlevé integrable and constructed multi-dromion solutions [29] employing Hirota method. Later, Z. J. Fang and W. F. Min have also derived (1,N) dromion solution through a direct method [30]. Estevez [31] has also investigated the 2LDW equation using singular manifold method

* Corresponding author.

E-mail address: senthilbdu@gmail.com (C.S. Kumar).



P-ISSN: 2394-1685
E-ISSN: 2394-1693
Impact Factor (ISRA): 5.38
IJPESH 2019; 6(2): 78-80
© 2019 IJPESH
www.kheljournal.com
Received: 21-01-2019
Accepted: 22-02-2019

Dr. R Vidhya Sree
Ph.D., Director of Physical
Education, Govt. College for
Women, Kumbakonam, Tamil
Nadu, India

International Journal of Physical Education, Sports and Health

Emotional balance through yoga

Dr. R Vidhya Sree

Abstract

In psychology, emotion is often defined as a complex state of feeling that results in physical and psychological changes that influence thought and behaviour. Emotionality is associated with a range of psychological phenomena, including temperament, personality, mood, and motivation. "The major theories of motivation can be grouped into three main categories: physiological, neurological, and cognitive. Physiological theories suggest that responses within the body are responsible for emotions. Neurological theories propose that activity within the brain leads to emotional responses. Finally, cognitive theories argue that thoughts and other mental activity play an essential role in forming emotions. Emotion as a conscious and subjective mental reaction toward a particular event and is usually accompanied by changes in the physiologic and behavioural aspects of a person. From this definition, we can deduce that an emotion has four components, namely: cognitive reactions, physiological reactions, behavioural reactions and affect. Cognitive reactions refer to a person's memory, thinking and perception of an event. Physiological reactions are primarily caused by changes in the hormonal levels in the body. On the other hand, behavioural reactions comprise the active expression of the emotion. Lastly, affect includes the positive or negative state of the emotion and is what makes an emotion a conscious and subjective experience. Asanas are one path to blissful contentment, working to bring us closer by focusing our minds and releasing any emotional or inner tension in our bodies. When we practice yoga, we're not only working on our physical body. We're also working on our energy and our mind. As we grow in our yoga practice, we can see the results of all this work in our life: we feel more comfortable in our bodies, we have more energy, our emotions become more stable and our minds more focused.

Keywords: emotion, reactions, adrenaline, amygdala, yoga, asanas (poses)

Introduction

Emotions are lower level responses occurring in the subcortical regions of the brain, the amygdala, and the ventromedial prefrontal cortices, creating biochemical reactions in your body altering our physical state. They originally helped our species survive by producing quick reactions to threat, reward, and everything in between in their environments. Emotional reactions are coded in our genes and while they do vary slightly individually and depending on circumstances, are generally universally similar across all humans and even other species. For example, you smile and your dog wags its tail.

The amygdala plays a role in emotional arousal and regulate the release of neurotransmitters essential for memory consolidation which is why emotional memories can be so much stronger and longer-lasting. Emotions precede feelings, are physical, and instinctual. Because they are physical, they can be objectively measured by blood flow, brain activity, facial micro-expressions, and body language.

The nervous system, central as well as peripheral, plays a vital role in the regulation of emotion. Thalamus: It is composed of a group of nerve cells and acts as a relay center of sensory nerves. Stimulation of thalamus produces fear, anxiety, and autonomic reactions. Hypothalamus: It is considered the primary center for regulation of emotion. It also regulates the homeostatic balance, controls autonomic activity and secretion of endocrine glands, and organises the somatic pattern of emotional behaviour. Limbic System: Along with thalamus and hypothalamus the limbic system plays a vital role in regulation of emotion. Amygdala is a part of limbic system, responsible for emotional control and involves formation of emotional memories. Cortex: Cortex is intimately involved in emotions. However, its hemispheres have a contrasting role to play. The left frontal cortex is associated with positive feelings whereas the right frontal cortex with negative feelings.

Correspondence

Dr. R Vidhya Sree
Ph.D., Director of Physical
Education, Govt. College for
Women, Kumbakonam, Tamil
Nadu, India

Clapping hands: Boon for your health

Dr. R Vidhya Sree

Director of Physical Education, Govt. College for Women Kumbakonam, Tamil Nadu, India

Abstract

Generally, we clap when we want to praise someone or something. It is a part of appreciation practice but there is a lot more to it than we think! Clapping has been proven to provide a lot of health benefits that most of us aren't aware of! Just like laughing therapy, clapping therapy is also becoming popular as an easy and effective healing practice. Our human body is very complex. Every organ is linked to each other. The palms have blood vessels and nerve endings. If you stimulate them, a lot of your health problems would be healed! This is surprisingly true. The benefit of clapping: it has been scientifically proven that clapping is the best exercise for various disorders. Clapping therapy is a simple exercise which can be done by anyone irrespective of their age. Daily 10 minutes of clapping helps to stay active the whole day. Our body has 340 known pressure points, 28 of which are in our hand. Stimulation of this points benefits in a great way. It is the best cure for digestive disorders, back pain, neck pain, joint pain and also improves immune system. Clapping is associated with all good things in life – promotions, good grades, celebrations, and more. It's associated with happiness in our life. There are as many as 30+ acupressure points in both palms. If we can activate pressure points by clapping them together with regular practice, we can yield stunning benefits.

Keywords: clping therapy, pressure points, types of claping, good health

Introduction

We often give a big round of applause to the people we want to appreciate and encourage. Clapping is thus, one of the highly practised ways of appreciating someone. But you all will be surprised to know that clapping is actually beneficial for your body. You can live a healthy and a cheerful life by practising the art of clapping. There are a lot of pressure points in our body and thus it leads to various health benefits. When we clap our hands these pressure points get a pressure and it affects our body in a good way. Let's discuss the benefits of clapping in detail.

"Clapping" a simple striking of hands but it's much more than we think. Normally people clap to appreciate others for their good works and achievements or when they are in mood of joy. People also clap while singing songs, bhajans, and prayers at holy places. It is scientifically proved that clapping is very effective exercise to cure many human diseases. We all think of clapping as a way to applaud somebody for a job well done in sports, award shows, theater or movies. However, there are a number of hidden benefits that clapping has been proven to have. In the recent years, there are different types of groups in parks who get together in the morning and/or evening. Some of these groups indulge in laughing, there is also another group who gather around to clap. Our body is a bundle of nerves and blood vessels spread all through the body and is one of the most complex networks we will ever know. Each organ is linked to the other and a bad effect on one organ can put the other in danger. The palms, basically, contain nerve and blood vessel endings and stimulating them through clapping helps improve the health of organs like kidney, digestive tract, and lower back. Clapping activates the receptors in the palms and cause activation of the large area of the brain which leads the improvement in health. There are 28 different acupressure points for

almost all organs on our palm which are activated by clapping and this action improves our health slowly but effectively. Daily 10-20 minutes of clapping in morning keeps us fit and active. According to several well established healing modalities the hands and palms have numerous reflex points that, when stimulated, engage the body's healing response and prompt a gradual improvement in any type of ailment. We have receptors in the hands that are connected to sensory fields in the brain, Clapping activates these hands receptors, which in turn activate a fair portion of the brain, which itself leads to the activation of various body systems and their associated healing response in ways that are experientially evident but that we still need to better understand. Clapping stimulates blood circulation, the lifeline of the human body, and this helps with literally everything.




Fig 1



PAPER

Facile precipitation synthesis, structural, morphological, photoluminescence and photocatalytic properties of Ni doped ZrO₂ nanoparticlesRECEIVED
23 June 2019REVISED
14 August 2019ACCEPTED FOR PUBLICATION
2 September 2019PUBLISHED
11 September 2019G Rajesh¹, S Akilandeswari^{1,2} , D Govindarajan¹  and K Thirumalai³¹ Department of Physics, Annamalai University, Annamalai Nagar, Chidambaram, Tamil Nadu, India² Department of Physics, Government College for Women (Autonomous), Kumbakonam, Tamil Nadu, India³ Department of Chemistry, Government Arts College, Thiruvannamalai, Tamil Nadu, IndiaE-mail: akilaphy2010@gmail.com**Keywords:** Ni doped ZrO₂, photoluminescence, methyl violet, methyl blue, photocatalysis**Abstract**

In this study, Pure ZrO₂ (600 °C) and Ni (0.02, 0.04, 0.06, 0.8 M) doped ZrO₂ (600 °C) nanoparticles were obtained by facile precipitation method. The obtained products have been characterized by XRD, FTIR, UV-DRS, PL, FESEM-EDX, TEM-SAED pattern and XPS. The tetragonal structure of ZrO₂ (600 °C) gradually increases with the increasing Ni doping concentration and then a small NiO diffraction peak was detected at increasing Ni content. The Ni (0.08 M) doped ZrO₂ (600 °C) nanoparticles were approximately in spherical morphology was confirmed by FESEM and TEM results. The energy gap value of the pure ZrO₂ (600 °C) and Ni doped ZrO₂ (600 °C) nanoparticles decreased from 5.12 to 3.2 eV with enhancing nickel concentration. The surface defects and oxygen vacancies were analyzed by PL spectroscopy. This peak intensity gradually decreased with increase in nickel concentration. The photocatalytic activities of the Ni (0.08 M) doped ZrO₂ (600 °C) catalysts were studied by the photodegradation of MV and MB aqueous solution under sunlight irradiation. The Ni (0.08 M) doped ZrO₂ (600 °C) catalyst exhibited good photocatalytic efficiency and the enhanced photocatalytic mechanism was discussed.

1. Introduction

Water is an important necessity of life and it is used for several households with industrial activities. Water is the most vital natural resources, unluckily exploited the most. The main reason is population growth, increased industrialization and urbanization. Nowadays, all over the world is the confronting water crises because of the unlimited and too much exploitation of groundwater by several industries: namely textile, paper, cosmetic, leather, plastics, food, printing and pharmaceutical and so on. These industries are generating large amounts of harmful colored dyes, surfactants, toxic metal components when released into water resources consequence in ecological problems [1].

The greatly important task to the scientist to remove hazardous effluents from contaminated water. Therefore, numerous physical, chemical and biological treatment have been used to remove the organic/inorganic pollutants from water by applying chemical coagulation, floatation, reverse osmosis, advanced oxidation, bioremediation, Filtration, membrane process, adsorption and chemical precipitation. Among the techniques, the photocatalytic activity one of the most important methods that can be applied efficiently to attain high elimination rate of colored dyes from water. Photocatalytic degradation methods, which are eco-friendly and low cost in comparison to the other techniques [1, 2]. The several metal oxide and sulfate materials play an important role in the photocatalytic degradation of colored dyes. For example, TiO₂ [3], CuO [4], ZnO [5], Co₃O₄ [6], Fe₃O₄ [7], WO₃ [8], CeO₂ [9], NiO [10], CdS [11], ZnS [12] and so on. Among the oxide and sulfate materials, zirconium oxide is a most important semiconductor material, since it posses unique properties like high chemical inertness, high surface area, good thermal stability, reusability and low cost. In addition, it has

SICO-CHEMICAL CHARACTERIZATION KAEMPFEROL-3-O-β-D-(6"-COUMAROYL)- GLUCOPYRANOSIDE ISOLATED FROM PLUMBAGO ZEYLANICA

K. Vimala, N. Hayaraja, *A. John Merina

Research Scholar, Assistant Professor, Principal (Retd),
Research Department of Chemistry, Government College for Women (Autonomous), Kumbakonam, Tamilnadu-
612 001, India

Research Department of Chemistry, Government Arts College (Autonomous), Kumbakonam, Tamilnadu-612 002,
India

A new flavonol glycoside was isolated from the flowers of *Plumbago zeylanica* (Plumbaginaceae). It has been isolated from ethyl acetate extract of methanolic extract of the flowers of plant. The structure of this compound was determined as kaempferol-3-O-β-D-(6"-coumaroyl)-glucopyranoside on the basis of UV, H-NMR and ¹³C-NMR spectral data.

Keywords: Plumbaginaceae, *Plumbago zeylanica*, Kaempferol-3-O-β-D-(6"-coumaroyl)-glucopyranoside

Introduction
Plumbago zeylanica is a multipurpose medicinal herb of family Plumbaginaceae. *P. zeylanica* is the most frequent plant used in traditional system of medicine. A native of South Asia, the species is distributed all through most of the tropics and subtropics, deciduous woodland, savannas and scrub lands from sea level up to 2000 m altitude. In India *Plumbago zeylanica* grows in wild part as wild species however it is additionally cultivated due to its broad therapeutic applications. It is used in indigenous medicine, and many times known as "Chithira malam". Five coumarins - isoselin, 5-methoxycoumarin, isobornyl coumarin, umbelliferone and umbelliferone have been isolated from the roots of *Plumbago zeylanica*. The root is used as laxative, expectorant, astringent, and in dysentery. Tincture of root bark is used as antiperiodic. The leaves are caustic and used in remedy of scabies. It is a therapy for skin sicknesses, infections and intestinal worm's viz. leprosy, scabies, ringworm, hookworm, dermatitis, acne, warts consequently in historic times. The ancient structures of medicine in extraordinary parts of the areas have been using all parts of *P. zeylanica* for a vary of treatments. Paste made from roots of the plant is useful to the pores and skin to deal with abscesses, pores and skin ailments inclusive of ulcers and scabies.* Hence, the focus of this letter about used to be to work out the chemical from *Plumbago zeylanica* blossoms.

MATERIALS AND METHODS

Isolation and fractionation

The fresh flowers (2 kg) of *Plumbago zeylanica* (Plumbaginaceae) gathered from Mayiladuthurai had been extracted with 40% EtOAc (4 X 500 mL) under reflux. The alcoholic extract was once concentrated in vacuo and the aqueous concentrate steadily fractionated with diethyl ether (4 X 250 mL) and ethyl acetate (5 X 500 mL). The benzene fractions did no longer yield any crystalline solids.

EtOAc fraction: Kaempferol-3-O-β-D-(6"-coumaroyl)-glucopyranoside

The EtOAc fraction was once focused in vacuo and left in an ice chest for few days. A yellow solid that separated was isolated and studied. It developed a green coloration with alc. Fe³⁺, pink colour with Mg-HCl. It regarded purple under UV, this behaviour on publicity to NH₃. It responded to Wilson's boric acid test and additionally answered Gibb's take a look at and Molisch's test.

It no longer reply Horhammer-Hansel test. It had λ_{max}^{MeOH} 265, 340 nm; ¹³C-NMR (100 MHz, DMSO-d₆) δ ppm: 56.4 (C-2), 57.4 (C-3), 77.4 (C-4), 161.5 (C-5), 98.7 (C-6), 164.1 (C-7), 93.6 (C-8), 156.4 (C-9), 104.0 (C-10), 120.8 (C-11), 130.9 (C-2'), 115.1 (C-3'), 117.4 (C-4'), 115.2 (C-5'), 130.9 (C-6'), 100.8 (C-1'), 74.1 (C-2'), 77.4 (C-3'), 69.8 (C-4'), 76.3 (C-5'), 60.7 (C-6'), 121.8 (C-7'), 115.1 (C-3'), 159.9 (C-4'), 116.2 (C-5'), 128.2 (C-6'), 164.1 (C-O), 115.1 (CH), 144.7 (CH) 29.2. III. RESULTS & DISCUSSION

The fresh flowers of *Plumbago zeylanica* have been observed to comprise kaempferol-3-O-β-D-(6"-coumaroyl)-glucopyranoside. The free aglycone from ether fraction has been characterized as kaempferol on the basis of m.p., R_f values, UV and ¹³C-NMR. The UV spectrum of the glycoside exhibited foremost absorption peaks at 340 nm (band I) and 265 nm (band II). Band I absorption of the glycoside is reminiscent of a flavonol skeleton. An assessment of band I absorption of the glycoside and its aglycone revealed that there may additionally be 3-glycosylation in the flavonol. The existing of free -OH at C-5 in the glycoside is evident from its fine response to Wilson's boric acid test. The equal absorption also seems from the fact that a bathochromic shift of 53 nm and may want to be observed in the glycoside and the aglycone respectively in the AlCl₃-HCl spectra. A bathochromic shift of 46 nm in the glycoside and 49 nm in the aglycone in NaOMe spectra is suggestive of the affords of a free-OH at C-4 in the aglycone. The presence of catechol type of substitution in B-ring is validated as there is no extra shift in AlCl₃ spectrum in contrast to AlCl₃.

CHEMICAL CHARACTERIZATION OF KAEMPFEROL- O- β -D-GLUCOPYRANOSIDE ISOLATED FROM DIPTYCHANDRA AURANTICA

¹K. VIMALA & ²A. JOHNSMIRVA

¹Research Scholar & ²Principal (Retd.)

^{1,2} Department of Chemistry

Government College for Women (Autonomous), Kumbakonam

Tamil Nadu, India

¹adhavimala31@gmail.com

Abstract

A new flavonol glycoside was isolated from the fresh yellow flowers of *Diptychandra aurantica* belongs to Fabaceae family. It has been isolated from ethyl acetate extract of the flowers of plant. structure of the compound was confirmed as kaempferol-3-O- β -D-glucopyranoside-7-O-rhamnopyranoside, on the basis of UV, ¹H-NMR and ¹³C-NMR spectral values.

Key words Fabaceae, *Diptychandra aurantica*, kaempferol-3-O- β -D-glucopyranoside-7-O-rhamnopyranoside

1. Introduction

Natural products are large source of synthetic and herbal medicinal drugs¹. *Diptychandra aurantica* is belonging to the fabaceae family. Economically necessary of family of flowering p trees and shrubs are used to medicinal purposes². Some natural products occurs in Fabaceae fam flowers contain flavonoids. In the present investigation, plants have been chosen and the glycoside has been isolated and its therapeutic uses have been researched.

2. Material and Methods

The following are the materials and methods have been used in the experiment

2.1 Extraction and Fractionation

Fresh flowers (3kg) of *Diptychandra aurantica* collected from Kumbakonam at District were extracted with 90% methanol (6 X 500 ml) under reflux.

SRIMATHI PASUPATHY* MANIVANNAN RAJAMANICKAM.

Department of Chemistry, Government Arts College (Autonomous), Kumbakonam.

Abstract

Ni -doped TiO₂ nanocomposite was synthesized by the sol-gel method. The structural properties of nano composites have been characterized by X-Ray diffraction (XRD), Fourier -Transform Infrared (FT-IR), Scanning Electron Microscope (SEM). The photocatalytic activity of the synthesized nano composites was measured by the degradation of Methylene blue (MB) under solar the effects of initial ion concentration, Catalyst dosage, pH on removal efficiency through batch studies were investigated. The results showed that the degradation efficiency is decreased with an increase in initial concentration of MB from 50-150 mg/l.


KeyWords : Homogeneous photocatalysis, Heterogeneous photocatalysis, Photo catalyst, Photo degradation, Ni doped TiO₂, Solar irradiation.

Introduction

Energy shortage and environmental issues are the most pivotal complications of our modernized world [1]. Even though science and technology are making our life very simple and stylish, Human beings could not healthily lead their life. According to the latest state of the environmental report, 17 rivers of West Bengal, 83% of open wells and most of the rivers in Kerala are highly contaminated by dyes, organic pollutants, and heavy metals. Those water sources unfit for even bathing. So the low availability of freshwater sources and the necessity of water for a high population pushed our researchers focusing on wastewater treatment. Several methods are available for treating polluted water [2, 3]. There are several methods available for treating the wastewater like Coagulation, Flocculation, Ion- Exchange method, Adsorption, Reverse Osmosis, Ozonation, Chlorination and Photocatalysis. High salinity of wastewater and recalcitrant nature of synthetic dyes are the primary reason for avoiding biological method. Chlorination and ozonation are extremely high costs [4-6]. Traditional methods such as adsorption, Ultrafiltration, Reverse osmosis, coagulation, ion-exchange methods are having issues in converting organic compounds from wastewater but discharging of secondary wastes [4]. Photocatalysis is a current trending scenario that has more advantages for the removal of pollutants even at low concentrations. Photooxidation is a complete oxidation of organic pollutants that occur within a few hours without the generation of any secondary toxins. Here the photocatalyst using for this process is highly effective and very cheap [7,8]. Homogeneous and Heterogeneous photocatalysis is the main classification of photocatalysis. Semiconductor

material TiO₂, ZnO, SnO₂, CeO₂ used as photocatalyst. Semiconductor photocatalysts are generally choosing for its favoring electronic structure. While irradiating with a light source the electrons are excited from empty valence band to conduction band [9-18]. To avoid the electron-hole formation(e⁻/h⁺) which is the reason for less photodegradation, doping of transition metals and non-metal ion are suggested. The transition metal such as Fe[19-21], ZnO [22,23], Cu [24,25], Mg [26], Cd[27] , Mn [28] are the most promising metals doped with TiO₂ Solar light and UV light are generally used as light source [29].Nickel is particularly suitable for industrial applications and its optimum level of cost, non - toxic nature, controlling morphology and improving bandgap [30]. The previous researches reveal that Ni doping in TiO₂ can improve the visible light responsive activity against organic and toxic pollutants, over and that Ni doping can introduce and impurity energy level in the band structure. Conduction band



Sign in to edit and save changes to this file. 

Pasupathy and Rajamanickam, IJPSR, 2019; Vol. 10(10): 4731-4737.

E-ISSN: 0975-8232; P-ISSN: 2320-5148

IJPSR (2019), Volume 10, Issue 10

(Research Article)



INTERNATIONAL JOURNAL
OF
PHARMACEUTICAL SCIENCES
AND
RESEARCH



Received on 16 May 2019; received in revised form, 15 September 2019; accepted, 24 September 2019; published 01 October 2019

SYNTHESIS OF PURE AND BIO MODIFIED CALCIUM OXIDE (CaO) NANOPARTICLES USING WASTE CHICKEN EGG SHELLS AND EVALUATION OF ITS ANTIBACTERIAL ACTIVITY

Srimathi Pasupathy* and Manivannan Rajamanickam

Department of Chemistry, Government Arts College (Autonomous), Kumbakonam - 612001, Tamil Nadu, India.

Keywords:

CaO,
XRD, Antimicrobial,
Bio-modified CaO, Nanoparticle

Correspondence to Author:
Srimathi Pasupathy

Research Scholar,
Department of Chemistry,
Government Arts College
(Autonomous), Kumbakonam -
612001, Tamil Nadu, India.

E-mail: samsathi87@gmail.com

ABSTRACT: Calcium oxide nanoparticles (CaO NPs) gain great value in the areas of energy storage and drug delivery systems. Due to good porosity, it finds its part in storage systems and its biocompatibility earns it a good value in drug delivery and gene transfection. Synthesis of nanoparticles by waste materials and plants of metal oxide is gaining considerable interest due to environmentally friendly reactants and room temperature synthesis. This is most using method of preparation of nanoparticles as it makes use of pollution-free chemicals and encourages the use of non-toxic solvents such as water and plant extracts. In this present study, the calcium oxide nanoparticles synthesized by the eco-friendly green synthesis using environmentally benign waste chicken egg shells. CaO nanoparticles show better antibacterial activity. CaO was bio modified by *Cissus quadrangularis*, *Acalypha indica*, *Solanum nigrum*, *Phyllanthus niruri* to enhance the antibacterial activity, then the antibacterial activity of pure CaO and bio modified CaO was investigated against *B. subtilis*, *S. aureus* and *E. coli* species. The synthesized nanoparticles have been characterized by XRD. Antimicrobial activity of synthesized nanoparticles were also analyzed in this study.

INTRODUCTION: Nanotechnology, a newly evolved discipline aims the creation, manipulation and application of structures in the nanometer size range. Over a past few decades, nanoparticles have been extensively applied in various fields of public interest. Nanotechnology has the potential to revolutionize the pharmaceutical industry with new tools for the molecular treatment of diseases, and rapid disease detection.

It advances materials with a nano-dimension provides several means for innovative design of nano-size drug delivery systems (Nano systems) to overcome biological barriers in order to direct the drug¹.

Nanoparticles have also been successfully used to enhance the immobilization and activity of catalysts, in medical and pharmaceutical nano-engineering for delivery of therapeutic agents, in chronic disease diagnostics, and in sensors. The increasing adaptability of clinical microbial strains to antimicrobial drug resistance demands highly effective compounds for the treatment of critical microbial infection. Nanoparticles have demonstrated antimicrobial activities; the development of novel applications in this field makes them an attractive



International Journal of Pharmaceutical Sciences and Research

4731

Pasupathy and Rajamanickam, IJPSR, 2019; Vol. 10(10): 4731-4737.

E-ISSN: 0975-8232; P-ISSN: 2320-5148

alternative to conventional dispersions. Metal oxide nanoparticles (NPs) are known to possess strong antimicrobial properties. Inorganic metal oxides are being increasingly used for antimicrobial applications and these nanoparticles are well known for inherent antimicrobial activity². In this study, a new antimicrobial nanoparticle (CaONPs) was synthesized and assessed for antimicrobial activities using a panel of bacterial and fungal pathogen.

CaO is of particular interest as it is regarded as a safe material to human beings and animals and it has excellent antimicrobial potential and adeptness to indolent microbial endotoxin^{1, 4}. Due to idiosyncratic structural and optical properties of CaO it acts as a potential drug delivery agent⁵.

organic antimicrobial agents. Basic metal oxides such as MgO and ZnO, have been shown to exhibit antibacterial activity, where the particle size of the oxides appears to have an impact¹².

Recent studies have shown that green biologically based methods using microorganisms and plants to synthesize nanoparticles are safe, inexpensive, and an environment-friendly alternative. Both microorganisms and plants have long demonstrated the ability to absorb and accumulate inorganic metallic ions from their surrounding environment. These attractive properties make many biological entities efficient biological factories capable of significantly reducing environmental pollution and reclaiming metals from industrial waste¹¹. Recently, the biological synthesis of nanoparticles

Soft Multi b-Connectedness

R. GOWRI¹ and **G. SAHITHYABHARATHI²**

¹ Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Tiruchirappalli.

E-mail gowrigck@rediffmail.com

² Research Scholar, Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India
Affiliated to Bharathidasan University, Tiruchirappalli.

E-mail gsahithyabharathi@gmail.com

Abstract

In this paper, we will research the notion of soft multi b-connectedness in soft multi topological spaces by means of soft multi b-open sets, soft multi b-closed sets, soft multi b-separated sets and study its basic properties.

AMS Subject Classification: 54A40, 03E70

Keywords: soft multi sets, soft multi topological spaces, soft multi b-connected, soft multi b-component, soft multi b-hyperconnected.

1 Introduction

In 2011, Cagman et al. [4] introduced soft topology. Topological structure of soft set were introduced by Sabir and Naz [14]. They defined the soft topological spaces which are defined over a initial universe with a fixed set of parameters. Tokat and Osmanoglu [16] introduced soft multi topology. In 2013, Anjan Mukherjee et al. [3] introduced topological structure formed by soft multi sets and soft multi compact space. The notion of soft b-open sets was initiated in [5]. In 2019, Gowri and Sahithyabharathi [8] introduced and studied the concept of Soft multi connectedness in soft multi topological spaces.

In this paper, we introduce the notion of connectedness based on soft multi b-open sets and give some basic definitions and theorems about it.

2 Preliminaries

Definition 2.1 [12] *Let U be an initial universe and E be a set of parameters. Let $P(U)$ denote the power set of U and A be a non-empty subset of E . A pair (F, A) is called a soft set over U , where F is a mapping given by $F : A \rightarrow P(U)$.*

In other words, a soft set over U is a parametrized family of subsets of the universe U . For $\epsilon \in A$. $F(\epsilon)$ may be considered as the set of ϵ - approximate elements of the soft set (F, A) .

Definition 2.2 [2] *Let $\{U_i : i \in I\}$ be a collection of universes such that $\bigcap_{i \in I} U_i = \emptyset$ and let $\{E_{U_i} : i \in I\}$ be a collection of sets of parameters. Let $U = \prod_{i \in I} P(U_i)$ where $P(U_i)$ denotes the power set of U_i , $E = \prod_{i \in I} E_{U_i}$ and $A \subseteq E$. A pair (F, A) is*

Some New Family on Root Cube Mean Labeling Of Graphs

R. GOWRI¹ and G.VEMBARASI²

Assistant professor, Department of Mathematics,
Government College for Women (Autonomous), Kumbakonam, India.

E.mail: gowrigck@rediffmail.com

Research Scholar, Department of Mathematics,
Government College for Women (Autonomous), Kumbakonam,India.

E.mail: vembagopal@gmail.com

Abstract

In this paper, we investigate Root Cube Mean Labeling of some new graphs. We prove that the graphs such as $L_n \odot \bar{K}_2$, Star graph $K_{1,n}$ ($n \geq 1$), Comb graph, (m,n) – Kite graph, $D(Q_4) \odot K_1$ graph are Root Cube Mean graph.

Keywords: Mean Labeling of graphs, Root Cube Mean Labeling of graphs.

AMS Subject Classification (2010):05C78

INTRODUCTION:

The graphs considered in this paper are finite and undirected. The cardinality of the vertex set of G is size of G which is denoted by p and q respectively. The concept of Root Square Mean Labeling of graphs was introduced by S.S.Sandya, et. al. [1]. R.Gowri and G.Vembarasi [5] introduced the concept of Root Cube Mean Labeling of graphs and proved Root Cube Mean Labeling of some more disconnected graph in [7]. In this paper we investigate new Familier of Root Cube Mean Labeling of graphs.

2 Preliminaries

Definition 2.1 [2]

A Triangular Ladder is a graph obtained from L_n by adding the edges $u_i v_{i+1}$, $1 \leq i \leq n-1$, where u_i and v_i , $1 \leq i \leq n-1$ are the vertices of L_n such that $u_1 u_2 \dots u_n$ and $v_1 v_2 \dots v_n$ are two paths of length n in the graph L_n .

Definition 2.2 [2]

A Double Triangular snake $D(T_n)$ consists of two triangular snake that have a common path.

Soft Multi ω Closed sets

R. GOWRI¹ and G. SAHITHYABHARATHI²

¹ Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India

² Research Scholar, Department of Mathematics,
Government College for Women(Autonomous), Kumbakonam, India

Abstract

In this paper we introduce soft multi ω closed sets in soft multi topological spaces and establishes the relation between other existing generalized closed sets in soft multi topological spaces. Also, properties of soft multi ω closed sets are analysed.

AMS Subject Classification: 54A40, 03E70

Keywords: soft sets, soft multi sets, soft multi topological spaces, soft multi semi open sets, soft multi semi closed sets, soft multi ω closed set, soft multi ω open set.

1 Introduction

In 2011, Alkhazaleh et al. [1] as a generalization of Molodtsov's soft set introduce the definition of a soft multiset, its basic operations such as complement, union and intersection etc. In 2011, Cagman et al. [3] introduced soft topology. Topological structure of soft set were introduced by Sabir and Naz [11]. They defined the soft topological spaces which are defined over a initial universe with a fixed set of parameters. Tokat and Osmanoglu [14] introduced soft multi topology. In 2013, Anjan Mukherjee et al. [2] introduced topological structure formed by soft multi sets and soft multi compact space. In 2013, Chen [4] introduced soft semi open and soft semi closed sets. Levine [7] introduced generalized closed and open sets in topological spaces. After then Yuksel et al [12] studied behaviour relative to soft subspaces of soft generalized closed sets and continued investigating the properties of soft generalized closed and open sets. In 2018, Gowri and Sahithyabharathi [5] [6] introduced and studied the concept of soft multi generalized closed sets and soft multi regular generalized closed sets in soft multi topological spaces.

In this paper, Soft multi ω closed sets and Soft multi ω open sets are introduced in soft multi topological spaces and some of its basic properties are discussed.

2 Preliminaries

Definition 2.1 [9] Let U be an initial universe and E be a set of parameters. Let $P(U)$ denote the power set of U and A be a nonempty subset of E . A pair (F, A) is called a soft set over U , where F is a mapping given by $F : A \rightarrow P(U)$.

In other words, a soft set over U is a parametrized family of subsets of the universe U . For $\epsilon \in A$. $F(\epsilon)$ may be considered as the set of ϵ - approximate elements of the soft set (F, A) .

Synthesis of Cu-Mn doped ZrO₂ Nanoparticles, and its application.

Akilandeswari.S¹, Kalaiarasi.I² and Gnanasheela.U³

¹Associate Professor, Government College for Women (Autonomous),
Kumbakonam, Tamil Nadu, India.

^{2,3}Assistant professor, Government College for Women (Autonomous),
Kumbakonam, Tamil Nadu, India.

Abstract

In this study, pristine ZrO₂ and Cu (0.01, 0.02, 0.03, and 0.08 M)-Mn (0.04 M) doped ZrO₂ photocatalysts were synthesized through chemical precipitation process and calcined at 600°C. The calcined products were analyzed by XRD, FTIR, UV-DRS, PL, FESEM-EDX, TEM-SAED pattern and XPS techniques. The XRD pattern of pristine ZrO₂ and Cu (0.01, 0.02, 0.03, and 0.08 M), Mn (0.04 M) doped ZrO₂ nanoparticles shows the formation of t-tetragonal phase and shifting of tetragonal phase to monoclinic phase. The Cu (0.06 M)-Mn (0.04 M) doped ZrO₂ nanoparticles were approximately in quasi spherical morphology and high agglomeration was confirmed by FESEM and TEM results. The surface defects and oxygen vacancies were analyzed by PL spectroscopy. The Cu (0.06 M)-Mn (0.04 M) doped ZrO₂ nanoparticles exhibited enlarged photocatalytic activity with 93 % and 95 % degradation of methyl violet and methyl blue dyes under sunlight irradiation.

Keywords: Photoluminescence, Methyl violet and Methyl blue, sunlight radiation,

Photocatalysts

1. Introduction

The various organic dyes are increasingly applied in many industries such as leather tanning, textile, plastic, paper, food, pharmaceutical, cosmetics, photo-electrochemical cells, hair coloring, rubber and agricultural research and so on [1-3]. These types of industries subsequent discharge their colored wastewater containing such as pigments and dyes. Colored wastewater (dyes) significantly attributed to an environmental pollution and health risk [4]. Almost these colored wastewaters are toxic and non-biodegradable. Various physical, biological and chemical techniques have been applied for its removals such as precipitation, activated carbon, ozonation, membrane separation, Fenton, photo Fenton catalytic reaction, bioremediation, adsorption, reverse osmosis, coagulation and ultra filtration [1-3, 5], etc. Among these techniques, photocatalysis is most important approach because it's low cost, eco-friendly and totally oxidizes the pollutants to H₂O and CO₂ [1-3].

There are many materials applied for photocatalytic degradation including TiO₂ [6] ZnO [7], WO₃ [8], ZrO₂ [9], SnO₂ [10], In₂O₃ [11], Fe₂O₃ [12], NiO [13], CuO [14] CdS [15], and PbS [16]. Among these, zirconium oxide (ZrO₂) is a most attractive material. ZrO₂ is a wide band gap (5.0–5.5 eV) semiconductor, and this oxide material had detected broad application in ceramics, catalysts, gas sensors, fuel cell, solid-state electrolytes, barrier coatings and optical devices [17-21] It is exhibit excellent thermal, mechanical, electrical and optical properties, such as low thermal conductivity, high hardness, high fracture toughness, high refractive index, optical transparency, high corrosion resistance and polymorphic nature [17-22], etc. Zirconium oxide (ZrO₂) exhibits three kinds of polymorphs as follows: Monoclinic (m-ZrO₂), tetragonal (t-ZrO₂) and cubic (c-ZrO₂). The monoclinic (m-ZrO₂) phase is stable below at <1170°C, t-tetragonal (t-ZrO₂) phase is stable at 1170–2370°C, and c-cubic (c-ZrO₂) phase is stable at >2370°C respectively [23]. These crystal phases strongly depend on thermal condition and preparation method [24].



A STUDY ON MUNICIPAL BORROWINGS IN TAMIL NADU

Dr. A. Thomas David¹ and **S. P. Balasangeetha²**

¹Associate Professor & Research Guide, PG &
Research Department of Economics, Government Arts
College (Autonomous) Kumbakonam.

²Ph.D Research Scholar, PG & Research Department of
Economics, Government Arts College (Autonomous)
Kumbakonam.



ABSTRACT:

The Local Authorities Loans Act, 1914 regulate the municipal borrowings. Most states generally allow municipalities to secure their borrowings against revenue/funds and not immovable property. In early days financial needs of the urban infrastructure meet by budgetary allocations and borrowings from state-owned institutions like HUDCO and LIC backed by state government guarantees. The commercial sources and volume of borrowings are Government Institutions, Capital Markets, Specialized Infrastructure Finance Entities, Sector Specific Municipal Development Funds/Facilities, and Tamil Nadu Urban Development Fund (TNUDF).

KEYWORDS: Municipalities, Borrowings, Tamilnadu.

INTRODUCTION:

In India urban areas are the drivers for sustaining economic growth and reducing incidence of poverty. It contributes about 65% of GDP in the year 2011. Urban economic activities depends on infrastructure such as power, telecom, roads, water supply and mass transportation, coupled with civic infrastructure such as sanitation and solid waste management. In large cities urban population increase day by day, which brings tremendous pressure on urban services such as water supply, sewerage, drainage, solid waste management and urban transportation networks and systems. Hence, it is mandatory for municipal governments to generate resources for meeting operations and maintenance expenses as well as administration. In this context, it is important to encourage municipal governments to leverage their resources and borrow from various sources. The Local Authorities Loans Act, 1914 regulate the municipal borrowings. Most states generally allow municipalities to secure their borrowings against revenue/funds and not immovable property.

OBJECTIVES OF THE STUDY

- To know the various Sources Municipal Borrowings.
- To understand the Municipal Borrowings in Tamil Nadu.

METHODOLOGY

This study is primarily based on secondary data collected from various published sources.



GOVERNMENT COLLEGE FOR WOMEN (AUTONOMOUS)

KUMBAKONAM – 612 001

Affiliated to Bharathidasan University

DST - CURIE Sponsored Institution

IV Cycle of Accreditation

☎ 0435 – 2401391

✉ principal@gwk.ac.in



CRITERION III – RESEARCH, INNOVATIONS AND EXTENSION

FRONT PAGE OF PUBLISHED PAPERS

2018-2019

BEHAVIOURAL ANOMALIES IN *CYPRINUS CARPIO* L. INDUCED BY MONOCROTOPHOS
INTOXICATION IN SUBLETHAL DOSES*Sharmila, G.¹ and A.V.Kavitha²¹P.G and Research Department of Zoology and Wildlife Biology
A.V.C. College (Autonomous), Mananmpandal – 609 305, Mayiladuthurai
Nagapattinam Distict, Tamil Nadu, India

*Corresponding author email id: viswanathansharmila@gmail.com

²P.G and Research Department of Zoology ,
Govt. College for Women (Autonomous), Kumbakonam - 612 001

ABSTRACT

Though plenty of research is available for short term impact of pesticides on fishes yet study on the quantification of behavioural implications were not carried out so far. In the present study, an attempt has been made to investigate the impact of chronic toxicity of Monocrotophos on the behavioural responses of freshwater fish, *Cyprinus carpio*. A static renewal bioassay was conducted to determine the acute toxicity of commercial grade organophosphorus pesticide, Monocrotophos (36% SL) to Common carp, *Cyprinus carpio*. The LC₅₀ concentrations of monocrotophos to carp fingerlings for 96 hrs was found to be 15.53 ppm. The fishes were exposed to the sublethal concentrations 1/5th and 1/10th of LC₅₀96 hrs for 30 days. The impact of toxicity on the behaviour and movement of fishes were assessed at five sampling points viz. at 0, 7, 14, 21 and 28 days of the exposure period. Even though the mortality was insignificant in the sublethal concentrations, considerable variations in behavioural responses such as lethargy, increased mucous secretion, surfacing, gulping of air, skin discolouration, respiratory distress and lack of appetite were observed. The one way Anova results showed that treated fishes showed significant differences in the movement when compared to their movement prior to intoxication at the same interval of time (F value = 3.328, $p < 0.05$).

KEYWORDS: Bioassay, Behaviour, *Cyprinus carpio*, Monocrotophos, Sublethal concentration, Movement

INTRODUCTION

Application of synthetic fertilizers, insecticides and pesticides are increasing now-a-days with increasing demand of agricultural food materials throughout the world. There is a high chance of aquatic ecosystem to be contaminated, located in industrial or agricultural areas, through runoff or ground water leaching of a variety of chemicals (Todd and Leuwen, 2002), thereby affecting many other organisms away from the primary target. It is estimated that only 0.1% of the pesticide reaches the specific target (Agular, 2002). Exposure of aquatic ecosystems to these pesticides is difficult to assess because of their short persistence in the water due to low solubility and rapid degradation. Yet monitoring of these pesticides is important because they are highly toxic and pose greater hazards to the aquatic organisms especially fishes (Talebi, 1998; Uner *et al.*, 2006; Banaee *et al.*, 2008). Pesticides, upon entering into the organs and tissues of fishes significantly damage certain physiological and biochemical processes.

Monocrotophos (3-(dimethoxyphosphinyloxy)-N-methylisocrotonamide) is a synthetic organophosphate, nonsystemic, broad spectrum insecticide and acaricide, acting as a cholinesterase inhibitor with contact, stomach and respiratory action. The major use of monocrotophos in farming is to protect crops against a variety of insect pests from soft bodied insect to leaf eating beetles.

Fishes, as they occupy the top most trophic level of aquatic food chain are able to take up and retain chemicals dissolved in water not only via active or passive processes but also by biomagnification. They represent good model systems to detect and document pollutants released into their environment. The deterioration of water quality can be detected easily by any change in the behaviour and physiology of fishes, hence they are excellent biological indicators. Moreover, as the mammalian and piscine systems exhibit similar toxicological and adaptive responses to oxidative stress, there is a growing concern in understanding the physiological mechanisms associated with fish responding to environmental stress. A long term exposure to sublethal concentrations of pesticides in aquatic environments cause more prominent structural and functional changes in aquatic organisms rather than mortality. Since monocrotophos has



Research Article

SUBLETHAL EFFECT OF ATRAZINE ON THE INTESTINE OF AN INDIAN EARTHWORM *LAMPITO MAURITII* (KINBERG) (ANNELIDA; OLIGOCHAETA)

*¹Kavitha, V. and ²Anandhan, R.

¹Department of Zoology, Government Women's College (Autonomous), Kumbakonam-612 001, Tamil Nadu, India

²Department of Zoology, Government Arts College (Autonomous), Kumbakonam-612 001, Tamil Nadu, India

Article History: Received 30th May 2018; Accepted 08th June 2018; Published 9th August 2018

ABSTRACT

The present study has investigated the toxic effect of Atrazine on the intestine of the earthworm *L. mauritii* in a laboratory experiment. A sub-lethal concentration of Atrazine (1/5th of 96 h LC₅₀ value-1.99 mg/kg) was applied for 30 days. The changes such as vacuolization, degenerated nuclei, damaged epithelial lining of villi and congestion of blood sinuses were observed in the intestine of 1st, 5th and 15th day of Atrazine exposure. In the 30th day of exposure, slight damages were observed. These results suggest that Atrazine could severely affect the intestine of 1st, 5th and 15th day of exposure when compared to 30th day. Histopathological study in *L. mauritii*'s intestine is a suitable parameter for detection of soil contamination by application of pesticides in agricultural field.

Keywords: Atrazine, Earthworm, Intestine, Soil, Toxicity.


INTRODUCTION

Increasing population growth and urbanization, especially in a developing country like India, necessitates producing more food. Food crops require fertile soil to grow. For terrestrial ecosystems, soil serves as a medium of entry to the nutrients. Continuous agricultural activities tend to decrease the soil fertility. However, the increasing application of herbicides and pesticides has also threatened the human environment and the ecosystems with deleterious consequences. Insecticide residues reach the soil in a variety of ways, causing toxicity to beneficial organisms. Earthworm represent the greater fraction of biomass of invertebrates in the ground (>80%). They can play a variety of important roles in agro ecosystems. Their feeding and burrowing activities incorporate organic residues and amendments into the soil, enhancing decomposition, humus formation, nutrient cycling, and soil structural development (Kladivko *et al.*, 1986). Therefore, earthworms can be used as bio indicators to detect pesticide contamination in agricultural soil. A pesticide is a chemical intended to kill, or disrupt the population of pest organisms. Pests are unwanted insects, mites, plants, disease causing organisms, and other organisms that interfere with health or

commerce. Pesticides are classified into insecticides, herbicides, fungicides, nematicides, rodenticides, etc. based on their action on type of pest. Insecticide target insects, herbicides target plants, fungicides target disease causing fungi, nematicides target nematodes and so on. The most common of these are herbicide which account for approximately 80% of all pesticide use. Atrazine is one of the most commonly used herbicides to prevent pre and post emergence broadleaf weeds in crops such as sugarcane and maize and on turf, such as golf courses and residential lawns. Atrazine is prepared from cyanotic chloride, which is treated sequentially with ethylamine and isopropylamine. The chemical name of atrazine is 2-chloro-4-ethylamino-6-isopropylamino-s-triazine. It is colourless solid and soluble in water.

Histology is the most useful tool for determining the influence of agricultural pesticides, industrial pollutants, organic wastes etc., at tissue level of an organism as it provides useful information concerned with the growth, damage and disorganization of tissues. Histopathological studies may signal a damaging effect of organisms resulting from prior or ongoing exposure to toxic agents. Earthworms have been shown to be affected by the

*Corresponding Author: Dr. V. Kavitha, Assistant Professor, Department of Zoology, Government College for Women (Autonomous), Kumbakonam-620 002, Tamil Nadu, India Email: kavitsamiksha@gmail.com



Toxicology

Nanosilver crystals capped with *Bauhinia acuminata* phytochemicals as new antimicrobials and mosquito larvicides

Naiyf S. Alharbi^{a,**}, Marimuthu Govindarajan^{b,c,*}, Shine Kadaikunnan^a, Jamal M. Khaled^a, Taghreed N. Almanaa^a, Sami A. Alyahya^d, Mohammed N. Al-anbr^a, Kasi Gopinath^e, Arumugam Sudha^f

^a Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^b Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India

^c Department of Zoology, Government College for Women, Kumbakonam, 612001, Tamil Nadu, India

^d National Center for Biotechnology, King Abdulaziz City for Science and Technology, Riyadh 11442, Saudi Arabia

^e Department of Packaging, Yonsei University, Gangwondo 26493, Republic of Korea

^f Department of Bioinformatics, Alagappa University, Karaikudi, India

ARTICLE INFO

Keywords:

Antibacterial activity
Antifungal activity
Insecticides
Mosquito control
Pesticides

ABSTRACT

To develop novel nanoformulated insecticides and antimicrobials, herein we produced Ag nanoparticles (AgNPs) using the *Bauhinia acuminata* leaf extract. This unexpensive aqueous extract acted as a capping and reducing agent for the formation of AgNPs. We characterized *B. acuminata*-synthesized AgNPs by UV–vis and FTIR spectroscopy, XRD and TEM analyses. UV–vis spectroscopy analysis of *B. acuminata*-synthesized AgNPs showed a peak at 441.5 nm. FTIR shed light on functional groups from the phytoconstituents involved in nanosynthesis. XRD of *B. acuminata*-synthesized AgNPs suggested a face-centered cubic structure, with a highly crystalline nature. TEM of *B. acuminata*-synthesized AgNPs revealed mean size of 25 nm, with round shape. AgNPs tested at 60 µg/mL inhibited the growth of 5 bacteria and 3 fungal pathogens. In the insecticidal assays on important mosquito species, LC₅₀ of the aqueous extract of *B. acuminata* leaves on the larvae of *Anopheles stephensi*, *Aedes aegypti*, and *Culex quinquefasciatus* were 204.07, 226.02, and 249.24 µg/mL, respectively. The *B. acuminata*-synthesized AgNPs exhibited higher larvicidal efficacy, with LC₅₀ values of 24.59, 27.19, and 30.19 µg/mL, respectively. Therefore, herein we developed a single-step, reliable, inexpensive, and environmentally non-toxic synthesis process to obtain AgNPs with high bioactivity against pathogens and vectors. Given the effective antimicrobial and larvicidal activity, nanoparticles fabricated using plant extracts and extremely low concentrations of trace elements, such as silver, can be exploited for multipurpose activities. Our results pointed out that *B. acuminata*-synthesized AgNPs have a promising potential in antimicrobial food packaging, as well as a foliar spray to control plant pathogens in the field, and to synergize the efficacy of fungicidal and larvicidal formulations.

1. Introduction

Over 5000 years ago, the Egyptians, Greeks, Persians, and Romans used to store food in silver-based vessels [1]. Similarly, ancient Indians used silver, gold, and copper-based jewelry such as bangles, belts, and anklets, as well as drinking vessels and kitchen utensils. These metals have good antibacterial activity, controlling microbial growth [1,2]. Furthermore, the ancient medicine system of Ayurveda uses powdered forms of copper, iron, silver, and gold in various formulations [2–4]. The shift in the use of metal vessels or adornments to metal powders


indicates a shift from the bulk level to nano-size. Thus, the traditional Indian system of medicine was rooted in knowledge about micro/nano-materials for drug carrier applications as early as around 2000 BCE [3,4]. For the preparation of nanomaterials, plant-based extracts were used for reducing and capping purposes, and the samples were then calcined to obtain the powder form. Such formulations were used as drug carriers and in the treatment of diseases like cancer and skin infection [4–6].

In the past decade, silver nanoparticles (AgNPs) have been subjected to extensive research owing to the typical physical and electrochemical

* Corresponding author at: Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, 608002, Tamil Nadu, India.

** Corresponding author.

E-mail addresses: nalharbi1@ksu.edu.sa (N.S. Alharbi), drgovind1979@gmail.com (M. Govindarajan).



Searching for crab-borne antimicrobial peptides: Crustin from *Portunus pelagicus* triggers biofilm inhibition and immune responses of *Artemia salina* against GFP tagged *Vibrio parahaemolyticus* Dahv2



Ravichandran Rekha^a, Baskaralingam Vaseeharan^{a,*}, Ramachandran Ishwarya^a, Mahalingam Anjugam^a, Naiyf S. Alharbi^b, Shine Kadaikunnan^b, Jamal M. Khaled^b, Mohammed N. Al-anbr^b, Marimuthu Govindarajan^{c,d}

^a Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Karaikudi, 630004, Tamil Nadu, India

^b Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, 11451, Saudi Arabia

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar, 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam, 612 001, Tamil Nadu, India

ARTICLE INFO

Keywords:

In vivo
Antibacterial
Antibiofilm
Crustin
Haemolymph
Portunus pelagicus

ABSTRACT

Marine organisms represent a huge source of novel compounds for the development of effective antimicrobial drugs. The present study focus on the purification of the antimicrobial peptide crustin from the haemolymph of the blue swimmer crab, *Portunus pelagicus*, by blue Sepharose CL-6B matrix assisted affinity column chromatography. Crustin showed a single band with a molecular mass of 17 kDa in SDS-PAGE analysis. The XRD analysis exhibited peaks at 32° and 45° while a distinct peak with a retention time of 1.8 min resulted in high performance liquid chromatography (HPLC) pointing out the crystalline nature and purity of crustin, respectively. Crustin purified from *P. pelagicus* (Pp-Cru) showed immunological activities, triggering encapsulation, phagocytosis on Sepharose beads and yeast (*Saccharomyces cerevisiae*) respectively. Furthermore, encapsulation of GFP tagged *V. parahaemolyticus* in *Artemia salina* and challenging study were assessed under CLSM and the potential of Pp-Cru was examined *in vivo*. In addition, the growth reduction and biofilm inhibition potential of Pp-Cru on *Staphylococcus aureus*, *Enterococcus faecalis* (Gram-positive bacteria) and *Pseudomonas aeruginosa*, *Escherichia coli* (Gram-negative bacteria) was evidenced by inverted and confocal laser scanning microscopic analysis, revealing that 100 µg/ml of Pp-Cru can disrupt the biofilm matrix thereby the thickness of biofilm was significantly reduced. Overall, the present investigation might provide a sensitive platform to realize the significant function of Pp-Cru in crustacean immune mechanism as well as its potential to bacterial growth inhibitor. The functional properties of purified Pp-Cru antimicrobial peptide may lead to a superior understanding of innate immune response in *P. pelagicus* species, which suggest the promising application for drug development in aquaculture.

1. Introduction

Natural products, including those isolated from marine organisms, are outstanding sources of molecules multipurpose applications (Gobi et al., 2016; Benelli, 2016, 2018a,b,c; Anjugam et al., 2017; Ishwarya et al., 2018). An antimicrobial peptide (AMP) is a molecule acting as an integral component of the humoral part participated in the innate immune defense mechanism of vertebrates and invertebrates. AMPs represent an evolutionary and protection strategy of an organism against invading pathogens, including bacteria and viruses (Malanovic and

Lohner, 2016; Abinaya et al., 2018). AMPs act as an effective defensive weapon and are present in higher concentration of haemocytes, playing an essential role in the flourishing evolution of intricate multicellular organisms. Since crustaceans lack in adaptive immune system responses, the AMPs have a major role as immune effectors against microbial infections (Antony et al., 2010). Crustacean-borne AMP has antibacterial activity which independently led to the lysis of bacterial cell wall. It can be able to function at very low concentrations as bactericidal agent against both Gram positive as well as Gram negative bacteria with haemocyte lysate supernatants (HLS) (Schnapp et al.,

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

<https://doi.org/10.1016/j.molimm.2018.07.024>

Received 26 February 2018; Received in revised form 9 July 2018; Accepted 14 July 2018

Available online 30 July 2018

0161-5890/ © 2018 Elsevier Ltd. All rights reserved.



Biocompatible properties of nano-drug carriers using TiO₂-Au embedded on multiwall carbon nanotubes for targeted drug delivery

Viswanathan Karthika^a, Periyannan Kaleeswarran^a, Kasi Gopinath^b, Ayyakannu Arumugam^{b,*}, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^e, Jamal M. Khaled^e, Mohammed N. Al-anbr^e, Giovanni Benelli^{f,g}

^a Department of Nanoscience and Technology, Alagappa University, Karaikudi, India

^b Department of Botany, Alagappa University, Karaikudi, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalaiagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^f Department of Agriculture, Food and Environment, University of Pisa, via del Borghetto 80, 56124 Pisa, Italy

^g The BioRobotics Institute, Sant'Anna School of Advanced Studies, viale Rinaldo Piaggio 34, 56025 Pontedera, Pisa, Italy

ARTICLE INFO

Keywords:

Anticancer drugs
Antimicrobial activity
Doxorubicin
Green synthesis
Nanoparticle characterization

ABSTRACT

Nanomaterial-based drug carriers have become a hot spot of research at the interface of nanotechnology and biomedicine because they allow efficient loading, targeted delivery, controlled release of drugs, and therefore are promising for biomedical applications. The current study made an attempt to decorate the multiwalled carbon nanotubes (MWCNT) with titanium dioxide-gold nanoparticles in order to enhance the biocompatibility for doxorubicin (DOX) delivery. The successful synthesis of nano drug carrier (NDC) was confirmed by XRD, XPS and UV-Visible spectroscopy. FESEM and TEM revealed that the morphology of NDC can be controlled by manipulating the reaction duration, MWCNT concentration and TiO₂-Au source concentration. Results showed that TiO₂ and Au nanoparticles were well coated on MWCNT. NDC had finely tuned biocompatible properties, as elucidated by hemolytic and antimicrobial assays. NDC also showed a high antioxidant potential, 80.7% expressed as ascorbic acid equivalents. Commercial DOX drug was utilized to treat A549 and MCF7 cancer cell lines showing improved efficiency by formulating it with NDC, which selectively delivered at the pH 5.5 with drug loading capacity of 0.45 mg/mL. The drug releasing capacity achieved by NDC was 90.66% for 10 h, a performance that far encompasses a wide number of current literature reports.

1. Introduction

Nanomaterials were widely used as drug carriers in the field of pharmacology due to their unique characteristics, which includes effectual loading capacity, targeted drug delivery and controlled drug release [1,2]. The current expansion in the area of controlled drug delivery has led to significant applications in biomedicine and nanotechnology. Various classes of nanoparticles have been successfully used for in the fight against parasites and vectors, as well as for other biological applications [3–6]. Doxorubicin hydrochloride (DOX) is an anticancer drug, also known as Adriamycin, which has shown high efficacy against several cancer cells. The nonspecific interaction between DOX and the physiological pathways can cause side effects, including allergic reactions such as anaphylaxis, heart damage (heart valves blockage), tissue damage, radiation recall, and treatment-related

leukemia. Also, it led to common clinical side effects includes vomiting, hair loss, bone marrow suppression, rash, and mouth inflammation [7]. In addition to this, DOX shows dose dependent cardiotoxicity, which makes an urge to develop a targeted DOX delivery system [8–11]. It is required the targeted releasing towards the cancer cells or tissue by different carrier biomolecules, dendrimers, liposome, and micelles [12–14]. For this purpose, synthetic polymers (polyethylene glycol), bio polymers (chitosan) [15–18], nanomaterials in form of metal [19], metal oxides [20], and carbon based materials (e.g., carbon nano tubes, CNT) [21], graphene [22], quantum nanodots [23] can be investigated. Among them, nanomaterials showed unique properties, are often cheap, highly stable and biocompatible. Hence, several nanoparticles have been considered for drug delivery, and carbon nanotubes are highly suitable for targeted drug delivery [24–26], as well as for biosensors [27], tissue engineering [28], bone regeneration [29], and

* Corresponding author.

E-mail address: ayyakannuarumugam@gmail.com (A. Arumugam).



Structural characterization of *Bacillus licheniformis* Dahb1 exopolysaccharide—antimicrobial potential and larvicidal activity on malaria and Zika virus mosquito vectors

Muthukumar Abinaya¹ · Baskaralingam Vaseeharan¹ · Mani Divya¹ · Sekar Vijayakumar¹ · Marimuthu Govindarajan^{2,3} · Naiyf S. Alharbi⁴ · Jamal M. Khaled⁴ · Mohammed N. Al-anbr⁴ · Giovanni Benelli^{5,6}

Received: 14 February 2018 / Accepted: 10 April 2018 / Published online: 27 April 2018
© Springer-Verlag GmbH Germany, part of Springer Nature 2018

Abstract

Microbial polysaccharides produced by marine species play a key role in food and cosmetic industry, as they are nontoxic and biodegradable polymers. This investigation reports the isolation of exopolysaccharide from *Bacillus licheniformis* Dahb1 and its biomedical applications. *Bacillus licheniformis* Dahb1 exopolysaccharide (BI-EPS) was extracted using the ethanol precipitation method and structurally characterized. FTIR and ¹H-NMR pointed out the presence of various functional groups and primary aromatic compounds, respectively. BI-EPS exhibited strong antioxidant potential confirmed via DPPH radical, reducing power and superoxide anion scavenging assays. Microscopic analysis revealed that the antibiofilm activity of BI-EPS (75 µg/ml) was higher against Gram-negative (*Pseudomonas aeruginosa* and *Proteus vulgaris*) bacteria over Gram-positive species (*Bacillus subtilis* and *Bacillus pumilus*). BI-EPS led to biofilm inhibition against *Candida albicans* when tested at 75 µg/ml. The hemolytic assay showed low cytotoxicity of BI-EPS at 5 mg/ml. Besides, BI-EPS achieved LC₅₀ values < 80 µg/ml against larvae of mosquito vectors *Anopheles stephensi* and *Aedes aegypti*. Overall, our findings pointed out the multipurpose bioactivity of BI-EPS, which deserves further consideration for pharmaceutical, environmental and entomological applications.

Keywords Exopolysaccharide · ¹H-NMR · Dengue fever · Insecticide · Drug development · Microbial pathogens · Pesticide · Yellow fever

Introduction

Microbial exopolysaccharides (EPS) are macromolecules of high molecular weight polymers with glycosidic linkages. Recently, natural polymers have been increasingly

requested for numerous industrial applications. EPS can be secreted by several microorganisms, including bacteria, fungi, and algae during their growth (Wang et al. 2010a, b; Ismail and Nampoothiri 2010; Poli et al. 2010; Zhang et al. 2016). As a general trend, it has been highlighted

Responsible editor: Philippe Garrigues

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s11356-018-2002-6>) contains supplementary material, which is available to authorized users.

✉ Baskaralingam Vaseeharan
vaseeharanb@gmail.com

¹ Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block, 6th floor, Burma Colony, Karaikudi, Tamil Nadu 630004, India


² Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalai Nagar, Tamil Nadu 608 002, India

³ Department of Zoology, Government College for Women, Kumbakonam, Tamil Nadu 612 001, India

⁴ Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

⁵ Department of Agriculture, Food and Environment, University of Pisa, via del Borghetto 80, 56124 Pisa, Italy

⁶ The BioRobotics Institute, Sant'Anna School of Advanced Studies, viale Rinaldo Piaggio 34, 56025 Pisa, Pontedera, Italy



Phenoloxidase activation, antimicrobial, and antibiofilm properties of β -glucan binding protein from *Scylla serrata* crab hemolymph

Mani Divya^a, Baskaralingam Vaseeharan^{a,*}, Mahalingam Anjugam^a, Arokiadhas Iswarya^a, Sivashanmugam Karthikeyan^b, Palaniyandi Velusamy^c, Marimuthu Govindarajan^{d,e}, Naiyf S. Alharbi^f, Shine Kadaikunnan^f, Jamal M. Khaled^f, Csaba Vágvölgyi^{f,g}

^a Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block 6th floor, Burma Colony, Karaikudi 630004, Tamil Nadu, India

^b School of bioscience & Technology, VIT University, Vellore 632014, Tamil Nadu, India

^c Department of Biotechnology, School of Bioengineering, SRM University, Chennai 603203, Tamil Nadu, India

^d Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalainagar 608 002, Tamil Nadu, India

^e Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^f Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^g Department of Microbiology, Faculty of Science and Informatics, University of Szeged, Közép fasor 52, H-6726 Szeged, Hungary

ARTICLE INFO

Article history:

Received 26 February 2018

Received in revised form 24 March 2018

Accepted 26 March 2018

Available online 27 March 2018

Keywords:

Arthropods

β -Glucan binding protein

Crabs

Encapsulation

Live and dead assay

SDS-PAGE

ABSTRACT

In this study, we purified β -GBP from hemolymph of *Scylla serrata* crabs using affinity chromatography. The purified *S. serrata* β -GBP (Ss- β -GBP) had 100 kDa molecular mass in the SDS-PAGE. MALDI-TOF/TOF analysis was conducted, revealing that the purified 100 kDa protein had 96% similarity with β -GBP of *Astacus leptodactylus*. Ss- β -GBP was characterized using high-performance liquid chromatography (HPLC), X-ray diffraction (XRD) analysis, circular dichroism (CD) and Fourier transform infrared (FTIR) spectroscopy, which confirmed the structure of the Ss- β -GBP. The purified Ss- β -GBP was functionally analyzed by yeast agglutination and phagocytic reaction assays. Moreover, the PO enhancing ability of Ss- β -GBP was evidenced through PO activity. Specifically, the antibacterial activity of the Ss- β -GBP against Gram-positive (*Enterococcus faecalis* and *Staphylococcus aureus*) and Gram-negative (*Escherichia coli* and *Pseudomonas aeruginosa*) bacteria was evaluated by determining its minimum inhibitory concentration (MIC) < 60 μ g/ml for all tested species. Furthermore, the antibiofilm efficacy of Ss- β -GBP at 50 and 100 μ g/ml was outlined using light microscopy and confocal laser scanning microscopy (CLSM). Bacterial viability assays also outlined the dose-dependent activity of Ss- β -GBP based on the ratio of live/dead bacterial cells. The results of this study revealed that crab-borne Ss- β -GBP might be widely used to suppress the growth of pathogenic bacteria.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

In response to the growing global population and the projected increasing protein-based food requirements of coming generations, crustaceans have been a focus as a source of animal proteins with several potential applications. For instance, crustacean-based proteins have host defense properties, including prevention of the growth of pathogenic microorganisms [1]. The innate immunity has a critical role in protecting against pathogenic bacteria and fungi, which is mediated by pattern recognition proteins, including Gram-negative bacteria-binding proteins (GNBP) [2,3], peptidoglycan recognition proteins (PGRPs) [4], lipopolysaccharide β -glucan binding proteins (LGBP) [6,7] and β -glucan binding proteins (β -GBP) [8]. β -GBPs are plasma proteins


that precisely bind to β -1, 3-glucan. They are numerous, playing a key role in crustacean effective defense system against invading pathogenic organisms [9]. Furthermore, GNBP led to the production of antibacterial peptides that appear to have an effective role to deactivate pathogenic bacteria [10]. LGBP from *Eriocheir sinensis* show binding activity towards Gram-positive bacteria, while the same is not true for peptidoglycans, because of the complex cell wall composition [11]. The interaction of LGBP with the microbial membrane and the subsequent antimicrobial activity may initiate the immune system mechanisms [12].

Crustacean proteins like lectin and pattern recognition proteins can kill microbial pathogens [13,14]. Furthermore, the antimicrobial activity and diversity of antimicrobial peptides (AMPs) have been investigated as potential and promising drugs for pharmaceutical industries [15].

AMPs are active molecules that play a crucial role by associating with selective cellular membranes to exhibit antimicrobial activity. The structural requirements for AMP activity include α -helical and β -sheet

* Corresponding author.

E-mail address: vaseeharanb@gmail.com. (B. Vaseeharan).



Identification, characterization and immune response of prophenoloxidase from the blue swimmer crab *Portunus pelagicus* and its antibiofilm activity

Sangily Jayanthi^a, Baskaralingam Vaseeharan^{a,*}, Ramachandran Ishwarya^a, Sivashanmugam Karthikeyan^b, Marimuthu Govindarajan^{c,d}, Naiyf S. Alharbi^e, Shine Kadaikunnan^e, Jamal M. Khaled^e, Csaba Vágvolgyi^{e,f}

^a Crustacean Molecular Biology and Genomics Division, Biomaterials and Biotechnology in Animal Health Lab, Department of Animal Health and Management, Alagappa University, Science Block 6th floor, Burma colony, Karaikudi 630004, Tamil Nadu, India

^b School of Bioscience & Technology, VIT University, Vellore 632014, Tamil Nadu, India

^c Unit of Vector Control, Phytochemistry and Nanotechnology, Department of Zoology, Annamalai University, Annamalinagar 608 002, Tamil Nadu, India

^d Department of Zoology, Government College for Women, Kumbakonam 612 001, Tamil Nadu, India

^e Department of Botany and Microbiology, College of Science, King Saud University, Riyadh 11451, Saudi Arabia

^f Department of Microbiology, Faculty of Science and Informatics, University of Szeged, Kozep Fásor 52, H-6726 Szeged, Hungary

ARTICLE INFO

Article history:

Received 16 February 2018

Received in revised form 5 March 2018

Accepted 6 March 2018

Available online 7 March 2018

Keywords:

Antibiofilm activity

Functional analysis

Portunus pelagicus

ABSTRACT

Prophenoloxidase is a conserved Cu-containing enzyme acting as a major defense molecule in the immune response of crustaceans. In the present research, we purified prophenoloxidase from the haemolymph of *Portunus pelagicus* (*Pp*-proPO) by Blue Sepharose CL-6B chromatography. *Pp*-proPO exhibited only one band with molecular weight of 75 kDa on SDS-PAGE. The purified *Pp*-proPO was characterized through X-ray diffraction (XRD) and high-performance liquid chromatography (HPLC). *Pp*-proPO showed phagocytic activity on the yeast *Saccharomyces cerevisiae* as well as encapsulation on sepharose CL-6B beads associated with CM sepharose and beads of sodium alginate. *Pp*-proPO also led to strong agglutination on human erythrocytes. Furthermore, *Pp*-proPO showed magnified PO activity when altered with activated particles acting as pathogen combined molecular patterns (PAMPs), metal ions or other chemicals. *Pp*-proPO showed relevant antibiofilm activity on Gram negative bacteria *Pseudomonas aeruginosa* and *Escherichia coli*. Overall, the above results allowed us to claim that *Pp*-proPO play a key role in immune defense mechanisms of *P. pelagicus* crabs, in particular towards microbial pathogens; notably we added basic information to the functional characterization of *Pp*-proPO, as well as to understand its immunological role in crustaceans defense systems.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction

The prophenoloxidase stimulating system (proPO) plays a crucial role in non-self detection systems characterizing arthropods, and is probably responsible for the non-self-concession method of crustaceans' defenses [1,2]. In invertebrates, proPO plays a main role in the immune recognition processes [3–5]. The proPO acts on cellular reply, with mechanisms comprising phagocytosis, encapsulation, and nodule formation, even as the humoral reaction contains the synthesis of an extensive range of effective antimicrobial proteins [6,7]. This proPO-activating enzyme (PPAE) - previously activated by microbial structure constituents [4]-can modify the inactive enzyme ancestor, proPO, to phenoloxidase (PO), which is the main enzyme concerned with native

immune system of invertebrates [1]. It has been reported for a range of arthropods that the proPO system can influence the pathogen survival [8,9]. Although, in crustaceans proPO is confined to haemocyte granules [10] and its isolation can be done from a haemocyte lysate supernatant [11,12].

Currently, the research on crustacean defense mechanisms is rapidly growing, in order to face a number of pathogens threatening the economic importance of crustacean aquaculture [13]. In this framework, the actuating device of proPO system in crustaceans, activated by constant factors, have been well recorded [4,14]. Indeed, the proPO system acts as a part of coagulation, with opsonic and antimicrobial properties [2], and is supposed to have a key role in host resistance mechanisms [15]. The pro-PO activating system can be physiologically important not only for defense reaction, but also for wound healing purposes. To elucidate the proPO activation process at molecular level, isolation and delineation of proPO, as well as proPO activating components engaged in the related activation process should be carried out.

* Corresponding author.

E-mail address: vaseeharanb@gmail.com (B. Vaseeharan).

PRIVACY PRESERVING ASSOCIATION RULE MINING FROM HIGHLY SECURED OUTSOURCED DATABASES

K. Bhuvaneshwari

Research Scholar, Department of Computer Science,
PRIST University, Thanjavur, India

K. Saravanan

DEAN, Faculty of Computer Science, PRIST University, Thanjavur, India

ABSTRACT

Database outsourcing is becoming more commercial in the recent distributed and parallel systems. This paper considers Association Rule Mining, Frequent Itemset Mining and Privacy Preserving Mining. There is always a controversy between security and the flexibility. For better mining approaches flexibility is more needed from the Database servers but it increases its security risks on distributed network. In current trend of system setup on distributed clouds the Database servers are separated from the service providing web servers. It also extends the resource of the web servers where they can access more than one Database to analyze and retrieve results. At this scenario the web server acts as the intermediate between the Database servers and the client applications. It is responsibility of the web server to preserve privacy of both client and the Database server. This paper concentrates on both client side and Database server side privacy by introducing the algorithms No-Cache Rules Mining on client and Encrypted Database Access on Database server to preserve both client and server privacy. By internally it follows the traditional Association Rule Mining, Frequent Itemset Mining but in different manner.

Key words: Data Mining, Knowledge Discovery, Frequent Itemset Mining, Encrypted Database Access and No-Cache Rules Mining.

Cite this Article: K. Bhuvaneshwari and K. Saravanan, Privacy Preserving Association Rule Mining From Highly Secured Outsourced Databases. *International Journal of Computer Engineering & Technology*, 8(4), 2017, pp. 98–107.

<http://www.iaeme.com/ijcet/issues.asp?JType=IJCET&VType=8&IType=4>

An Improved Naive Bayes Classifier on Imbalanced Attributes

S Geetha, Government Arts College (A), Kumbakonam, India

R Maniyosai, Government Arts College (A), Kumbakonam, India

ABSTRACT

Data plays a major and prominent role in this modern information era. Classification is a data mining task to discover the hidden information from large amounts of data stored in the repository. This process becomes extremely challenging in case of highly imbalanced dataset. Prediction from imbalanced attributes cannot be done accurately in the following case: During the training phase, the categorical variable is not observed but the test phase encounters the categorical variable and hence it assigns zero probability which leads to false prediction. To overcome this scenario, this article proposes a novel smoothing technique called optimized laplace smoothing estimation. This technique adds a bias value function to improve the accuracy of imbalanced attributes. For example, a child dataset has more attributes and the classification model is used to predict the child weight. Some of the attribute values may not be present in the child dataset due to which Naive Bayes assigns a zero for incomplete and an empty attribute. This leads to inaccurate prediction. In such cases, Naive Bayes can be further tuned by adding some new parameters as well as altering the existing optimization method. Experimental analysis shows that this novel smoothing technique enhances the classification accuracy by means of accurate predictions for imbalanced attributes.


KEYWORDS

Child Dataset, Classification, Imbalanced Attributes, Naive bayes

1. INTRODUCTION

Classification is a supervised (Caruana & Niculescu-Mizil, 2006; John & Langley, 1995) learning algorithm where the attribute labels are known. But in some cases (Nguyen et al., 2011), the attribute label is missing or the attribute label is incomplete which leads to an imbalance in the attributes. This dataset imbalance affects the prediction accuracy. Some of the applications which face this attributes imbalance problem are intrusion detection in networks, medical diagnosis, credit card fraud detection, document classification etc (Gonzalez-Abril et al., 2008). The main objective of classification algorithm is that the model constructed using the training data should improve the prediction accuracy. The performance of the classification algorithms achieves better classification accuracy since it does not consider the imbalance between the attributes in the dataset while constructing the classification model.

DOI: 10.4018/IJOI.2019040101



An Improved Naive Bayes Classifier on Imbalanced Attributes

S Geetha (Government Arts College (A), Kumbakonam, India) and R Maniyosai (Government Arts College (A), Kumbakonam, India)

Source Title: International Journal of Organizational and Collective Intelligence (IJOICI) (/journal/international-journal-organizational-collective-intelligence/1140) 9(2)

Copyright: © 2019

Pages: 15

DOI: 10.4018/IJOICI.2019040101

OnDemand:
(Individual Articles)

\$29.50

() Available

[Current Special Offers](#)

Abstract

Data plays a major and prominent role in this modern information era. Classification is a data mining task to discover the hidden information from large amounts of data stored in the repository. This process becomes extremely challenging in case of highly imbalanced dataset. Prediction from imbalanced attributes cannot be done accurately in the following case: During the training phase, the categorical variable is not observed but the test phase encounters the categorical variable and hence it assigns zero probability which leads to false prediction. To overcome this scenario, this article proposes a novel smoothing technique called optimized laplace smoothing estimation. This technique adds a bias value function to improve the accuracy of imbalanced attributes. For example, a child dataset has more attributes and the classification model is used to predict the child weight. Some of the attribute values may not be present in the child dataset due to which Naive Bayes assigns a zero for incomplete and an empty attribute. This leads to inaccurate prediction. In such cases, Naive Bayes can be further tuned by adding some new parameters as well as altering the existing optimization method. Experimental analysis shows that this novel smoothing technique enhances the classification accuracy by means of accurate predictions for imbalanced attributes.

Article Preview

Top

2. Related Works

In this section, various existing works have been discussed.

Complete Article List

Search this Journal:

Reset



ECONOMIC ANALYSIS OF COTTON PRODUCTION AMONG COTTON FARMERS IN NAGAPPATTINAM DISTRICT

R. Sivabalan¹ & Dr. T. Mahenthiran²

¹Research Scholar, Assistant Professor, Department of Economics

Rajah Serfoji Government College (A), Thanjavur

²Assistant Professor and Research Advisor,

Post Graduate and Research Department of Economics,

Rajah Serfoji Government College (Autonomous),

Thanjavur-613 005, Tamil Nadu, India,

Received: 30 Nov 2018

Accepted: 31 Nov 2018

Published: 02 Dec 2018

ABSTRACT

The thrust of this study is on economic analysis of cotton production. The specific objectives were to describe socio-economic characteristics, resource utilization and production technologies and to determine profit in cotton production. Primary data were used for the study through administration of structured questionnaire to collect information from 220 farmers using the list from reconnaissance survey of Tamil Nadu State Agricultural Development Project to randomly select 15% of farmers from selected villages of the four local government areas in the study area. The analytical tools used to achieve the stated objectives in this study were descriptive statistics and net farm income model. The study found that 68% of the farmers were within the average age of 50 years and 30% of the farmers had no formal education while majority of about 74% were married. Also, about 90% were males and 30% having about 15 years of experience in farming. In resource utilization and production technologies, about 35% used seed retained from previous planting while 36% procured fertilizers used from Agricultural Development Projects. Analysis of net farm income showed a profit of 51, 414.51/ha if all labour were valued and the returns per man-day of 966.80 while the returns to investment showed that a farmer gains 1.11 per Naira invested in cotton production. It was recommended that appropriate inputs delivery network need to be put in place by government and agro-service agencies, adequate and intensive research and extension service delivery programme should pursue a consistent and systematic campaign for cotton production while an enabling marketing policy should be instituted by government through product marketing corporation which will serve as a clearing house for cotton marketing in Nagappattinam District.

Key Words: Economic analysis, cotton production, cotton farmers, Nagappattinam District

INTRODUCTION

In Nagappattinam District, the agricultural sector has been invaluable in supporting economic growth and development since independence in 1960. Indeed, before the discovery and exploration of petroleum, the Nagappattinam District economy depended on funds generated from agricultural export expansion for the development of other sectors of the economy. Due to its important role in nation building, the agricultural sector has continued to be a target of government policies overtime (Eyo, 2008).

A Comparative Study of Image Filtering Techniques for Stone Inscriptions

S.Sundareswar

Research Scholar

Department of Computer Science

Research and Development Centre

Bharathiar University

Coimbatore, Tamil Nadu, India.

profresearch2014@gmail.com

Dr.I.Kaspar Raj

Director(i/c),Computer Centre

Department of Computer Science

The Gandigram Rural Institute Deemed University

Gandhigram

Dindigul, Tamil Nadu, India

kasparraj@gmail.com

Abstract – Stone Inscriptions are used to know about the past life of our ancestors and various types of the ancient characters belonging to various centuries. It can be used to establish literacy, archaeological, and historical antiquity on the basis of engravings. Tamil language has the largest number of inscriptions which includes knowledge rich contents. Importance of inscriptions to mankind is remarkable. Nowadays modern readers are faced many difficulties to interpret ancient script from stone inscriptions. Enhancement of inscription image is very important before segmentation process. The need for digitizing this inscriptions arise which help in sharing the content. Enhancement of such inscription image is difficult and still challenging process. The objective of the analysis issues provides to historical document preservation. In this paper discusses the various filtering techniques to enhance the Tamil Brahmi inscription images. The existing filtering techniques are also discussed in this paper to enhance the quality of the inscription images.

Keywords - Tamil Brahmi Inscriptions, Image Quality Factors, Image Noises, Filtering Techniques.

1. INTRODUCTION

In Tamil Nadu, old inscriptions are found in three types of scripts, namely vatteluttu, Tamil and Grantha. The vatteluttu and Tamil Scripts were used to write the Tamil language and the Grantha script was used to write Sanskrit. The Tamil and Grantha scripts are becoming visible only from the 7th century A.D. These inscriptions were engraved on stone and copper plates recording information mostly about constructions of templates, donations to temples etc [1]. In this research work contributes an experiment method

is concerned for enhancement of Tamil Brahmi stone inscriptions using filtering techniques.

This paper is organized as follows: In Section 1 the details of Tamil Brahmi inscription is given. In Section 2 the details of image quality factors are discussed. The types of noise are highlighted in Section 3, Section 4 discusses the various filtering techniques and Section 5 tells about the quality measures. Section 6 describes experimental results and discussion. The proper conclusion is provided in Section 7.

Some of the Tamil Brahmi Stone Inscription images are given below in Figure 1,2,3,4




Fig. 1.



Fig. 2

Secure Association Rule Mining Using Array Mapping Table

K. Bhuvaneshwar

Research Scholar,
Department of Computer Science,
PRIST University,
Thanjavur, India.
bonish88@gmail.com

K. Saravanan

DEAN, Faculty of Computer Science,
PRIST University,
Thanjavur, India.
ks_tnj@yahoo.co.in

Abstract-Database outsourcing is becoming additional commercial approach in the recent distributed and parallel systems. This paper considers Association Rule Mining, Frequent Itemset Mining, and Privacy Protective Mining. There is always a controversy between security, flexibility optimizes space complexity and time complexity. For enhanced mining approaches flexibility is more needed from the Database servers but it increases its security risks on a distributed network. In recent trend of system setup on distributed clouds, the Database servers are separated from the service providing web servers. It also extends the resource of the web servers where they will access more than one Database to analyze and retrieve results. At this state, the web server acts as the midway between the Database servers and the consumer applications. This paper concentrates on both the consumer side and the Database server side privacy by introducing the algorithms No-Cache Rules Mining on the client and Encrypted the Database Access on Database server to preserve both client and server privacy. Primary work to information planning and gathering the required information, optional to create items that decide the rule constraints for information, third mine k-visit itemsets utilizing the new database and fourth deliver the affiliation decide that sets up the learning base and offers better outcomes. Another Technique examined in this paper is the Array Mapping Table used to advance space multifaceted nature and time unpredictability.

Keywords-Data Mining, Association rules, Apriori algorithm, AMT

I. INTRODUCTION

Data mining systems are regularly arranged with regards to the targets they take after and furthermore the outcomes they offer, which acquires PC as an apparatus and makes utilization of the expertise and learning noteworthiness to comprehend and portray the issue. An assortment of information mining strategies, for example, choice trees, affiliation rules, and neural systems are as of now displayed and turn into the purpose of consideration for quite a long

while. Affiliation decide mining procedure is that the best information preparing strategy to go looking covered up or wanted example among the enormous amount of data. It's responsible to initiate connection connections among various information traits in an expansive arrangement of things in a database. A lot of captivating association or associated a case of the affiliation rules mining is the commercial center wicker bin examination. A run of the mill case of the affiliation decides mining is that the market wicker bin examination. Affiliation rules investigation helps to seek out the association among entirely unexpected stock (things) in dealings and order of the client as per purchasing designs. Affiliation Analysis is the recognizable proof of covered illustration or condition that happens a significant part of the time together in a given data. Affiliation Rule mining procedures find captivating affiliations and connections among the known enlightening gathering. An affiliation govern is an administer, which includes certain affiliation relationship with articles or things, for example, the interrelationship of the data thing as whether they happen at the same time with other data thing and how frequently. These tenets are enlisted from the data and, affiliation rules are discovered with help of probability. It has a mentionable measure of reasonable applications, including request, XML mining, spatial data examination, and offer market and proposal systems. These administer measure with help to ensure each dataset treated comparatively in a built up show. The perspective of affiliation govern mining suggests the assistance conviction level chart and thick affiliation control mining to the divulgence of relentless thing sets. Manage support and sureness are two measures of captivating quality. Affiliation rules are seen as drawing in if a base help and a base assurance confine is satisfied. Boolean affiliation run mining is more broadly used than various sorts of affiliation manage mining. Affiliation lead mining system should be possible in four phases. In any case data arranging and pick the required data second make itemsets that choose

An Efficient Lymph Disease Prediction Model Using Naïve Bayes Tree Classifier

¹K. Bhuvaneshwari

¹Assistant Professor, Department of Computer Application, Idhaya College for Women
Kumbakonam

Email: bonish88@gmail.com

Abstract

This research expects to build up a model to upgrade lymphatic diseases analysis by the utilization of irregular forest group machine-learning method prepared with a basic examining plan. This examination has been completed in two noteworthy stages: highlight choice and characterization. In this study, Naïve Bayes tree (NBtree) is employed to classify the data in Lymph disease identification. Exploratory outcomes show that the proposed method accomplishes an amazing improvement in characterization precision rate.

Keywords: Classification, Lymph; NB; Decision trees

1. Introduction

In these days, Computer-Aided Diagnosis (CAD) applications have turned out to be one of the key research themes in medical biometrics diagnostic tasks. Medical diagnosis relies on the experience of the doctor adjacent to the current data. Therefore, various articles recommended a few methodologies to process the doctor's investigation and judgment tasks about genuine clinical evaluations [1]. With sensible achievement, machine-learning systems have been connected in developing the CAD applications because of its solid ability of removing complex connections in the medical data [2]. Crude medical data requires some powerful characterization procedures to help the computer-based examination of such voluminous and heterogeneous data. Precision of clinically analyzed cases is especially critical issue to be considered amid order. Much of the time, the measure of medical datasets is typically incredible, which straightforwardly influences the intricacy of the data mining technique [3]. In this way, the huge scale medical data is viewed as a wellspring of huge difficulties in data mining applications, which includes extricating the most spellbinding or discriminative highlights. In this way, include decrease has a noteworthy job in wiping out unimportant highlights from medical datasets [4], [5]. Dimensionality decrease technique expects to diminish computational unpredictability with the conceivable focal points of upgrading the general arrangement execution. It incorporates dispensing with unimportant

Credit Card Fraud Detection using Feature Augmentation based Boosted Ensemble (FABE)

V. Sobanadevi^{1*}, G. Ravi²

¹ Research Scholar, Department of Computer Science, **Jamal Mohamed College, Trichy, India**

² Associate Professor and Head, Department of Computer Science, **Jamal Mohamed College, Trichy, India**

*Corresponding Author: ammu.soba1987@gmail.com, ravi_govindaraman@yahoo.com

Available online at: www.ijcseonline.org

Received:/2018, Revised:2018, Accepted:2018, Published: 30/Sep/2018

Abstract— Fraud detection in credit card transactions have become mandatory for the financial services industry due to the huge levels of automations observed in the industry. This work presents a Feature Augmentation based Boosted Ensemble (FABE) for credit card fraud detection on huge data. The proposed model integrates two major components; feature augmentation and ensemble creation. Feature augmentation phase performs feature reduction, feature transformation and feature engineering. Feature reduction aids in effective elimination of unnecessary features, while feature transformation and feature engineering aids in creation of new features that can aid in better predictions. The ensemble creation phase models a boosted ensemble using Decision Trees. Multiple training data bags are created, and multiple base learners are created. The learner with highest weight and lowest error levels is iteratively modelled and used as the final learner. Experiments were performed and comparisons with existing models in literature exhibit the high-performance levels of the proposed FABE model.

Keywords— Credit card fraud detection; Ensemble model; Feature Augmentation; Feature Reduction; Feature Engineering; Boosting

I. INTRODUCTION

Big data has become a vital part of the current scenario due to the increase adoption levels of online based systems. Such huge increase in the amount of data poses several unique challenges [1] in fields that require analysis of such huge data. Most real-time data processing applications are currently facing this issue, hence processing Big Data has become a domain in itself. Credit Card Fraud Detection (CCFD) is one of the major use case and has a huge requirement for Big Data processing models that operate in real-time [2]. The huge volume and velocity at which the transactions are generated poses a huge challenge for the prediction models. Significance is to be provided for all transactions irrespective of the transaction value. Both high and low valued transactions are to be provided with the same significance, as fraudsters usually check with the low valued transactions before moving to high valued transactions [3]. Several other issues form major components of credit card transaction data. They are data imbalance and noise [4].

Further, the credit card transactions, being dictated by human behavior, are also affected by concept drift [5].

The problem of anomaly detection in credit card transaction is usually modelled as a supervised learning problem. Machine learning has been the go-to solution for such decision-making problems. The process of classification identifies rules that partition data into sub-sections that involves decisions, which in turn divides them and finally completes by predicting any one of the existing classes [6]. The credit card fraud detection problem can be modelled in several ways even in the supervised context. The two broad categories are models specifically build for the data [7] and models extending generic models [8,9] to perform classification. However, in the current context such simple extensions were found to be insufficient. Currently the data is more complex, as human interaction is the only process that can generate a credit card transaction and human interactions tend to be complex. The generated transactions tend to get more complex, as more customers are involved. Hence simple and single machine learning models are not sufficient

Received: 4 August Revised: 9 August Accepted: 19 August

An economic analysis of women in informal sector - a study in Ariyalur district, Tamil Nadu**Kalaivanan.D**

M.Phil research scholar,

Department of Economics, Annamalai University
Annamalai Nagar, Chidambaram - 608 002**Dr. A. Jayakodi**

Assistant Professor of Economics

Department of Economics, Annamalai University
Annamalai Nagar, Chidambaram - 608 002**Abstract**

The present paper aims at understanding of the status of women in informal sector. In this chapter can well study in three distinct economic sectors. Agriculture, business, and service sector workers, and especially analysis of income, expenditure, savings and debts of women in informal sector in ariyalur district. A small survey has been conducted in this district to find the realities. 90 informal sector women (30 sampling from each sector. Agriculture, Business, Service sector, viz,,)(15 sampling from each block. Ariyalur, Thirumanur, Jayankondam, Andimadam, T.Palur, Sendurai) of ariyalur district were randomly selected with the help of black development officers and village administrations. However it may appear presumptuous on part of the author to claim its generalization for the country, but, a numbers of similar studies conducted in other parts of this district by other scholars support the contention.

Key Words: Working and Living condition, Informal sector, Women, Workers

Introduction

A great majority of the people lives under the poverty line. They are deprived of adequate access to the basic needs of life such as food, dress, shelter, health, education, security, employment, equality and justice. Issues of sustainable livelihood, lack of social and political participation of the vulnerable groups are the major problems in the developing and under developed countries. The government makes some policies for poor people to uplift from the below of poverty line. But, which one of the government has not given sufficient attention to the full implementation of it. So, they are living as third-class people in the country. But unfortunately, among the lower-class and upper-class differences, no one can imagine. But, can well understand the status of women in the society, because, they are only being bottom of the society. So the study deals with the Status of Women in Society. What is Informal Sector? , What they are doing there?, Is it solve their problem?. Come let's see.

Objectives

The following are the objectives of the study:

1. To study the socio-economic status of women in informal sector in the study area.
2. To make a study on the working conditions of women in informal sector.
3. To identify the women workers exploitation and gender discrimination.
4. To offer concrete suggestions for the betterment of women workers and conclusion based on the outcomes of the study.

Methodology

a) Reasons for Choosing the Aria

1. It is well-known fact that Ariyalur district is one of the most backward districts in Tamilnadu.
2. In Ariyalur district, women constitute more than half work in informal sector.
3. In 2011, woman population is higher than the men population. Sex ratio is 1015 females for every 1000 males.
4. In 2011-12 ariyalur district per capita income (Rs.19467). This onlylowest district in per capita income compared to other district.
5. In 2011, total workers in Ariyalurdistrict 34.30per cent were engaged in agriculture as agricultural labourers. But more than 80 per cent of the workers depend on informal sector.

A Study on Impact of Goods and Service Tax on Indian Economy

R.Asokan^{1,3}, A.Jayakodi^{2,3}

¹Assistant professor, Department of Economics, Government Arts College for Women, Salem-8.

²Assistant professor, Department of Economics, Annamalai University, Annamalai Nagar-608002.

³Assistant professor, Department of Economics, Government College for women (Autonomous), Kumbakonam.

Abstract- Goods and Services Tax is the largest indirect tax reform of India. Goods and Services Tax is a single tax system on the contributor of goods and services. It is a destination based tax. It has subsumed taxes like Central Excise Law, Service Tax Law, VAT, Entry Tax, Octroi, etc. Goods and Services Tax is guessed to bring together state economies and develop overall economic growth of the country. The implementation of a tax reform like Goods and Services Tax in a varied country like India necessitated a lot of efforts on the state and national level and drew a lot of challenges like the reconciliation of interests of diverse States with that of the Centre. Some of these issues included origin-based versus destination-based taxation, rate structure and compensation, dispute settlement, inclusion of Alcohol and Petroleum products under Goods and Services Tax. However, there is a positive impact of Goods and Services Tax is on the logistics industry in India had to maintain multiple warehouses across states to evade Goods and Services Tax and state entry taxes on inter-state movement. Most of the times, these warehouses were enforced to operate below their capacity, thus increasing their operating costs. With Goods and Services Tax, restrictions on inter-state movement are lessened and the logistics sector can start consolidating warehouses across the country. Diminution in unnecessary logistics costs will increase profits for businesses involved in supply of goods through transportation. Factors like reduced logistics costs, simplified procedures, lower tax burden, removal of cascading effect, etc. make the Goods and Services Tax regime especially beneficial to startups in the country. Threshold of Rs. Twenty lakhs is beneficial for small businesses. Goods and Services Tax also has helped widen the tax base, with the number of registrations crossing ten million. International Monetary Fund has projected that the Goods and Services Tax reform in India is projected to drive the medium term economic growth rate to beyond 8 per cent.

Keywords: Indirect Tax; Goods and Service Tax; Single tax system; Central GST; State GST

I. INTRODUCTION

Goods and Service Tax is the largest tax reform has established on the idea of “one nation, one market, and one tax” is finalized in India. The moment that the Union government was waiting for a decade has finally

arrived. The single biggest indirect tax management has jerked into force, dismantling all the inter-state obstructions with admiration to trade. The Goods and Service Tax squash, with a single stroke, has converted India into an amalgamated market of 1.3 billion nationals. Primarily, the 2.4 trillion dollar economy is attempting to change itself by doing away with the internal tariff obstacles and subsuming Union, state and local taxes into a unified Goods and Service Tax. The thought of introducing Goods and Service Tax was first recommended by the then Union Finance Minister, P. Chidambaram in his Budget for 2006-07. The discussion on Goods and Service Tax took precise pronouncement with the introduction of the Constitutional Bill of 122nd Amendment, 2014. The Bill was passed by the Parliament on 8 August 2016. This was followed by the authorization of the Bill by more than fifteen states. On 12 April 2017, the Union Government passed four Goods and Service Tax bills are listed below.

- a. Central Goods and Services Tax (CGST) Bill
- b. Integrated Goods and Services Tax (IGST) Bill
- c. Union Territory Goods and Services Tax (UTGST) Bill
- d. The Goods and Services Tax (Compensation to States) Bill

In the Indian economy, the service sector contributes to over fifty five per cent. Separate taxation of goods and services is neither viable nor desirable. Goods and Services Tax in India has been launched to reduce the tax burden that's on both companies and consumers. In the earlier system, there were multiple taxes added at each step of the supply chain, without taking credit for taxes paid at previous stages. As a result, the end cost of the product does not clearly show the actual cost of the product and how much tax was applied. The tax structure was multifaceted. Goods and Service Tax incorporated most of the taxes into one single tax, where the customers are benefited. This system provides Input Tax credit remunerated on the purchase of goods and services, which can be equalize with the tax to be remunerated on the supply of goods and services. As an end result, this diminishes the overall expenditure, with the end

IJSRCSAMS

International Journal of Scientific Research and Reviews

A Peer View about Corporate Social Responsibility in India

A. Jayakodi*

Department of Economics, Government College for Women, Kumbakonam, Tamil Nadu

ABSTRACT

In india, when new economic policies were launched in 1991, the largest foreign companies (agricultural allied industries, soft drinks, other service sectors) came into existence, and local authorities had to face a stiff competition with their new rivals. Because of this, companies can only be successful in earning profit, and the interest in social care decrease, furthermore, companies may try to misuse employers and natural resources to maximize its profits, or look forward to any organization that will systematically process it from service loyalty. In any case, the role of human resources and natural resources are remarkable, and the need for the government is to organize them. So, what is the CSR Act 2014 brought about by the government, the present paper discuss about the benefits for this community by CRS.

KEYWORDS: Corporate, Social, Responsibility, Community, Welfare.

***Corresponding author**

Dr . A. Jayakodi

Assistant Professor

Department of Economics

Government College for Women

Kumbakonam, Tamil Nadu

Email: jayakodi1973@gmail.com

Studies on physico-chemical parameters in the Nagappattinam marine water samples, south east coast of Tamil Nadu, India

¹C. Maheswari, ²A. Muruganandam and ³R. Rengarajan

¹Research Scholar, P.G and Research Department of Botany, M.R. Government Arts College, Mannarkudi, Tamilnadu (India)

²Assistant Professor, P.G and Research Department of Botany, M.R. Government Arts College, Mannarkudi, Tamilnadu (India)

³Assistant Professor, Department of Zoology, Govt. Arts College, Ariyalur, Tamilnadu (India)

ARTICLE DETAILS

Article History

Published Online: 10 November 2018

Keywords

Physico-chemical parameters, Monthly variations, Maximum and minimum, Water quality

Corresponding Author

Email: rengarajan250276@gmail.com

ABSTRACT

Monthly fluctuations of physico-chemical characteristics were carried out in Nagappattinam marine water samples, South east coast of Tamil Nadu, India, for a period of twelve months (August 2016 to July 2017). Nine various physico-chemical parameters were analyzed by using standard methods (APHA, 1998). Water temperature varied from 25.65 to 33.40°C, Turbidity was from 23.57 to 45.36 cm, pH ranged from 7.20 to 8.52, dissolved oxygen content varied between 3.87 to 6.17 mg/L, salinity (28.10 to 32.09 mg/L), calcium (33.67 to 69.98 mg/L), phosphate (0.09 to 0.13 mg/L), nitrate (0.14 to 0.38 mg/L) and ammonia (0.11 to 0.56 mg/L) also varied independently. The maximum and minimum were noted during the study period August 2016 to July 2017.

1. Introduction

The marine environment includes the adjacent coastal areas supports productive and protective habitats such as mangroves, coral reefs and sand dunes. The marine environment is facing a number of pressures, arising out of the needs of people, and the multiple uses that coastal and marine areas can be put to. These pressures contribute to the depletion of marine resources and degradation of the marine environment. In the absence of good management, these pressures may result in severe stress (James Balgan Anand and Mary Jelastin Kala, 2015).

The coastal ecosystems provide food and other incomes, also used for waste disposal, recreation and inspiration. Coastal environment is vital for all human activities including industrial growth. Without the coastal environment, success of any community or nation is impossible. The marine environment forms an essential constituent of the global life. In all the countries, the human activities can affect the physical characteristics of the coastal water. The Coastal ecosystem is the vibrant host for fauna and flora and it is the most important resource to provide a good platform for the coastal life. There are various sources which are responsible to change the biodiversity of the coastal ecosystem (Srivastava *et al.*, 2003).

Marine habitat plays a major role in forecasting, localizing, and manipulating the marine resources (Asha and Diwakar, 2007). The coastal biodiversity is most concentrated in the

Rahul Kundu, 2012). Coastal waters are considered to be the pillar and essential of marine life. The study of hydrographic properties of coastal environments is important, because the variations in the instantly influence on the floral and faunal production. To maintain optimum level of water quality parameters is better for the species survival and healthy ecosystem (James Balgan Anand and Mary Jelastin Kala, 2015).

James Balgan Anand *et al.* (2015) were noted the environmental factors of coastal areas are very important, because the variations in the physico-chemical properties, such as temperature, salinity, pH, dissolved oxygen and nutrients will impact the life span of the fauna and flora in the sea. It regulates the species richness in the coastal area. The variations in the nutrients load along the coastal waters is due to the natural weathering, riverine, land and anthropogenic input. However, the natural seasonal changes keep the coastal waters well mixed and aerated, which help to scatter the nutrients useful for the Biota. The data suggests that during the monsoon period, a significant increase of fresh water and land side input into the coastal area and have elevated nutrient concentration compared with other seasons. In the present investigation attempts to provide such vital information for future references. All the physico-chemical parameters were studied from Nagappattinam marine water samples, south east coast of Tamil nadu, India for a period of August 2016 to July 2017.

Monthly fluctuation of phytoplanktons in Vettar estuary and Nagapattinam marine water samples South east coast of Tamil Nadu India

¹C. Maheswari*, ¹A. Muruganandam, and ²R. Rengarajan

1. P.G and Research Department of Botany, M.R. Government Arts College, Mannarkudi, Tamil Nadu, India.

2. Department of Zoology, Govt. Arts College, Ariyalur, Tamil Nadu, India.


Corresponding author: Dr. A. Muruganandam, P.G and Research Department of Botany, M.R. Government Arts College, Mannarkudi, Tamilnadu, India.

ABSTRACT: A study was undertaken to record the monthly fluctuation in phytoplanktons population in Vettary estuary and Nagapattinam marine water samples south east coast of Tamil Nadu, India, for a period of August 2015 to July 2016. Four groups of phytoplankton were recorded in the samples. They are i. Diatoms, ii. Dinoflagellates, iii. Blue greens and iv. Green algae. During the study, a total of 67 species, belong to 38 families and 57 genera were recorded in respective study area. Diatoms were found to be the dominant group with 52 species. Dinoflagellates formed second dominant group with 10 species. Blue greens came next in the order with 3 species followed by green algae with 2 species in Vettar estuary and Nagapattinam marine area. With respect to population density, in station-1 the density varied from 225 to 5,450 cells/L with minimum during July-18 (premonsoon) and maximum during February-18 (post monsoon), in station-2, the population density it fluctuated between 103 and 7,346 cells/L with minimum during December-15 (monsoon) and maximum during February-16 (summer), in station-3, fluctuated from 102 to 7,961 cells/L with minimum during April-15 (summer) and maximum during May-16 (summer) and in station-4, density of phytoplankton varied from 525 to 7,240 cells/L with minimum during March-16 (post monsoon) and maximum during January-16.

Index Terms: *Phytoplanktons, Diversity, Marine, Estuary, Water samples.*


1. INTRODUCTION

Phytoplankton plays a vital role in the transfer of energy from primary to secondary level; hence, it acts as an integral part of marine and estuarine ecosystem¹. Phytoplankton is a pre dominant type of plants found in aquatic system and its community and relative abundance undergoes continuous Changes at varying scales, and also used as good indicator of water Quality². Phytoplankton species are predominantly



Contents lists available at ScienceDirect

Commun Nonlinear Sci Numer Simulat

journal homepage: www.elsevier.com/locate/cnsns

Research paper

New classes of solutions in the coupled \mathcal{PT} symmetric nonlocal nonlinear Schrödinger equations with four wave mixing

P.S. Vinayagam^a, R. Radha^b, U. Al Khawaja^{a,*}, Liming Ling^c^a Department of Physics, United Arab Emirates University, Al-Ain P.O.Box 15551, United Arab Emirates^b Centre for Nonlinear Science (CeNSc), PG and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam, 612001, India^c Department of Mathematics, South China University of Technology, Guangzhou, 510640, China

ARTICLE INFO

Article history:

Received 27 March 2017

Revised 31 July 2017

Accepted 23 November 2017

Available online 24 November 2017

MSC:

37K40

35Q51

35Q55

Keywords:

Coupled nonlinear Schrödinger system

Soliton

Breathers

Darboux transformation

Lax pair

Four-wave mixing

ABSTRACT

We investigate generalized nonlocal coupled nonlinear Schrödinger equation containing Self-Phase Modulation, Cross-Phase Modulation and four wave mixing involving nonlocal interaction. By means of Darboux transformation we obtained a family of exact breathers and solitons including the Peregrine soliton, Kuznetsov-Ma breather, Akhmediev breather along with all kinds of soliton-soliton and breather-soliton interactions. We analyze and emphasize the impact of the four-wave mixing on the nature and interaction of the solutions. We found that the presence of four wave mixing converts a two-soliton solution into an Akhmediev breather. In particular, the inclusion of four wave mixing results in the generation of a new solutions which is spatially and temporally periodic called “Soliton (Breather) lattice”.

© 2017 Elsevier B.V. All rights reserved.

1. Introduction

Since the invention of the laser, optical solitons [1] have played an important role in nonlinear physics. The optical soliton in fibres is probably the best studied form of solitons because of its remarkable behavior that agrees well with theoretical predictions and its potential as optical information carrier. The propagation of optical pulses through optical birefringent fibres is described by the celebrated Manakov model of the following form [2],

$$i q_{1t} + q_{1xx} + 2(g_{11}|q_1|^2 + g_{12}|q_2|^2)q_1 = 0, \quad (1a)$$

$$i q_{2t} + q_{2xx} + 2(g_{21}|q_1|^2 + g_{22}|q_2|^2)q_2 = 0, \quad (1b)$$

where, q_1 and q_2 are wave envelopes, x , t are space and time variables and i is the imaginary unit. The interaction coefficients g_{11} and g_{22} correspond to the Self-Phase Modulation (SPM) and g_{12} and g_{21} represent the Cross-Phase Modulation

* Corresponding author.

E-mail address: u.alkhawaja@uaeu.ac.ae (U. Al Khawaja).

R. Radha* and C. Senthil Kumar

Digging into the Elusive Localised Solutions of (2+1) Dimensional sine-Gordon Equation

<https://doi.org/10.1515/zna-2017-0447>

Received December 10, 2017; accepted February 19, 2018; previously published online April 6, 2018

Abstract: In this paper, we revisit the (2+1) dimensional sine-Gordon equation analysed earlier [R. Radha and M. Lakshmanan, J. Phys. A Math. Gen. **29**, 1551 (1996)] employing the Truncated Painlevé Approach. We then generate the solutions in terms of lower dimensional arbitrary functions of space and time. By suitably harnessing the arbitrary functions present in the closed form of the solution, we have constructed dromion solutions and studied their collisional dynamics. We have also constructed dromion pairs and shown that the dynamics of the dromion pairs can be turned ON or OFF desirably. In addition, we have also shown that the orientation of the dromion pairs can be changed. Apart from the above classes of solutions, we have also generated compactons, rogue waves and lumps and studied their dynamics.

Keywords: Compactons; Dromions; Lumps; Rogue waves; Truncated Painlevé Approach.

1 Introduction

The advent of localised solutions in terms of doubly periodic Jacobian elliptic functions [1–3] using the Painlevé Truncation Approach has completely revived the interest in the study of (2+1) dimensional integrable models and has given a filip to the identification of more general localised structures. The fact that the exponentially localised solutions called ‘dromions’ generated by Boiti et al. [4] fits into this category of solutions only as a special case has given a new dimension to the investigation of (2+1) dimensional integrable models in an effort to generate other elusive localised solutions like rogue waves, lumps etc. Rogue waves which are another interesting class of solutions, finds application in various fields [5, 6] such

as hydrodynamics [7–10], nonlinear optics [11–18], Bose Einstein condensates [19, 20], plasma physics [21]. The important feature of rogue waves is that they come from nowhere and disappear with no trace. Lumps [22] which are algebraically localised solutions do not interact with each other. In this context, it would be interesting to revisit the (2+1) sine-Gordon equation in an attempt to extract such localised solutions in it.

2 (2+1) Dimensional sine-Gordon Equation

Konopelchenko and Rogers [23, 24] have proposed an interesting symmetric generalisation of the sine-Gordon equation to (2+1) dimensions through a reinterpretation and generalisation of a class of infinitesimal Bäcklund transformations originally introduced in gas dynamics by Loewner [25] as far back in 1952 to give the system of equations

$$\left[\frac{\phi_\xi}{\sin\theta} \right]_\xi - \left[\frac{\phi_\eta}{\sin\theta} \right]_\eta + \frac{(\phi_\eta \theta_\xi - \phi_\xi \theta_\eta)}{\sin^2\theta} = 0, \quad (1)$$

$$\left[\frac{\phi'_\xi}{\sin\theta} \right]_\xi - \left[\frac{\phi'_\eta}{\sin\theta} \right]_\eta + \frac{(\phi'_\xi \theta_\eta - \phi'_\eta \theta_\xi)}{\sin^2\theta} = 0, \quad (2)$$

where $\theta_t = \phi + \phi'$. If we assume that $\phi' = 0$ and that $\theta_t = \phi$ is independent of y , then (1, 2) becomes trivial and (1) gives the (1+1) dimensional sine-Gordon equation

$$\theta_{\xi t} = \sin\theta. \quad (3)$$

Eventhough the (2+1) dimensional sine-Gordon equation has more representations, a more convenient and elegant representation is given by

$$\theta_{xyt} + \frac{1}{2}\rho_x \theta_y + \frac{1}{2}\rho_y \theta_x = 0, \quad (4)$$

$$\rho_{xy} = \frac{1}{2}(\theta_x \theta_y)_t, \quad (5)$$

where

$$\rho_x = -\frac{[\phi_y - \phi'_y] + \theta_{yt} \cos\theta}{\sin\theta}, \rho_y = \frac{[\phi_x - \phi'_x] - \theta_{xt} \cos\theta}{\sin\theta}, \quad (6)$$


*Corresponding author: R. Radha, Centre for Nonlinear Science (CeNSc), Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, India, E-mail: vittal.cnls@gmail.com

C. Senthil Kumar: Department of Physics, VMKV Engineering College, Salem 636308, India



Contents lists available at ScienceDirect

Computers and Mathematics with Applications

journal homepage: www.elsevier.com/locate/camwa

Drone like dynamics of dromion pairs in the (2+1) AKNS equation



R. Radha^{a,*}, C. Senthil Kumar^b, K. Subramanian^{c,d}, T. Alagesan^c

^a Centre for Nonlinear Science (CeNSc), Post-Graduate and Research Department of Physics, Government College for Women (Autonomous), Kumbakonam 612001, India

^b Department of Physics, VMKV Engineering College, Salem 636308, India

^c Department of Physics, Presidency College (Autonomous), Chennai 600005, India

^d Department of Physics, Jerusalem College of Engineering, Chennai 600100, India

ARTICLE INFO

Article history:

Received 16 April 2017

Received in revised form 13 October 2017

Accepted 16 December 2017

Available online 3 January 2018

Keywords:

Truncated Painlevé approach

Drones

Dromion triplet pairs

Rogue waves

Lumps

ABSTRACT

We employ Truncated Painlevé Approach (TPA) to the (2+1) dimensional AKNS equation and construct the solutions in closed form in terms of lower dimensional arbitrary functions of space and time. The highlight of our investigation is that we are able to generate dromions undergoing inelastic and elastic collisions. We observe that the conventional dromions undergo inelastic collision not only exchanging their energy, but also their phase while the dromion pair undergoes elastic collision. In particular, we observe that, we are able to turn ON or OFF the dynamic property of dromion pair by selectively choosing the lower dimensional arbitrary functions with a suitable initial condition. Similar to “drones”, Unmanned Aerial Vehicles (UAVs), dromion pairs can be driven anywhere in the two dimensional plane by selectively giving the initial conditions. In addition to dromions, we have also generated a wide class of localized solutions such as rogue waves and lumps. We observe that while the rogue waves are found to be unstable and stationary, lumps do not interact with other, when they travel in the two dimensional plane.

© 2017 Published by Elsevier Ltd.

1. Introduction

Rogue waves, also known as Freak waves occur in deep ocean [1–4]. The important feature of rogue waves is that they “come from nowhere and disappear with no trace” i.e. they appear for a short duration of time, like a shock wave. Recently, Rogue waves have received much attention in many fields such as hydrodynamics [5], Nonlinear optics [6–8], Bose Einstein Condensates [9,10], Plasma Physics [11], etc. In contrast to the Rogue waves, there is another interesting class of solutions which occur in (2+1) dimensional integrable systems, known as dromions [12–18]. They originate at the cross point of the intersection of two nonparallel ghost solitons, decay exponentially in all directions and are driven by lower dimensional boundaries or velocity potentials. There exists another class of localized solutions called lumps [19,20] which decay algebraically and do not interact with each other. The quest towards unearthing localized solutions in (2+1) dimensional nonlinear partial differential equations (pdes) with exotic behaviour continues even today.

In this paper, we consider the (2+1) dimensional AKNS equation [21] and employ Truncated Painlevé Approach (TPA) [22–26] to construct its solutions in closed form in terms of lower dimensional arbitrary functions of space and time. For a similar (2+1) dimensional Nonlinear equation [27], Lax pair is obtained using Singular Manifold method and Solitonic solutions such as Line solitons and dromions are constructed using Darboux Transformation. Even though the (2+1)

* Corresponding author.

E-mail address: vittal.cnls@gmail.com (R. Radha).



P-ISSN: 2394-1685
E-ISSN: 2394-1693
Impact Factor (ISRA): 5.38
IJPEHS 2018; 5(6): 01-02
©2018 IJPESH
www.kheljournal.com
Received: 01-09-2018
Accepted: 02-10-2018

Dr. R Vidhya Sree
Director of Physical Education,
Govt. College for Women,
Kumbakonam, Tamil Nadu,
India

International Journal of Physical Education, Sports and Health

“8” shaped walk maintain good health

Dr. R Vidhya Sree

Abstract

We all know that Yoga is a great technique used to do breath control and adoption of specific bodily postures. Yoga is widely practiced for health and body relaxation. Similarly Healthy 8 is a walking technique introduced by the Siddhas in India a few thousand years ago. Every one knows that regular walking is a great way to keep one's health fit but walking "8" shaped will give you tremendous health benefits. This is considered to be a supreme walk. Large amount of Oxygen will be inhaled during this walk. Phelgm in the lungs will come out slowly. Body gets energized due to huge quantity of oxygen consumption. One will feel energized the whole day. As per the guidance given by the siddhas, 6 to 8 feet wide and 12 to 16 feet long. Directions North "N" to South "S". By doing this "S" will be aligned to South direction. Start walking from No. 1 and walk towards 2, 3, 4, 5, 6 and come back to number 1. Basically complete the "8" shaped walk. Continuously walk 8 for 10-15 min. After completing 15 min, walk towards 1-6-5-4-3-2-1 for another 10-15 min. Walk at your normal speed. Walking every day can greatly improve your health, help you lose weight, and get you feeling more confident about yourself and your body. It is an essential part of your physical and mental health. Daily walking in 8 will make you feel better and become more productive and happy.


Keywords: Yogis, infinity walk, proper breathing, bare foot, acupressure points, good health

Introduction

Walking is one of the best exercise and will maintain good health. We should walk with free mind without any distraction. If it is done properly excellent health will be maintained. In this modern world Morning walk is becoming fashion and many of the people walk chatting with friends in person or through mobile. Walking with talking.

"Walking" is a common recommendation by any health care professional as one of the best exercise that can help us to maintain good physical and mental health. But in our busy daily schedules, many of us have an excuse to skip this walking. However, we try to compensate by doing a heavy walking or other exercise during weekend!. It may be a justification from our point of view, but the body still needs a daily exercise routine to maintain a proper health. So what's the alternate? Good news is, there is an alternate solution available from our ancient practice called "8 Walking" invented by Tamil Siddars (Yogis). Western world call it as "Infinity walking". Among the walking exercise best walking exercise is "8 shape walk method". It is supreme of all. This is one of the best methods which give miraculous benefits as suggested by the Yogis and Siddhars. It should be practiced daily for 15-30 minutes.

Only at the time of driving license we think about 8 and drive the vehicle in the 8 shape circle and get license. Driving in the 8 shape line by vehicle we get the driving license. But if we walk in the 8 shape line we will get the life long license for good health. We will see the method of 8 walk, how it should be done properly, and what are all the benefits.



Correspondence
Dr. R Vidhya Sree
Director of Physical Education,
Govt. College for Women,
Kumbakonam, Tamil Nadu,
India



Sign in to edit and save changes to this file. ✓

SYNTHESIS OF BIODIESEL FROM NEEM OIL VIA TRANSESTERIFICATION PROCESS USING MG DOPED TITANIUM DI OXIDE

P.Srimathi pasupathy¹ R.Manivannan Rajamanickam²

1. Research Scholar, Assistant Professor of Chemistry,

Department of Chemistry,

Government Arts College (Autonomous), Kumbakonam, Tamil Nadu

Abstract

The Mg doped TiO₂ is synthesized by sol-gel method. The prepared solid nanocatalyst was characterized by XRD, FT-IR. The catalyst showed the highest yield of 89% for transesterification reaction at 65°C, 6 hour with 150mg catalysts loading. The reusability of the catalyst was moderate and future studies focusing on regeneration and enhancing the reusability of the catalyst are recommended.

Introduction

Biodiesel fuel is well known as a biodegradable, renewable, nontoxic, and eco-friendly that can substitute petro-diesel perfectly (Meng et al. 2013) which can be used directly or diesel mixture in engine with little changes (Dehkordi and Ghazem et al. 2012). The majority of world's energy is supplied through petrochemical sources, coal and natural gases. New renewable sources have been widely studied due to the reduction of petroleum reserves and increase in environmental and health concerns [1]. The demand of petroleum-based fuels also increases with increase in the industrialization and motorization of the world. So there is an alternate solution is need to alleviate all the problems. Biodiesel is the opt way to rectify the fuel demand and the environmental issues produced by petroleum productions, bio-diesel is also reduce the dependency on natural resources and greenhouse gas emissions [2]. Conversion of crude oil, coal and natural gas to fuels and chemical feed stocks, production of a variety of petrochemical and chemical products and emission control of CO, hydrocarbons, and NO_x are rely on catalytic technologies [3]. The application of nanotechnology in the domain of harmful dye degradation, pesticide degradation, Water purification, Biodiesel synthesis, in drug synthesis are highly markable. The particle size of the catalyst is one of the most important factors for their catalytic activity [4]. Various Nano catalysts have been investigated for biodiesel production such as Wen et al Catalytic technologies are critical to present and future energy, chemical process and environmental industries. [5-7] The main objective of this work is to synthesize the maximum amount of biodiesel in the presence of Mg doped TiO₂ nanocatalyst, here the efficacy of reusability of nanocatalyst also analysed.

2.Methods

2.1 Materials and Methods

Refined Neem oil (NO) was purchased from local market. Methanol (>99% purity) and all other materials were of analytical grade purchased from E-Merck, India and were used as received without purification. Mg doped TiO₂ was prepared by sol-gel method. Titanium tetra n-butoxide [ti(O-Bu)₄] Magnesium nitrate obtained from E-merck AR grade, were used as titanium and magnesium sources for preparing Pure TiO₂ (anatase form) and Mg²⁺ doped TiO₂ photocatalysts. All other chemicals and reagents are of merck (India) Analytical grade.

Initially 21ml of Titanium tetra-n-butoxide [ti(O-Bu)₄] was dissolved in 80 ml of absolute ethanol (100%) and the resulting solution was stirred vigorously. Then 2ml of water and 0.5 ml of acetic acid 50% were added to another 80ml of ethanol to make ethanol-water-acetic acid solution. The later solution was slowly added to the ti(O-Bu)₄ ethanol solution under vigorous stirring. When the resulting mixture turned to sol, the magnesium nitrate solution was added drop-wise. The resulting transparent colloidal suspension was stirred for more than 2 hours and aged at 25°C until the formation of gel. The gel was dried at 70°C in Vacco pressure of 600psi and then ground. The resulting powder was calcined at 400°C for 2 hours in 50°C increments for every 30 minutes until reaching 400°C after calcination period, the furnace was allowed to cool for 2 hours. The calcined powders were crushed using pestle and mortar under similar conditions in attempt to produce powders with similar fineness and particle size distribution.

2.2 Transesterification of Neem oil

The production of biodiesel from ethanol and Neem oil has been used in the transesterification process. Mg doped TiO₂ have been selected as catalyst. First 20 ml of ethanol was mixed with Mg doped TiO₂ with different ratio (0.50mg - 300mg) This is an exothermic reaction so the mixture would get hot. Then the mixture was preheated about 65°C. Then the mixture was shaken

for 5 minute in a glass beaker. After completion of the methanolysis reaction (6 hour), the catalyst was recovered from the reaction mixture by centrifugation and excess methanol was recover in distillation unit. When the product was cooled, an oil phase clearly separated from a glycerol phase. The top oil phase i.e. biodiesel was collected. The percentage of biodiesel yield was defined as

Yield of Biodiesel= volume of Biodiesel (V₁)

Volume of Neem oil

1 of 4

2.3 Characterization of Catalyst

The crystallinity was determined by XRD using Bruker D8 advance X ray diffractometer equipped with Cu ka



Sign in to edit and save changes to this file.



PHOTO-TRIGGERED DEGRADATION OF RHODAMINE B DYE IN AQUEOUS SOLUTION UNDER SUNLIGHT USING SYNTHESIZED CaO NANOPARTICLES FROM WASTE CHICKEN EGG SHELLS

K.SANTHANALAKSHMI¹, B.AKILANDESHWARI², P.SRIMATHI PASUPATHY³.

1. Research Scholar, 2. Research Scholar, 3. Lecturer in Chemistry,

Department of Chemistry,

Government College for Women (Autonomous), Kumbakonam, Tamil Nadu

ABSTRACT

All over the world pollution is the major problem in Nowadays. Next generation people are must concentrate on environment to lead a life in pollution free world. Dyes are the major polluting agent of water body so. In the present study, CaO photocatalyst was synthesized from waste chicken egg shells by physical method and it was treated with Rhodamine B under sunlight and results shows the optimum dye concentration of Rhodamine B is 10ppm at optimum catalyst 0.5 mg, optimum calcination temperature is 500°C, at 90 minutes at 200 rpm is shows better results.

Key words: Dyes, Degradation, Rhodamine B, Photocatalyst, calcination

INTRODUCTION

The dye is generally applied in an aqueous solution, and may require a mordant to improve the fastness of the dye on the fiber [1]. The dye manufacturing industry represents a relatively small part of the overall chemical industries. Generally these dyes are playing a vital role in polluting rivers, and fresh water bodies. The production of dyes is nearly 800,000 tons per year in all over the world. About 10-15% of synthetic dyes are lost during different processes of textile industry. Sewage is another name of waste water from domestic and industrial processes. Agricultural pollution, Oil pollution, Radioactive substance, river dumping and Marine dumping are the main sources of water pollution. Water pollution statistics 2018-19 government organization announced 663,000,000 peoples are consuming Without Clean Drinking water. [2]. There are so many dyes which is used in textile industries are highly risk and it makes food track infection, Wheezing, and many other diseases. Among the textile dyes Rhodamine B is water-soluble, yielding a red colloidal solution; its solubility is greater in organic solvents. However, the use of Rhodamine B has long been abandoned, primarily because of its carcinogenic properties.[4]. The untreated waste water from textiles draining into the fresh water bodies bringout serious pollution and threatening of environmental and human health.[5-10], so we are in hectic condition which remove the polluting dyes without distributing the economical wealth of our country. Eventhough many types of removal methods are available like coagulation, Flocculation, adsorption, ion exchange method, Nanochemistry leads a very good progress in environmental health. Using nano particles as a catalyst is now very popular method. In this project CaO nanoparticles are used as a photo nanocatalyst to remove Rhodamine B.

MATERIALS AND METHODS

Synthesis of CaO from nanocatalyst:

CaO was synthesized from chicken eggshells by physical method. Collected eggshells were washed dried it in open air for 48 hrs. After that the eggshell was crushed using pestle & mortar by physical method, for 30 minutes. Crushed powder was formed and this powder was treated at above 700° C for 7 hrs. The gaseous state CO₂ was evaporated and form pure CaO Nanoparticles.

EXPERIMENT - I

The Photocatalytic experiments were carried out in the presence of direct sunlight. The experiments were carried out between 10.00am-2.00pm. In all photocatalytic experiments 100ml of 10ppm Rhodamine B was taken in 100ml beaker. The UV - Visible spectrophotometer was used for the determination of absorbance in the range of 560 nm. A known concentration of CaO was added into a beaker, containing Rhodamine B, and kept in the direct sunlight, for Photocatalytic activity.

EXPERIMENT - II

The Nanoparticle with high Photocatalytic activity was selected for further studies. Photocatalytic suspension from 0.5 - 2.5g of CaO. Before irradiation, Photocatalyst suspension was stirred in the dark to ensure the adsorption equilibrium and it was kept in sunlight for the Photocatalytic degradation upto 240 minutes, the suspension was sampled at regular time intervals of 30 minutes and centrifuged for 5 minutes to remove Photocatalyst. Absorbance was monitored by UV - Visible spectrophotometer at 560nm. The data obtained from the Photocatalytic degradation experiments were used to calculate the degradation efficiency:

$$D = \frac{A_0 - A_t}{A_0} \times 100$$

A_t = Adsorbance at time

Derivations on GK algebra

R.Gowri¹, J.kavitha²

¹Assistant Professor, Department of Mathematic, Government college for women (Autonomous), Kumbakonam, India.

²Assistant Professor, Department of Mathematics, D.G.Vaishnav college(Autonomous), Chennai, India

Abstract: In this paper, we introduce the concept of derivation of GK algebra and also obtain some properties about this concept.

I. INTRODUCTION

The structure of GK algebra was introduced by us in 2018[4]. A GK algebra is an algebra of non-empty set X together with a binary operation $*$ and a constant 1 , satisfying the following axioms

- (i) $x*x=1$
- (ii) $x*1=x$
- (iii) $x*y=1$ and $y*x=1$ implies $x=y$
- (iv) $(y*z)*(x*z)=y*x$
- (v) $(x*y)*(1*y)=x$ for all x,y,z in X .

Let the commutative ring R with identity. Let X be an algebra over R . An R linear mapping $f:X \rightarrow X$ is called a derivation if $f(xy)=f(x)y+xf(y)$ for all x,y in X . The derivation of ring is wide area, many researchers started their work in this concept[1],[2],[3]. Inspired by these works, in a same way, we introduce the concept of derivation on GK algebra and discuss some properties in this paper.

II. PRELIMINARIES

A. Definition:2.1[4]

A non-empty set X with fixed constant 1 and a binary operation $*$ is called GK algebra if it satisfying the following axioms

- (i) $x*x=1$
- (ii) $x*1=x$
- (iii) $x*y=1$ and $y*x=1$ implies $x=y$
- (iv) $(y*z)*(x*z)=y*x$
- (v) $(x*y)*(1*y)=x$ for all x,y,z in X .

B. Definition:2.2[4]

GK algebra X is said to be commutative if it satisfies for all x,y in X , $(x*y)*y=(y*x)*x$.

C. Definition:2.3[3]

Let X be a d-algebra. A map $\theta:X \rightarrow X$ is a left-right derivation (l,r)- derivation of X it satisfies the identity $\theta(x*y)=(\theta(x)*y) \wedge (x*\theta(y))$ for all x,y in X . If θ satisfies the identity $\theta(x*y)=(x*\theta(y)) \wedge (\theta(x)*y)$ for all x,y in X then θ is a right-left derivation (r,l) derivation of X .

D. Definition:2.4[2]

Let $(X,*,0)$ be a TM algebra. A self map $d:X \rightarrow X$ is said to be a (l,r) derivation of X if $d(x*y)=(d(x)*y) \wedge (x*d(y))$. A self map $d:X \rightarrow X$ is said to be (r,l) derivation on X if $d(x*y)=(x*d(y)) \wedge (d(x)*y)$.

A NEW FLAVONOID ISOLATED FROM *THEVETIA PERUVIANA* AND EVALUATION OF ANTI-BACTERIAL ACTIVITY OF QUERCETIN- 3-O-RUTINOSIDE

V. Anu,¹S. Ilayaraja,¹A. John Merina

¹Research Scholar, ²Assistant Professor, ³Principal (Retd)

¹PG & Research Department of Chemistry, Government College for Women (Autonomous), Kumbakonam, Tamilnadu-612 001

India

²PG & Research Department of Chemistry, Government Arts College (Autonomous), Kumbakonam, Tamilnadu-612 002, India.

Abstract : The isolation of Quercetin-3-O-rutinoside flavonoid was obtained from *Thevetia peruviana* and the structure was established based on analysis of UV, ¹H NMR and ¹³C NMR spectroscopy methods. Moreover, the absence of toxicity of plant extracts and the isolation of active compounds are important to propose these plants as alternative approaches to resistance management and the results indicate the potential utility of isolated compounds, in the treatment of various bacterial infections.

IndexTerms - *Thevetia peruviana*, Apocynaceae, Flavonoid, Quercetin-3-O-rutinoside.

I. INTRODUCTION

Phytochemicals derived from plants have delivered the basis for numerous commercial prescriptions used today for the treatment of a wide range of diseases. Phytochemical technique mainly applies to the quality control of herbal medicine of different chemical constituents such as tannins, saponins, alkaloids, volatile oils, flavonoids and anthraquinones. It is essential to use the phytochemical methods to screen and analyze bioactive components, not only for the quality control of crude drugs, but also for the interpretation of their therapeutic mechanisms.¹ The study of the distribution of flavonoids in plants is a continuing exercise and known flavonoids are being regularly exposed from new sources.²

Thevetia peruviana belongs to the family Apocynaceae and it often known as yellow oleander.³ *Thevetia peruviana* has been deliberated as a hypothetically important plant for industrial and pharmacological application. Research admired that phytochemicals, working together with nutrients found in fruits, vegetables and nuts, may help slow the aging process and reduce the risk of numerous diseases such as high blood pressure, pain, asthma, cancers, heart diseases and urinary tract infections.⁴

The ethno-medical uses of *Thevetia peruviana* is seeming in treating the external wounds, infected area, ring worms, tumours etc., the use of grinded leaves of *Thevetia peruviana* in ethno-veterinary medicine is the evidence for its plentiful use for healing of wounds.⁵ In the present study, yellow flowers of *Thevetia peruviana* were selected and the flavonol glycoside has been isolated and its therapeutic applications have been investigated.

II. MATERIALS AND METHODS

2.1. Extraction and fractionation

The fresh flowers (2 kg) of *Thevetia peruviana* (Apocynaceae) collected from Kumbakonam were extracted with 85 % MeOH (5 X 500 mL) under reflux. The alcoholic extract was concentrated *in vacuo* and the aqueous concentrate successively fractionated with peroxide - free ether (4 X 250 mL) and ethyl acetate (8 X 250 mL). The EtOAc fraction alone was taken up for the study.

2.2. Ether fraction: flavonol - quercetin

The Et₂O fraction was concentrated *in vacuo* and left in an ice chest for about a week. A yellow solid that separated was filtered and studied. It came out as pale yellow needles m.p. 315-317°C on crystallization from MeOH. It was soluble in organic solvents and sparingly soluble in hot water. It gave a red colour with Mg-HCl, olive green with alc.Fe³⁺, golden yellow colour with NH₃ and NaOH and appeared yellow under UV and UV/NH₃. It answered the Horhammer-Hansal, Wilson's boric acid and Gibb's tests. It had λ_{max}^{MeOH} nm 256, 268 sh, 370; +NaOMe 261 sh, 321, 422; +AlCl₃ 266, 302, 457; + (AlCl₃ - HCl) 268, 303, 352, 427; +NaOAc 274, 327, 390 and + (NaOAc - H₃BO₃) 261, 303 sh, 385 nm. The aglycone part of the compound was identified as quercetin and the same was confirmed by CO, mixed PC and m.m.p. with an authentic sample of quercetin from *Wrightia tinctoria*.⁶

2.3. Ethyl acetate fraction: Quercetin-3-O-rutinoside

The ethyl acetate fraction was concentrated *in vacuo* and left in an ice-chest for few days. A yellow solid that separated was filtered and studied. It developed a green colour with alc. Fe³⁺ and a pink colour with Mg-HCl. It appeared deep purple under UV that turned yellow on exposure to NH₃. It responded to Wilson's boric acid, Molisch and Gibb's tests, but did not answer the