

Date: 4.11.24

Defn:

The real-valued

function  $f$  on the interval $J \subset \mathbb{R}$  is said to be strictly

increasing if

$$f(x) < f(y) \quad (x < y; x, y \in J)$$

Similarly,  $f$  is said to be strictly

decreasing if

$$f(x) > f(y) \quad (x < y; x, y \in J)$$

Thus, if  $f$  is nonincreasing on  $J$ then  $f$  is strictly increasingon  $J$  if and only if  $f$  isstrictly decreasing on  $J$ .Date: 4.2 Metric Space

Refer the book

4.3

Page - (111)

2.2A

Let  $\{S_n\}_{n=1}^{\infty}$  be a sequence ofreal numbers. We say that  $S_n$  approachesthe limit  $L$  (as  $n$  approaches infinity),if for every  $\epsilon > 0$  there is a positiveinteger  $N$  such that

$$|S_n - L| < \epsilon \quad (n \geq N)$$

If  $S_n$  approaches the limit  $L$ we write,  $\lim_{n \rightarrow \infty} S_n = L$ 

$$(a) S_n \rightarrow L \quad (n \rightarrow \infty)$$

2.3 A

Q) of the sequence of real numbers  $\{s_n\}_{n=1}^{\infty}$  has the limit  $L$ ,

We say that  $\{s_n\}_{n=1}^{\infty}$  is convergent to  $L$ .

If  $\{s_n\}_{n=1}^{\infty}$  does not have

a limit, we say that  $\{s_n\}_{n=1}^{\infty}$  is divergent.

Ex: 1) The sequences  $1, 1, 1, \dots$  and  $1, \frac{1}{2}, \frac{1}{3}, \dots$  are convergent.

2) The sequences  $1, 2, 3, \dots$  and  $-1, +1, -1, +1, \dots$  are divergent.

2.10 A Let  $\{s_n\}_{n=1}^{\infty}$  be a sequence

of real numbers. Then  $\{s_n\}_{n=1}^{\infty}$

is called a Cauchy

sequence if for any  $\epsilon > 0$  there exists an  $N \in \mathbb{I}$  such that

$$|s_m - s_n| < \epsilon \quad (m, n \geq N)$$

2.10 B If the sequence of real numbers  $\{s_n\}_{n=1}^{\infty}$  converges,

then  $\{s_n\}_{n=1}^{\infty}$  is a Cauchy sequence.

4.3 C Defn:

4.3 E: 1

Let  $(M, \epsilon)$  be a metric

space. If  $\{s_n\}_{n=1}^{\infty}$  is a convergent

sequence of points of  $M$ ,

then  $\{s_n\}_{n=1}^{\infty}$  is Cauchy.

Date: \_\_\_\_\_  
Proof: Let  $L = \lim_{n \rightarrow \infty} s_n$

Then, Given  $\epsilon > 0$  there exists an  $N \in \mathbb{I}$  such that,

$$\rho(s_{m+1}) < \epsilon/2 \text{ for every } m, \rho(s_n, L) < \epsilon/2$$

Thus, if  $m, n \geq N$ , we have

$$\rho(s_m, s_n) = \rho(s_{m+1}, s_{n+1}) \\ |s_m - s_n| = |s_{m+1} - s_{n+1}|$$

$$\text{i.e., } \rho(s_m, s_n) = \rho(s_{m+1}, s_{n+1}) + \rho(s_n, L) \\ \leq \rho(s_{m+1}, s_{n+1}) + \rho(s_n, L) \\ \leq \epsilon/2 + \epsilon/2$$

Which proves that  $\{s_n\}_{n=1}^{\infty}$  is Cauchy.

5.2 Theorem:

The real-valued function

$f$  is continuous at  $a \in \mathbb{R}^1$

iff the inverse image under

$f$  of any open ball  $B[f(a); \epsilon]$  about  $f(a)$  contains an open

ball  $B[a; \delta]$  about  $a$ .

$$\text{i.e., } f^{-1}[B[f(a); \epsilon]] \supset B[a; \delta].$$

Proof:

Let the real valued function

$f$  is continuous at  $a \in \mathbb{R}^1$ .

Let us consider that,

$$x \in B[a; \delta]$$